

EXERCISE 8.1

Q.1. The angles of a quadrilateral are in the ratio 3 : 5 : 9 : 13. Find all the angles of the quadrilateral.

Sol. Suppose the measures of four angles are $3x$, $5x$, $9x$ and $13x$.

$$\therefore 3x + 5x + 9x + 13x = 360^\circ \quad [\text{Angle sum property of a quadrilateral}]$$

$$\Rightarrow 30x = 360^\circ$$

$$\Rightarrow x = \frac{360^\circ}{30} = 12^\circ$$

$$\Rightarrow 3x = 3 \times 12^\circ = 36^\circ$$

$$5x = 5 \times 12^\circ = 60^\circ$$

$$9x = 9 \times 12^\circ = 108^\circ$$

$$13x = 13 \times 12^\circ = 156^\circ$$

$\therefore$ the angles of the quadrilateral are **36° , 60° , 108° and 156°** Ans.

Q.2. If the diagonals of a parallelogram are equal, then show that it is a rectangle.

Sol. **Given :** ABCD is a parallelogram in which $AC = BD$.

To Prove : ABCD is a rectangle.

Proof : In $\triangle ABC$ and $\triangle BAD$

$$AB = AB \quad [\text{Common}]$$

$$BC = AD \quad [\text{Opposite sides of a parallelogram}]$$

$$AC = BD \quad [\text{Given}]$$

$$\therefore \triangle ABC \cong \triangle BAD \quad [\text{SSS congruence}]$$

$$\angle ABC = \angle BAD \quad \dots(i) \quad [\text{CPCT}]$$

Since, ABCD is a parallelogram, thus,

$$\angle ABC + \angle BAD = 180^\circ \quad \dots(ii)$$

[Consecutive interior angles]

$$\angle ABC + \angle ABC = 180^\circ$$

$$\therefore 2\angle ABC = 180^\circ \quad [\text{From (i) and (ii)}]$$

$$\Rightarrow \angle ABC = \angle BAD = 90^\circ$$


This shows that ABCD is a parallelogram one of whose angle is 90° .

Hence, ABCD is a rectangle. **Proved.**

Q.3. Show that if the diagonals of a quadrilateral bisect each other at right angles, then it is a rhombus.

Sol. **Given :** A quadrilateral ABCD, in which diagonals AC and BD bisect each other at right angles.

To Prove : ABCD is a rhombus.



Proof : In $\triangle AOB$ and $\triangle BOC$

$$AO = OC$$

[Diagonals AC and BD bisect each other]

$$\angle AOB = \angle COB \quad [\text{Each} = 90^\circ]$$

$$BO = BO \quad [\text{Common}]$$

$$\therefore \triangle AOB \cong \triangle BOC \quad [\text{SAS congruence}]$$


$$AB = BC \quad \dots(i) \quad [\text{CPCT}]$$

Since, ABCD is a quadrilateral in which

$$AB = BC \quad [\text{From (i)}]$$

Hence, ABCD is a rhombus.

[$\therefore$ if the diagonals of a quadrilateral bisect each other, then it is a parallelogram and opposite sides of a parallelogram are equal] **Proved.**



Q.4. Show that the diagonals of a square are equal and bisect each other at right angles.

Sol. Given : ABCD is a square in which AC and BD are diagonals.

To Prove : $AC = BD$ and AC bisects BD at right angles, i.e. $AC \perp BD$.

$$AO = OC, OB = OD$$

Proof : In $\triangle ABC$ and $\triangle BAD$,

$$AB = AB \quad [\text{Common}]$$

$$BC = AD \quad [\text{Sides of a square}]$$

$$\angle ABC = \angle BAD = 90^\circ \quad [\text{Angles of a square}]$$

$$\therefore \triangle ABC \cong \triangle BAD \quad [\text{SAS congruence}]$$

$$\Rightarrow AC = BD \quad [\text{CPCT}]$$

Now in $\triangle AOB$ and $\triangle COD$,


$$AB = DC \quad [\text{Sides of a square}]$$

$$\angle AOB = \angle COD \quad [\text{Vertically opposite angles}]$$

$$\angle OAB = \angle OCD \quad [\text{Alternate angles}]$$

$$\therefore \triangle AOB \cong \triangle COD \quad [\text{AAS congruence}]$$

$$\angle AO = \angle OC \quad [\text{CPCT}]$$



Similarly by taking $\triangle AOD$ and $\triangle BOC$, we can show that $OB = OD$.

$$\text{In } \triangle ABC, \angle BAC + \angle BCA = 90^\circ \quad [\because \angle B = 90^\circ]$$

$$\Rightarrow 2\angle BAC = 90^\circ \quad [\angle BAC = \angle BCA, \text{ as } BC = AD]$$

$$\Rightarrow \angle BCA = 45^\circ \text{ or } \angle BCO = 45^\circ$$

$$\text{Similarly } \angle CBO = 45^\circ$$

In $\triangle BCO$.

$$\angle BCO + \angle CBO + \angle BOC = 180^\circ$$

$$\Rightarrow 90^\circ + \angle BOC = 180^\circ$$

$$\Rightarrow \angle BOC = 90^\circ$$


$$\Rightarrow BO \perp OC \Rightarrow BO \perp AC$$

Hence, $AC = BD$, $AC \perp BD$, $AO = OC$ and $OB = OD$. **Proved.**

Q.5. Show that if the diagonals of a quadrilateral are equal and bisect each other at right angles, then it is a square.

Sol. Given : A quadrilateral ABCD, in which diagonals AC and BD are equal and bisect each other at right angles,

To Prove : ABCD is a square.



Proof : Since ABCD is a quadrilateral whose diagonals bisect each other, so it is a parallelogram. Also, its diagonals bisect each other at right angles, therefore, ABCD is a rhombus.

$\Rightarrow AB = BC = CD = DA$ [Sides of a rhombus]

In $\triangle ABC$ and $\triangle BAD$, we have

$AB = AB$ [Common]

$BC = AD$ [Sides of a rhombus]

$AC = BD$ [Given]

$\therefore \triangle ABC \cong \triangle BAD$ [SSS congruence]

$\therefore \angle ABC = \angle BAD$ [CPCT]

But, $\angle ABC + \angle BAD = 180^\circ$ [Consecutive interior angles]

$\angle ABC = \angle BAD = 90^\circ$

$\angle A = \angle B = \angle C = \angle D = 90^\circ$ [Opposite angles of a $\parallel^{\text{gm}}$]

$\Rightarrow$ ABCD is a rhombus whose angles are of 90° each.

Hence, ABCD is a square. **Proved.**

Q.6. Diagonal AC of a parallelogram ABCD bisects $\angle A$ (see Fig.). Show that

(i) it bisects $\angle C$ also,

(ii) ABCD is a rhombus.

Given : A parallelogram ABCD, in which diagonal AC bisects $\angle A$, i.e., $\angle DAC = \angle BAC$.

To Prove : (i) Diagonal AC bisects $\angle C$ i.e., $\angle DCA = \angle BCA$

(ii) ABCD is a rhombus.

Proof : (i) $\angle DAC = \angle BCA$

$\angle BAC = \angle DCA$

But, $\angle DAC = \angle BAC$

$\therefore \angle BCA = \angle DCA$

Hence, AC bisects $\angle DCB$

Or, AC bisects $\angle C$ **Proved.**

(ii) In $\triangle ABC$ and $\triangle CDA$

$AC = AC$ [Common]

$\angle BAC = \angle DAC$ [Given]

and $\angle BCA = \angle DCA$ [Proved above]


$\therefore \triangle ABC \cong \triangle ADC$ [ASA congruence]

$\therefore BC = DC$ [CPCT]

But $AB = DC$ [Given]

$\therefore AB = BC = DC = AD$

Hence, ABCD is a rhombus **Proved.**




[$\therefore$ opposite angles are equal]

Q.7. ABCD is a rhombus. Show that diagonal AC bisects $\angle A$ as well as $\angle C$ and diagonal BD bisects $\angle B$ as well as $\angle D$.

Sol. Given : ABCD is a rhombus, i.e.,

$AB = BC = CD = DA$.

To Prove : $\angle DAC = \angle BAC$,



$$\angle BCA = \angle DCA$$

$$\angle ADB = \angle CDB, \angle ABD = \angle CBD$$

Proof : In $\triangle ABC$ and $\triangle CDA$, we have

$$AB = AD$$

[Sides of a rhombus]

$$AC = AC$$

[Common]

$$BC = CD$$

[Sides of a rhombus]

$$\triangle ABC \cong \triangle ADC$$

[SSS congruence]

$$\text{So, } \left. \begin{array}{l} \angle DAC = \angle BAC \\ \angle BCA = \angle DCA \end{array} \right\} \text{ [CPCT]}$$

Similarly, $\angle ADB = \angle CDB$ and $\angle ABD = \angle CBD$.

Hence, diagonal AC bisects $\angle A$ as well as $\angle C$ and diagonal BD bisects $\angle B$ as well as $\angle D$. **Proved.**

Q.8. *ABCD is a rectangle in which diagonal AC bisects $\angle A$ as well as $\angle C$. Show that :*

(i) ABCD is a square (ii) diagonal BD bisects $\angle B$ as well as $\angle D$.

Sol. Given : ABCD is a rectangle in which diagonal AC bisects $\angle A$ as well as $\angle C$.

To Prove : (i) ABCD is a square.
(ii) Diagonal BD bisects $\angle B$ as well as $\angle D$.

Proof : (i) In $\triangle ABC$ and $\triangle ADC$, we have
 $\angle BAC = \angle DAC$ [Given]
 $\angle BCA = \angle DCA$ [Given]
 $AC = AC$

$$\therefore \triangle ABC \cong \triangle ADC \quad \text{[ASA congruence]}$$

$$\therefore AB = AD \text{ and } CB = CD \quad \text{[CPCT]}$$

But, in a rectangle opposite sides are equal,

$$\text{i.e., } AB = DC \text{ and } BC = AD$$

$$\therefore AB = BC = CD = DA$$

Hence, ABCD is a square **Proved.**

(ii) In $\triangle ABD$ and $\triangle CDB$, we have

$$AD = CD$$

$$AB = CD$$

$$BD = BD$$


[Sides of a square]

[Common]

$$\therefore \triangle ABD \cong \triangle CBD \quad \text{[SSS congruence]}$$

$$\text{So, } \left. \begin{array}{l} \angle ABD = \angle CBD \\ \angle ADB = \angle CDB \end{array} \right\} \text{ [CPCT]}$$

Hence, diagonal BD bisects $\angle B$ as well as $\angle D$ **Proved.**



Q.9. *In parallelogram ABCD, two points P and Q are taken on diagonal BD such that $DP = BQ$ (see Fig.). Show that :*


$$(i) \triangle APD \cong \triangle CQB$$

$$(ii) AP = CQ$$

$$(iii) \triangle AQB \cong \triangle CPD$$

$$(iv) AQ = CP$$

$$(v) APCQ \text{ is a parallelogram}$$



Sol. Given : ABCD is a parallelogram and P and Q are points on diagonal BD such that DP = BQ.

To Prove : (i) $\triangle APD \cong \triangle CQB$

(ii) $AP = CQ$

(iii) $\triangle AQB \cong \triangle CPD$

(iv) $AQ = CP$

(v) APCQ is a parallelogram

Proof : (i) In $\triangle APD$ and $\triangle CQB$, we have

$AD = BC$ [Opposite sides of a || gm]

$DP = BQ$ [Given]

$\angle ADP = \angle CBQ$ [Alternate angles]

$\therefore \triangle APD \cong \triangle CQB$ [SAS congruence]

(ii) $\therefore AP = CQ$ [CPCT]

(iii) In $\triangle AQB$ and $\triangle CPD$, we have

$AB = CD$ [Opposite sides of a || gm]


$DP = BQ$ [Given]

$\angle ABQ = \angle CDP$ [Alternate angles]

$\therefore \triangle AQB \cong \triangle CPD$ [SAS congruence]

(iv) $\therefore AQ = CP$ [CPCT]


(v) Since in APCQ, opposite sides are equal, therefore it is a parallelogram. **Proved.**



Q.10. ABCD is a parallelogram and AP and CQ are perpendiculars from vertices A and C on diagonal BD (see Fig.). Show that

(i) $\triangle APB \cong \triangle CQD$

(ii) $AP = CQ$



Sol. Given : ABCD is a parallelogram and AP and CQ are perpendiculars from vertices A and C on BD.

To Prove : (i) $\triangle APB \cong \triangle CQD$

(ii) $AP = CQ$

Proof : (i) In $\triangle APB$ and $\triangle CQD$, we have


$\angle ABP = \angle CDQ$ [Alternate angles]

$AB = CD$ [Opposite sides of a parallelogram]

$\angle APB = \angle CQD$ [Each = 90°]

$\therefore \triangle APB \cong \triangle CQD$ [ASA congruence]

(ii) So, $AP = CQ$ [CPCT] **Proved.**



Q.11. In $\triangle ABC$ and $\triangle DEF$, $AB = DE$, $AB \parallel DE$, $BC = EF$ and $BC \parallel EF$. Vertices A, B and C are joined to vertices D, E and F respectively (see Fig.). Show that

(i) quadrilateral ABED is a parallelogram


(ii) quadrilateral BEFC is a parallelogram

(iii) $AD \parallel CF$ and $AD = CF$


(iv) quadrilateral ACFD is a parallelogram

(v) $AC = DF$

(vi) $\triangle ABC \cong \triangle DEF$



Sol. Given : In $\triangle ABC$ and $\triangle DEF$, $AB = DE$, $AB \parallel DE$, $BC = EF$ and $BC \parallel EF$. Vertices A, B and C are joined to vertices D, E and F.



- To Prove :**
- (i) ABED is a parallelogram
 - (ii) BEFC is a parallelogram
 - (iii) $AD \parallel CF$ and $AD = CF$
 - (iv) ACFD is a parallelogram
 - (v) $AC = DF$
 - (vi) $\triangle ABC \cong \triangle DEF$


Proof :

- (i) In quadrilateral ABED, we have
 $AB = DE$ and $AB \parallel DE$. [Given]
 $\Rightarrow$ ABED is a parallelogram.
 [One pair of opposite sides is parallel and equal]
- (ii) In quadrilateral BEFC, we have
 $BC = EF$ and $BC \parallel EF$ [Given]
 $\Rightarrow$ BEFC is a parallelogram.
 [One pair of opposite sides is parallel and equal]
- (iii) $BE = CF$ and $BE \parallel CF$ [BEFC is parallelogram]
 $AD = BE$ and $AD \parallel BE$ [ABED is a parallelogram]
 $\Rightarrow AD = CF$ and $AD \parallel CF$
- (iv) ACFD is a parallelogram.
 [One pair of opposite sides is parallel and equal]
- (v) $AC = DF$ [Opposite sides of parallelogram ACFD]
- (vi) In $\triangle ABC$ and $\triangle DEF$, we have
 $AB = DE$ [Given]
 $BC = EF$ [Given]
 $AC = DF$ [Proved above]
 $\therefore \triangle ABC \cong \triangle DEF$ [SSS axiom] **Proved.**

Q.12. ABCD is a trapezium in which $AB \parallel CD$ and $AD = BC$ (see Fig.).

Show that


- (i) $\angle A = \angle B$
- (ii) $\angle C = \angle D$
- (iii) $\triangle ABC \cong \triangle BAD$
- (iv) diagonal $AC =$ diagonal BD



Sol. Given : In trapezium ABCD, $AB \parallel CD$ and $AD = BC$.

- To Prove :**
- (i) $\angle A = \angle B$
 - (ii) $\angle C = \angle D$
 - (iii) $\triangle ABC \cong \triangle BAD$
 - (iv) diagonal $AC =$ diagonal BD

Constructions : Join AC and BD. Extend AB and draw a line through C parallel to DA meeting AB produced at E.




Proof :

- (i) Since $AB \parallel DC$
 $\Rightarrow AE \parallel DC$... (i)
 and $AD \parallel CE$... (ii) [Construction]
 $\Rightarrow ADCE$ is a parallelogram [Opposite pairs of sides are parallel]
 $\angle A + \angle E = 180^\circ$... (iii) [Consecutive interior angles]
 $\angle B + \angle CBE = 180^\circ$... (iv) [Linear pair]
 $AD = CE$... (v) [Opposite sides of a $\parallel^m$]
 $AD = BC$... (vi) [Given]
 $\Rightarrow BC = CE$ [From (v) and (vi)]
 $\Rightarrow \angle E = \angle CBE$... (vii) [Angles opposite to equal sides]
 $\therefore \angle B + \angle E = 180^\circ$... (viii) [From (iv) and (vii)]
 Now from (iii) and (viii) we have
 $\angle A + \angle E = \angle B + \angle E$
 $\Rightarrow \angle A = \angle B$ **Proved.**
- (ii) $\left. \begin{array}{l} \angle A + \angle D = 180^\circ \\ \angle B + \angle C = 180^\circ \end{array} \right\}$ [Consecutive interior angles]
 $\Rightarrow \angle A + \angle D = \angle B + \angle C$ [$\because \angle A = \angle B$]
 $\Rightarrow \angle D = \angle C$
 Or $\angle C = \angle D$ **Proved.**
- (iii) In $\triangle ABC$ and $\triangle BAD$, we have
 $AD = BC$ [Given]
 $\angle A = \angle B$ [Proved]
 $AB = CD$ [Common]
 $\therefore \triangle ABC \cong \triangle BAD$ [ASA congruence]
- (iv) diagonal $AC =$ diagonal BD [CPCT] **Proved.**

EXERCISE 8.2

Q.1. *ABCD is a quadrilateral in which P, Q, R and S are mid-points of the sides AB, BC, CD and DA respectively. (see Fig.). AC is a diagonal. Show that :*




(i) $SR \parallel AC$ and $SR = \frac{1}{2}AC$

(ii) $PQ = SR$

(iii) $PQRS$ is a parallelogram.

Given : ABCD is a quadrilateral in which P, Q, R and S are mid-points of AB, BC, CD and DA. AC is a diagonal.



To Prove : (i) $SR \parallel AC$ and $SR = \frac{1}{2}AC$

(ii) $PQ = SR$

(iii) $PQRS$ is a parallelogram

Proof : (i) In $\triangle ABC$, P is the mid-point of AB and Q is the mid-point of BC.

$$\therefore PQ \parallel AC \text{ and } PQ = \frac{1}{2}AC \quad \dots(1)$$

[Mid-point theorem]

In $\triangle ADC$, R is the mid-point of CD and S is the mid-point of AD

$$\therefore SR \parallel AC \text{ and } SR = \frac{1}{2}AC \quad \dots(2)$$

[Mid-point theorem]

(ii) From (1) and (2), we get

$$PQ \parallel SR \text{ and } PQ = SR$$

(iii) Now in quadrilateral PQRS, its one pair of opposite sides PQ and SR is equal and parallel.

$\therefore PQRS$ is a parallelogram. **Proved.**

Q.2. *ABCD is a rhombus and P, Q, R and S are the mid-points of the sides AB, BC, CD and DA respectively. Show that the quadrilateral PQRS is a rectangle.*

Sol. Given : ABCD is a rhombus in which P, Q, R and S are mid points of sides AB, BC, CD and DA respectively :

To Prove : PQRS is a rectangle.

Construction : Join AC, PR and SQ.

Proof : In $\triangle ABC$

P is mid point of AB [Given]

Q is mid point of BC [Given]

$\Rightarrow PQ \parallel AC$ and $PQ = \frac{1}{2} AC$... (i) [Mid point theorem]

Similarly, in $\triangle DAC$,

$SR \parallel AC$ and $SR = \frac{1}{2} AC$... (ii)

From (i) and (ii), we have $PQ \parallel SR$ and $PQ = SR$

$\Rightarrow$ PQRS is a parallelogram

[One pair of opposite sides is parallel and equal]

Since ABQS is a parallelogram

$\Rightarrow AB = SQ$ [Opposite sides of a $\parallel$ gm]


Similarly, since PBCR is a parallelogram.

$\Rightarrow BC = PR$

Thus, $SQ = PR$ [AB = BC]

Since SQ and PR are diagonals of parallelogram PQRS, which are equal.

$\Rightarrow$ PQRS is a rectangle. **Proved.**




Q.3. *ABCD is a rectangle and P, Q, R and S are mid-points of the sides AB, BC, CD and DA respectively. Show that the quadrilateral PQRS is a rhombus.*

Sol. Given : A rectangle ABCD in which P, Q, R, S are the mid-points of AB, BC, CD and DA respectively, PQ, QR, RS and SP are joined.

To Prove : PQRS is a rhombus.

Construction : Join AC



Proof : In $\triangle ABC$, P and Q are the mid-points of the sides AB and BC.

$$\therefore PQ \parallel AC \text{ and } PQ = \frac{1}{2} AC \quad \dots(i) \quad [\text{Mid point theorem}]$$

Similarly, in $\triangle ADC$,

$$SR \parallel AC \text{ and } SR = \frac{1}{2} AC \quad \dots(ii)$$

From (i) and (ii), we get

$$PQ \parallel SR \text{ and } PQ = SR \quad \dots(iii)$$

Now in quadrilateral PQRS, its one pair of opposite sides PQ and SR is parallel and equal [From (iii)]

$\therefore$ PQRS is a parallelogram.

$$\text{Now } AD = BC \quad \dots(iv)$$

[Opposite sides of a rectangle ABCD]

$$\therefore \frac{1}{2} AD = \frac{1}{2} BC$$

$$\Rightarrow AS = BQ$$

In $\triangle APS$ and $\triangle BPQ$

$$AP = BP$$

$$AS = BQ$$

$$\angle PAS = \angle PBQ$$

$$\triangle APS \cong \triangle BPQ$$

$$\therefore PS = PQ \quad \dots(v)$$

From (iii) and (v), we have

PQRS is a rhombus **Proved.**

Q.4. ABCD is a trapezium in which $AB \parallel DC$, BD is a diagonal and E is the mid-point of AD. A line is drawn through E parallel to AB intersecting BC at F (see Fig.). Show that F is the mid-point of BC.

Sol. Given : A trapezium ABCD with $AB \parallel DC$, E is the mid-point of AD and $EF \parallel AB$.

To Prove : F is the mid-point of BC.

Proof : $AB \parallel DC$ and $EF \parallel AB$

$\Rightarrow AB, EF$ and DC are parallel.




Intercepts made by parallel lines AB, EF and DC on transversal AD are equal.

$\therefore$ Intercepts made by those parallel lines on transversal BC are also equal.

i.e., $BF = FC$

$\Rightarrow$ F is the mid-point of BC.


Q.5. In a parallelogram ABCD, E and F are the mid-points of sides AB and CD respectively (see Fig.). Show that the line segments AF and EC trisect the diagonal BD.



Sol. Given : A parallelogram ABCD, in which E and F are mid-points of sides AB and DC respectively.

To Prove : $DP = PQ = QB$

Proof : Since E and F are mid-points of AB and DC respectively.



$$\Rightarrow AE = \frac{1}{2} AB \text{ and } CF = \frac{1}{2} DC \quad \dots(i)$$

$$\text{But, } AB = DC \text{ and } AB \parallel DC \quad \dots(ii)$$

[Opposite sides of a parallelogram]

$$\therefore AE = CF \text{ and } AE \parallel CF.$$

$\Rightarrow$ AECF is a parallelogram.

[One pair of opposite sides is parallel and equal]

In $\triangle BAP$,

E is the mid-point of AB

$EQ \parallel AP$

$\Rightarrow$ Q is mid-point of PB

[Converse of mid-point theorem]

$\Rightarrow PQ = QB$

$\dots(iii)$

Similarly, in $\triangle DQC$,

P is the mid-point of DQ

$DP = PQ$

$\dots(iv)$

From (iii) and (iv), we have

$DP = PQ = QB$

or line segments AF and EC trisect the diagonal BD. **Proved.**


Q.6. Show that the line segments joining the mid-points of the opposite sides of a quadrilateral bisect each other.

Sol. Given : ABCD is a quadrilateral in which EG and FH are the line segments joining the mid-points of opposite sides.

To Prove : EG and FH bisect each other.

Construction : Join EF, FG, GH, HE and AC.

Proof : In $\triangle ABC$, E and F are mid-points of AB and BC respectively.



$$\therefore EF = \frac{1}{2} AC \text{ and } EF \parallel AC \quad \dots(i)$$

In $\triangle ADC$, H and G are mid-points of AD and CD respectively.

$$\therefore HG = \frac{1}{2} AC \text{ and } HG \parallel AC \quad \dots(ii)$$

From (i) and (ii), we get

$EF = HG$ and $EF \parallel HG$

$\therefore$ EFGH is a parallelogram.

[$\because$ a quadrilateral is a parallelogram if its one pair of opposite sides is equal and parallel]

Now, EG and FH are diagonals of the parallelogram EFGH.

$\therefore$ EG and FH bisect each other.

[Diagonal of a parallelogram bisect each other] **Proved.**

Q.7. *ABC is a triangle right angled at C. A line through the mid-point M of hypotenuse AB and parallel to BC intersects AC at D. Show that*

(i) *D is the mid-point of AC.*

(ii) *MD ⊥ AC*

(iii) *CM = MA = $\frac{1}{2}$ AB*


Sol. Given : A triangle ABC, in which $\angle C = 90^\circ$ and M is the mid-point of AB and $BC \parallel DM$.

To Prove : (i) D is the mid-point of AC

[Given]

(ii) $DM \perp BC$

(iii) $CM = MA = \frac{1}{2} AB$



Construction : Join CM.

Proof : (i) In $\triangle ABC$,

M is the mid-point of AB.

[Given]

$BC \parallel DM$

[Given]

D is the mid-point of AC

[Converse of mid-point theorem] **Proved.**

(ii) $\angle ADM = \angle ACB$ [$\because$ Corresponding angles]

But $\angle ACB = 90^\circ$

[Given]

$\therefore \angle ADM = 90^\circ$

But $\angle ADM + \angle CDM = 180^\circ$

[Linear pair]

$\therefore \angle CDM = 90^\circ$

Hence, $MD \perp AC$ **Proved.**

(iii) $AD = DC$...(1) [$\because$ D is the mid-point of AC]

Now, in $\triangle ADM$ and $\triangle CMD$, we have

$\angle ADM = \angle CDM$

[Each = 90°]

$AD = DC$

[From (1)]

$DM = DM$

[Common]

$\therefore \triangle ADM \cong \triangle CMD$

[SAS congruence]

$\Rightarrow CM = MA$

...(2) [CPCT]

Since M is mid-point of AB,

$\therefore MA = \frac{1}{2} AB$

...(3)

Hence, $CM = MA = \frac{1}{2} AB$ **Proved.** [From (2) and (3)]