

Ostwald dilution law : The dissociation of weak electrolyte obeys law of mass action. The application of law of mass action to weak electrolyte is known as Ostwald dilution law. **It is to be noted here, that either acids or bases are weak electrolytes but salts are usually strong electrolytes.**

Consider a weak electrolyte say HA (an acid)

$$HA \rightleftharpoons H^+ + A^-$$

Before dissociation	1	0	0
After dissociation	$1 - \alpha$	α	α

Where α is degree of dissociation.

Let C be the concentration of HA in mol litre⁻¹ before its dissociation then, $[HA] = C(1 - \alpha)$; $[H^+] = C\alpha$; $[A^-] = C\alpha$

According to law of mass action

$$K_a = \frac{[H^+][A^-]}{[HA]} = \frac{(C\alpha \cdot C\alpha)}{C(1 - \alpha)} = \frac{C\alpha^2}{1 - \alpha}$$

If α is small, then $1 - \alpha \approx 1$.


$$K_a = C\alpha^2 \quad \dots(1)$$

or

$$\alpha = \sqrt{\left(\frac{K_a}{C}\right)} \quad \dots(2)$$

where K_a is dissociation constant of acid.

Similarly, for a weak base



$\therefore$

$$\alpha = \sqrt{\left(\frac{K_b}{C}\right)} \quad \dots(4)$$

where K_b is dissociation constant of base.

Note: 1. K_a and K_b are just the equilibrium constants and hence depend only on temperature.

2. Greater are the values of K_a or K_b more is the strength of acid or base respectively.

3. Relations 1 to 4 are valid for mono basic acid and mono acid base.

4. Different expressions have to be derived for dibasic or tribasic acids and diacidic or triacidic base.


Acid and base

Arrhenius concept



- (1) An acid furnishes H^+ ions in solution and a base furnishes OH^- ions in solutions
- (2) The strength of acid depends upon its tendency to furnish H^+ ions.

Bronsted concept

- (1) An acid is a proton donor; a base is a proton acceptor.
- (2) The strength of an acid depends upon its tendency to donate proton. More is the tendency to donate proton, more is acidic nature.
- (3) Water is amphoteric as it donates as well as accepts proton.



- (4) Each cation is acid and each anion is base.
- (5) A pair of acid and base which differ by a proton is known as conjugate pair of acid and base.



Lewis concept

- (1) Acids are electron pair acceptor, e.g., BF_3 , $AlCl_3$.
- (2) Bases are electron pair donor, e.g., NH_3 , PCl_3 .

Relative strength of Acids and Bases :


For weak acids :

$$\begin{aligned} \text{Relative strength} &= \frac{\text{Strength of I acid}}{\text{Strength II acid}} \\ &= \frac{[H^+] \text{ furnished by I acid}}{[H^+] \text{ furnished by II acid}} = \frac{C_1\alpha_1}{C_2\alpha_2} \\ &= \frac{C_1}{C_2} \times \sqrt{\frac{K_{a1}C_2}{K_{a2}C_1}} = \sqrt{\frac{K_{a1}C_1}{K_{a2}C_2}} \dots(5) \\ &= \sqrt{\frac{K_{a1}}{K_{a2}}} \text{ (if } C_1 = C_2) \dots(6) \end{aligned}$$


For strong acids :

$$\text{Relative strength} = \frac{K_1}{K_2} = \frac{\text{rate constant for a reaction catalysed by I acid}}{\text{rate constant for a reaction catalysed by II acid}}$$

Note : For rate constant of acid catalysed reactions, see ester hydrolysis and sugar hydrolysis in **chemical kinetics**.



Ionic product of water : Pure water, a weak electrolyte dissociates as



$$\therefore K = \frac{[\text{H}^+][\text{OH}^-]}{[\text{H}_2\text{O}]} \quad (\text{According to law of mass action})$$

$$\text{or } K_w = K[\text{H}_2\text{O}] = [\text{H}^+][\text{OH}^-] \quad \dots(7)$$

$\therefore [\text{H}_2\text{O}]$ is constant since only one molecule out of 550 million molecules of H_2O dissociates, i.e., $\alpha_{\text{H}_2\text{O}} = \frac{1}{550 \times 10^6}$

$$\alpha_{\text{H}_2\text{O}} = 1.8 \times 10^{-9} = 1.8 \times 10^{-7} \%$$

here K_w is a characteristic constant for water, known as ionic product of water. The numerical value of K_w is $10^{-14} \text{ mol}^2 \text{ litre}^{-2}$ at 25°C . The value of K_w however increases with increase in temperature.

pH and pOH values : Sorenson used a new term to express $[\text{H}^+]$ as pH defined as the negative power raised on 10 in order to express $[\text{H}^+]$

$$\text{i.e., } [\text{H}^+] = 10^{-\text{pH}} \quad \dots(8)$$

$$\text{or } \text{pH} = -\log [\text{H}^+] \quad \dots(9)$$

(i.e., negative logarithm of magnitude of $[\text{H}^+]$)

Thus, for pure water $\text{pH} = 7$ i.e., Neutral solution
 $\text{pH} = 0$ to 7 i.e., Acidic solution
 $\text{pH} = 7$ to 14 i.e., Alkaline solution



$$\text{Also } \therefore [\text{H}^+][\text{OH}^-] = 10^{-14}$$

$$\therefore (-\log \text{H}^+) + (-\log \text{OH}^-) = 14$$

$$\therefore \text{pH} + \text{pOH} = 14 \quad \dots(10)$$

Common ion effect : The phenomenon in which degree of dissociation of a **weak electrolyte** is suppressed due to the presence of **common ion** present in it.

Consider a weak electrolyte along with common ion, e.g.,



$$\text{For acetic acid } K_a = \frac{[\text{CH}_3\text{COO}^-][\text{H}^+]}{[\text{CH}_3\text{COOH}]}$$

Thus, $[\text{CH}_3\text{COO}^-]$ in solution becomes more in presence of CH_3COONa . However K_a is constant. Thus, in order to have K_a constant, $[\text{H}^+]$ in solution must decrease or $[\text{CH}_3\text{COOH}]$, i.e., undissociated acid must increase or in other words degree of dissociation of CH_3COOH decreases.

Note : In presence of a common ion (from strong electrolyte) present with weak electrolyte, the concentration of common ion is derived from strong electrolyte.


Buffer solutions : Solutions which possess reserve acidic nature or alkaline nature or solutions which resist change or do not show significant change in pH due to dilution or addition of small quantities of acid or alkali are known as buffer solutions.

Types of buffer solutions :


(1) **Simple buffer** : Salts of weak acid and weak base, e.g., $\text{CH}_3\text{COONH}_4$, NH_4CN , etc.

(2) **Mixed buffers** : These are of two types.

(a) **Acidic buffer mixtures** : A weak acid + its salts or its conjugate base





(b) **Basic buffer mixtures** : A weak base + its salt or its conjugate acid



Henderson equation for pH and pOH of mixed buffers

Acidic buffer mixtures : Consider a buffer mixture of weak acid + its conjugate base i.e., $\text{CH}_3\text{COOH} + \text{CH}_3\text{COONa}$



Applying law of mass action to Eq. (B)

$$K_a = \frac{[\text{H}^+][\text{CH}_3\text{COO}^-]}{[\text{CH}_3\text{COOH}]} \quad \text{or } [\text{H}^+] = \frac{K_a[\text{CH}_3\text{COOH}]}{[\text{CH}_3\text{COO}^-]}$$

$$\text{or } -\log \text{H}^+ = -\log K_a + \log \frac{[\text{CH}_3\text{COO}^-]}{[\text{CH}_3\text{COOH}]}$$

$$\text{or } \text{pH} = -\log K_a + \log \frac{[\text{Conjugate base}]}{[\text{Acid}]} \quad \dots(11)$$


$$\text{pH} = \text{p}K_a + \log \frac{[\text{Conjugate base}]}{[\text{Acid}]} \quad \dots(12)$$

Similarly for basic buffers i.e., A weak base + its conjugate acid


$$\text{pOH} = -\log K_b + \log \frac{[\text{Conjugate acid}]}{[\text{Base}]} \quad \dots(13)$$

$$\text{or } \text{pOH} = \text{p}K_b + \log \frac{[\text{Conjugate acid}]}{[\text{Base}]} \quad \dots(14)$$



Addition of acid or base to a buffer : The anion of salt consumes H^+ given by acid and thus, [anion] in buffer decreases and [acid] in buffer increases and pH changes slightly.



The undissociated acid consumes OH^- given by alkali and thus, [anion] in buffer increases and [acid] in buffer decreases which results a change in pH slightly.



Similarly, acid or alkali addition brings in following changes in basic buffer solutions.



Note: No doubt pH of buffer solutions experimentally remains unchanged on addition of small amount of acid or base but it change theoretically to a very small extent which however cannot be detected by pH meters.

Buffer capacity : The property of a buffer solution to which it resist the change in pH on addition of acid or alkali. It is expressed as :

$$\phi = \frac{d(b)}{dpH} \quad \dots(15)$$

where b is no. of mole of acid or base added to one litre solution and dpH is change in pH.

e.g., suppose pH of buffer changes from 4.745 to 4.832 when 0.01 mole of KOH is added to 0.5 litre buffer.



$$\phi = \frac{0.01 \times 2}{0.087} = 0.230$$

pK_a and pK_b for a conjugate acid-base pair

$$pK_a + pK_b = pK_w = 14 \quad \dots(16)$$

- Note :** 1. Stronger is acid, weaker is its conjugate base.
2. Higher is the value of pK_a of an acid, lower is acid strength and higher is basic strength of its conjugate base.

Solubility product : Consider a sparingly soluble salts in water. Let solubility of it be s mol litre⁻¹.



According to law of mass action,

$$K = \frac{[A^{+y}]^x [B^{-x}]^y}{[A_x B_y]} \text{ or } K[A_x B_y] = [A^{+y}]^x [B^{-x}]^y$$


$$K_{sp} = [A^{+y}]^x [B^{-x}]^y \quad \dots(17)$$

where K_{sp} is a characteristic constant for a given solute known as solubility product and defined as the product of ionic concentrations with suitable powers. K_{sp} however changes with temperature.


Putting $[A^{+y}] = xs$ and $[B^{-x}] = ys$ in Eq. (17);

$$K_{sp} = (xs)^x (ys)^y = x^x \cdot y^y (s)^{x+y} \quad \dots(18)$$

e.g., For AgCl



For Ag₂CrO₄



Now if


$[A^{+y}]^x [B^{-x}]^y > K_{sp}$; The solid $A_x B_y$ will precipitate out

$[A^{+y}]^x [B^{-x}]^y < K_{sp}$; No precipitation

$[A^{+y}]^x [B^{-x}]^y = K_{sp}$; No precipitation upto this limit or precipitation will just start at this condition. This is a limiting case.

Effect of common ion on solubility

Consider saturated solution of AgCl. If a salt having either of the ion common to AgCl say KCl is added to it, then



$$\text{For AgCl: } K_{sp} = [Ag^+][Cl^-]$$

$[Cl^-]$ increases in solution due to presence of KCl and thus to have K_{sp} constant, $[Ag^+]$ will decrease or AgCl will precipitate out from solution, i.e., solubility of AgCl will decrease with increasing concentration of KCl in solution.

Let 0.1 M KCl(aq.) solution with AgCl(aq.). If solubility of AgCl is s mol litre⁻¹, then

$$\text{For AgCl: } K_{sp} = [Ag^+][Cl^-]$$

$$K_{sp} = s(s + 0.1)$$

s being small in comparison to 0.1 and thus may be neglected therefore,



$$K_{sp} = s \times 0.1 \text{ or } s_{AgCl} = \frac{K_{sp}}{0.1}$$

where s is solubility of AgCl in presence of 0.1 M KCl(aq.).

Note: 1. A solute on addition if undergoes complex formation with ions of sparingly soluble salt, the solubility of salt increases.

2. The solubility of sparingly soluble salt say AgCN is also influenced if the salt undergoes hydrolysis. Normally solubility of salts increases with hydrolysis.

Simultaneous solubility : The phenomenon when two or more sparingly soluble salts having one ion common (either cation AgCl and AgBr or anion BaSO₄ and CaSO₄) leads to simultaneous solubility equilibria. For example.



$$K_{sp} AgCl = [Ag^+][Cl^-]$$

$$K_{sp} AgBr = [Ag^+][Br^-]$$


Thus if both are present $[Ag^+]$ should be same

$$\text{or } \frac{K_{sp} AgCl}{K_{sp} AgBr} = \frac{[Cl^-]}{[Br^-]}$$

Hydrolysis : The phenomenon in which anion or cation furnished by a salt interacts with H₂O to produce acidic nature or alkaline nature is known as salt hydrolysis.

The nature of aqueous solution of salt to be acidic or alkaline depends upon the nature of salt.


Salts of strong acid and weak base : e.g., NH₄Cl, NH₄NO₃, CuSO₄, ZnCl₂, FeCl₃, etc.



NH_4Cl on dissolution in water furnish NH_4^+ and Cl^- ion. Cl^- with H^+ gives HCl , a strong acid which returns back all the H^+ to solution. On the other hand NH_4^+ interacts with OH^- to give NH_4OH , a weak electrolyte having low degree of dissociation. Also due to common ion effect of unhydrolysed NH_4^+ , the degree of dissociation of NH_4OH is further suppressed and thus in solution the $[\text{H}^+]$ becomes more and it acquires acidic character. It is thus suggested that acidic character of NH_4Cl solution is due to hydrolysis of NH_4^+ ion or the hydrolysis of weak component of salt.

Hydrolysis constant and degree of hydrolysis :

Consider a salt say NH_4Cl in water. Let C mol litre⁻¹ be the concentration of salt and h the degree of hydrolysis of salt.



∴ According to law of mass action,

$$K = \frac{[\text{NH}_4\text{OH}][\text{H}^+]}{[\text{NH}_4^+][\text{H}_2\text{O}]} \quad \text{or} \quad K[\text{H}_2\text{O}] = \frac{[\text{NH}_4\text{OH}][\text{H}^+]}{[\text{NH}_4^+]} \quad \dots(19)$$

or $K_H = \frac{[\text{NH}_4\text{OH}][\text{H}^+]}{[\text{NH}_4^+]} \quad \dots(19)$

Now for weak base $\text{NH}_4\text{OH} \rightleftharpoons \text{NH}_4^+ + \text{OH}^-$

$$K_b = \frac{[\text{NH}_4^+][\text{OH}^-]}{[\text{NH}_4\text{OH}]} \quad \dots(20)$$

By Eqs. (19) and (20), $K_H \times K_b = [\text{H}^+][\text{OH}^-] = K_w$

or $K_H = \frac{K_w}{K_b} \quad \dots(21)$

Further, $[\text{NH}_4^+] = C(1-h)$

$$[\text{NH}_4\text{OH}] = Ch$$

$$[\text{H}^+] = Ch$$

∴ By Eqs. (19), $K_H = \frac{Ch \cdot Ch}{C(1-h)} = \frac{Ch^2}{(1-h)} \quad \dots(22)$

∴ h is small ∴ $1-h \approx 1$ or $K_H = Ch^2$

or $h = \sqrt{\left(\frac{K_H}{C}\right)} \quad \dots(23)$

Also pH is derived by $[\text{H}^+]$

$$[\text{H}^+] = Ch$$

By Eq. (23), $[\text{H}^+] = C \sqrt{\left(\frac{K_H}{C}\right)} = \sqrt{(K_H \cdot C)} \quad \dots(24)$

By Eqs. (21) and (24), $[\text{H}^+] = \sqrt{\left(\frac{K_w}{K_b} \cdot C\right)}$

∴ $-\log \text{H}^+ = \frac{1}{2} [\log K_b - \log K_w - \log C]$

∴ $\text{pH} = \frac{1}{2} [\log K_b - \log K_w - \log C] \quad \dots(25)$

Thus, following results can be written for case I and similar derivations can be made for case II and case III.

Case I : Salt of strong acid and weak base : e.g., NH_4Cl

$$K_H = \frac{K_w}{K_b}; h = \sqrt{\left(\frac{K_H}{C}\right)} \quad \text{and} \quad \text{pH} = \frac{1}{2} [\log K_b - \log K_w - \log C]$$

$$= \frac{1}{2} [\text{p}K_w - \text{p}K_b - \log C]$$

Case II : Salts of weak acid + Strong base : e.g., CH_3COONa , KCN , etc.

$$K_H = \frac{K_w}{K_a}; h = \sqrt{\left(\frac{K_H}{C}\right)} \quad \text{and} \quad \text{pOH} = \frac{1}{2} [\log K_a - \log K_w - \log C]$$

$$= \frac{1}{2} [\text{p}K_w - \text{p}K_a - \log C]$$

Case III : Salts of weak acid + weak base : e.g., $\text{CH}_3\text{COONH}_4$, NH_4CN , etc.

$$K_H = \frac{K_w}{K_a \cdot K_b}; h = \sqrt{K_H} \quad (\text{if } h \text{ is small}) \quad \text{or} \quad \frac{h}{1-h} = \sqrt{K_H}$$

$$\text{pH} = \frac{1}{2} [\log K_b - \log K_a - \log K_w] = \left[\frac{1}{2} \text{p}K_w + \frac{1}{2} \text{p}K_a - \frac{1}{2} \text{p}K_b \right]$$

Case IV : Salts of strong acid + strong base : e.g., NaCl , KNO_3 , etc.

This category of salt does not undergo salt hydrolysis.

NOTE : (i) Eq. (21) and analogous equation for other category should be used only when the degree of hydrolysis is negligible.

(ii) All the derivations made for salt hydrolysis are for uni-univalent salts. For other types of salts (say uni-bivalent, e.g., $\text{Na}_2\text{C}_2\text{O}_4$ or bi-bivalent CaC_2O_4 , one should derive the similar equations following the same above methods). The term ' C ' in the above equations however represents the concentration of ion that undergoes hydrolysis.

(iii) The pH of acidic salts of polyprotic acids in water say NaHS is given by : $\text{pH} = \frac{[\text{p}K_{a_1} + \text{p}K_{a_2}]}{2}$

● NUMERICAL PROBLEMS ●

- Calculate the amount of acetic acid present in one litre of its solution having $\alpha = 1\%$ and $K_a = 1.8 \times 10^{-5}$.
- 0.16 g of N_2H_4 are dissolved in water and the total volume made upto 500 mL. Calculate the percentage of N_2H_4 that has reacted with water in this solution. The K_b for N_2H_4 is 4.0×10^{-6} M. (Roorkee 1998)
- Nicotinic acid ($K_a = 1.4 \times 10^{-5}$) is represented by the formula $HNiC$. Calculate its per cent dissociation in a solution which contains 0.10 mole of nicotinic acid per 2.0 litre of solution. (Roorkee 1993)
- Saccharin ($K_a = 2 \times 10^{-12}$) is a weak acid represented by formula $HSaC$. 4×10^{-4} mole of saccharin is dissolved in 200 cm^3 water of pH 3. Assuming no change in volume, calculate the concentration of SaC^- ions in the resulting solution at equilibrium. (Roorkee 1994)
- Acetyl salicylic acid, i.e., aspirin ionises in water as :
 $HC_9H_7O_4 + H_2O \rightleftharpoons H_3O^+ + C_9H_7O_4^-$;
 $K_a = 2.75 \times 10^{-9}$
 If two tablets of aspirin each of 0.32 g is dissolved in water to produce 250 mL solution, calculate $[H^+]$, $[OH^-]$ and $[C_9H_7O_4^-]$ in solution.
- The ionisation constant of NH_4^+ in water is 5.6×10^{-10} at 25°C . The rate constant for the reaction of NH_4^+ and OH^- to form NH_3 and H_2O at 25°C is $3.4 \times 10^{10} \text{ litre mol}^{-1} \text{ sec}^{-1}$. Calculate the rate constant for proton transfer from water to NH_3 . (IIT 1996)
- Calculate the concentration of fluoroacetic acid which is required to get $[H^+] = 1.50 \times 10^{-3}$ M. K_a of acid $= 2.6 \times 10^{-3}$.
- An aqueous solution contains 10% ammonia by mass and has a density of 0.99 g cm^{-3} . Calculate hydroxyl and hydrogen ion concentration in this solution. K_a for $NH_4^+ = 5.0 \times 10^{-10}$ M. (Roorkee 1995)
- Liquid ammonia ionises to a slight extent. At -50°C , its ionisation constant, $K_{NH_3} = [NH_4^+][NH_2^-] = 10^{-30}$. How many amide ions are present per cm^3 of pure liquid ammonia? Assume $N = 6.0 \times 10^{23}$.
- The ionisation constant for pure formic acid, $K = [HCOOH_2^+][HCOO^-]$ has been estimated as 10^{-6} at room temperature. What percentage of formic acid molecules in pure formic acid are converted to formate ion? The density of formic acid is 1.22 g/cm^3 .
- Calculate the dissociation constant of NH_4OH at 25°C , if ΔH° and ΔS° for the given changes are as follows :
 $NH_3 + H^+ \rightleftharpoons NH_4^+; \quad \Delta H^\circ = -52.2 \text{ kJ mol}^{-1}$
 $H_2O \rightleftharpoons H^+ + OH^-; \quad \Delta H^\circ = 56.6 \text{ kJ mol}^{-1}$
 $NH_4OH \rightleftharpoons NH_4^+ + OH^-; \quad \Delta S^\circ = -76.53 \text{ JK}^{-1} \text{ mol}^{-1}$
- Prove that degree of dissociation of a weak acid is given by:

$$\alpha = \frac{1}{1 + 10^{(pK_a - pH)}}$$
 where K_a is its dissociation constant.
- Determine degree of dissociation of 0.05 M NH_3 at 25°C in a solution of pH = 11.
- In the qualitative analysis, Bi^{3+} is detected by the appearance of the precipitates of bismuthyl hydroxide $[BiO(OH)_5]$. Calculate the pH when the following equilibrium exists K_b for $BiO(OH) = 4 \times 10^{-10}$.
- Calculate the concentration of all species of significant concentrations present in 0.1 M H_3PO_4 solution. $K_1 = 7.5 \times 10^{-3}$, $K_2 = 6.2 \times 10^{-8}$, $K_3 = 3.6 \times 10^{-13}$. Also calculate pH of solution.
- Calculate the $[OH^-]$ of $[NH_2 \cdot C_2H_4NH_3]^+$ and $[H_3N - C_2H_4NH_3]^{2+}$ in 0.15 M ethylene diamine (aq.); if
 $NH_2C_2H_4NH_2 + H_2O \rightleftharpoons NH_2C_2H_4NH_3^+ + OH^-$;
 $K_1 = 8.5 \times 10^{-5}$
 $NH_2C_2H_4NH_3^+ + H_2O \rightleftharpoons [NH_3C_2H_4NH_3]^{2+} + OH^-$; $K_2 = 2.7 \times 10^{-8}$
- Calculate pH of (1) 10^{-3} N HNO_3 , (2) 10^{-3} M H_2SO_4 , (3) 10^{-3} N H_2SO_4 , (4) 0.01 N HCl , (5) 10^{-8} N HCl , (6) 10^2 M HCl .
- Calculate pH for acid solutions having $[H^+]$ as (a) $[H^+] = 0.05$ M, (b) $[H^+] = 5.0$ M, (c) $[H^+] = 10^{-8}$ M.
- Calculate pH for.
 (a) 0.001 N $NaOH$ (b) 0.01 N $Ca(OH)_2$
 (c) 0.01 M $Ca(OH)_2$ (d) 10^{-8} M $NaOH$
 (e) 10^2 M $NaOH$ (f) 0.0008 M $Mg(OH)_2$
 Assume complete ionisation of each. (Roorkee 1992)
- Calculate pH of basic solutions having $[OH^-]$ as
 (a) $[OH^-] = 0.05$ M, (b) $[OH^-] = 5.0$ M,
 (c) $[OH^-] = 10^{-8}$ M.
- Calculate pH of (Given K_a $CH_3COOH = K_a$ $NH_4OH = 2.0 \times 10^{-3}$)
 (a) 0.002 N acetic acid having 2.3% dissociation.
 (b) 0.002 N NH_4OH having 2.3% dissociation.

22. The average concentration of SO_2 in the atmosphere over a city on a certain day is 10 ppm, when the average temperature is 298 K. Given that the solubility of SO_2 in water at 298 K is $1.3653 \text{ mol litre}^{-1}$ and the pK_a of H_2SO_3 is 1.92, estimate the pH of rain on that day. (IIT 2000)
23. Calculate $[\text{H}^+]$ and $[\text{CHCl}_2\text{COO}^-]$ in a solution that is 0.01 M HCl and 0.01 M in CHCl_2COOH . K_a for CHCl_2COOH is 5×10^{-2} .
24. A solution contains 0.09 M CHCl_2COOH and 0.1 M CH_3COOH . The pH of this solution is 1. If K_a for acetic acid is 10^{-5} , calculate K_a for CHCl_2COOH .
25. What is $[\text{H}^+]$ for a solution in which
(i) $\text{pH} = 3$ (ii) $\text{pH} = 4.75$?
26. The pH of 0.05 M aqueous solution of diethyl amine is 12.0. Calculate K_b . (Roorkee 1993)
27. The K_w for $2\text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{O}^+ + \text{OH}^-$ changes from 10^{-14} at 25°C to 9.62×10^{-14} at 60°C . What is pH of water at 60°C ? What happens to its neutrality?
28. Ionic product of water (K_w) is 1×10^{-14} at 25°C . What are dissociation constant of water and autoprotolysis constant of water?
29. Will the pH of water be same at 4°C and 25°C ? Explain. (IIT 2003)
30. The pH of pure water at 25°C and 35°C are 7 and 6 respectively. Calculate the heat of formation of water from H^+ and OH^- .
31. A solution of HCl has a $\text{pH} = 5$. If one mL of it is diluted to 1 litre, what will be pH of resulting solution?
32. 2.0 g of diborane (B_2H_6) reacts with water to produce 100 mL solution. If K_a for H_3BO_3 is 7.3×10^{-10} , calculate pH of solution.
33. 100 mL of HCl gas at 25°C and 740 mm pressure were dissolved in one litre of water. Calculate the pH of solution. Given, V.P. of H_2O at 25°C is 23.7 mm.
34. Find the concentrations of H^+ , HCO_3^- and CO_3^{2-} in a 0.01 M solution of carbonic acid if the pH of solution is 4.18. $K_1 = 4.45 \times 10^{-7}$, $K_2 = 4.69 \times 10^{-11}$.
35. Calculate the $[\text{Cl}^-]$, $[\text{Na}^+]$, $[\text{H}^+]$, $[\text{OH}^-]$ and pH of resulting solution obtained by mixing 50 mL of 0.6 N HCl and 50 mL of 0.3 N NaOH .
36. Calculate the pH of solution obtained by mixing 10 mL of 0.1 M HCl and 40 mL of 0.2 M H_2SO_4 .
37. What is the pH of 1 M solution of acetic acid? To what volume one litre of this solution be diluted so that pH of the resulting solution will be twice of the original value? $K_a = 1.8 \times 10^{-5}$. (IIT 1990)
38. Calculate the pH of a solution which contains 100 mL of 0.1 M HCl and 9.9 mL of 1.0 M NaOH .
39. What will be the resultant pH when 200 mL of an aqueous solution of HCl ($\text{pH} = 2.0$) is mixed with 300 mL of an aqueous solution of NaOH ($\text{pH} = 12.0$)? (IIT 1998)
40. 500 mL of 0.2 M aqueous solution of acetic acid is mixed with 500 mL of 0.2 M HCl at 25°C .
(i) Calculate the degree of dissociation of acetic acid in the resulting solution and pH of the solution.
(ii) If 6 g of NaOH is added to the above solution, determine the final pH. [Assume there is no change in volume on mixing; K_a of acetic acid is $1.75 \times 10^{-5} \text{ mol L}^{-1}$.] (IIT 2002)
41. An aqueous solution of aniline of concentration 0.24 M is prepared. What concentration of sodium hydroxide is needed in this solution so that anilinium ion concentration remains at $1 \times 10^{-8} \text{ M}$? K_a for $\text{C}_6\text{H}_5\text{NH}_3^+$ is $2.4 \times 10^{-5} \text{ M}$. (Roorkee 1996)
42. Calculate $[\text{H}^+]$ in a solution containing 0.1 M HCOOH and 0.1 M HOCN . K_a for HCOOH and HOCN are 1.8×10^{-4} and 3.3×10^{-4} .
43. What is the concentration of acetic acid which has same pH as 0.5 M HCOOH solution. K_a $\text{HCOOH} = 2.4 \times 10^{-4}$ and K_a $\text{CH}_3\text{COOH} = 1.8 \times 10^{-5}$.
44. What are $[\text{H}^+]$, $[\text{A}^-]$ and $[\text{B}^-]$ in a solution that is 0.03 M HA and 0.1 M HB ? K_a for HA and HB are 1.38×10^{-4} and 1.05×10^{-10} respectively.
45. Calculate $[\text{H}^+]$, pH , $[\text{CH}_3\text{COO}^-]$ and $[\text{C}_7\text{H}_5\text{O}_2^-]$ in a solution containing 0.02 M CH_3COOH and 0.01 M $\text{C}_6\text{H}_5\text{COOH}$. $K_{\text{CH}_3\text{COOH}} = 1.8 \times 10^{-5}$ and $K_{\text{C}_6\text{H}_5\text{COOH}} = 6.4 \times 10^{-5}$.
46. What concentration of HCOO^- is present in a solution of 0.015 M HCOOH and 0.02 M HCl . K_a $\text{HCOOH} = 1.8 \times 10^{-4}$.
47. A solution contains 0.1 M H_2S and 0.3 M HCl . Calculate the conc. of S^{2-} and HS^- ions in solution. Given K_{a1} and K_{a2} for H_2S are 10^{-7} and 1.3×10^{-13} respectively. (Roorkee 1992)
48. The K_a for formic acid and acetic acid are 2.1×10^{-4} and 1.1×10^{-5} respectively. Calculate relative strength of acids.
49. Calculate the pH of a solution of given mixtures:
(a) $(2\text{g CH}_3\text{COOH} + 3\text{g CH}_3\text{COONa})$ in 100 mL of mixture; $K_a = 1.8 \times 10^{-5}$
(b) 5 mL of 0.1 M NH_4OH + 250 mL of 0.1 M NH_4Cl ; $K_b = 1.8 \times 10^{-5}$
(c) (0.25 mole of acid + 0.35 mole of salt) in 500 mL mixture; $K_a = 3.6 \times 10^{-4}$

50. Calculate the pH of a buffer solution prepared by dissolving 30 g of Na_2CO_3 in 500 mL of an aqueous solution containing 150 mL of 1 M HCl. K_a for $\text{HCO}_3^- = 5.63 \times 10^{-11}$.
51. 0.15 mole of pyridinium chloride has been added into 500 cm³ of 0.2 M pyridine solution. Calculate pH and hydroxyl ion concentration in the resulting solution assuming no change in volume.
(K_b for pyridine $= 1.5 \times 10^{-9}$ M) (Roorkee 1995)
52. Calculate the mass of $(\text{NH}_4)_2\text{SO}_4$ in g which must be added to 500 mL of 0.2 M NH_3 to yield a solution of pH = 9.35. K_b for $\text{NH}_3 = 1.78 \times 10^{-5}$.
53. What volume of 0.1 M sodium formate solution should be added to 50 mL of 0.05 M formic acid to produce a buffer solution of pH = 4.0; pK_a of formic acid = 3.80?
(Roorkee 1990)
54. How many mole of HCl will be required to prepare one litre of buffer solution (containing NaCN + HCl) of pH 8.5 using 0.01 g formula mass of NaCN? $K_{\text{HCN}} = 4.1 \times 10^{-10}$.
55. A 40 mL solution of weak base BOH is titrated with 0.1N HCl solution. The pH of solution is found to be 10.04 and 9.14 after the addition of 5.0 mL and 20.0 mL of acid respectively. Find out K_b for weak base.
(IIT 1991)
56. A weak acid HA after treatment with 12 mL of 0.1 M strong base BOH has a pH of 5. At the end point, the volume of same base required is 26.6 mL. Calculate K_a of acid.
57. A solution of weak acid was titrated with base NaOH. The equivalence point was reached when 36.12 mL of 0.1 M NaOH have been added. Now 18.06 mL 0.1 M HCl were added to titrated solution, the pH was found to be 4.92. What is K_a of acid?
58. Calculate $[\text{H}^+]$ in a 0.20 M solution of dichloroacetic acid ($K_a = 5 \times 10^{-2}$) that also contains 0.1 M sodium dichloroacetate. Neglect hydrolysis of sodium salt.
59. Calculate the change in pH of 1 litre buffer solution containing 0.1 mole each of NH_3 and NH_4Cl up on addition of:
(i) 0.02 mole of dissolved gaseous HCl.
(ii) 0.02 mole of dissolved NaOH. (Roorkee 1992)
Assume no change in volume. $K_{\text{NH}_3} = 1.8 \times 10^{-5}$
60. 20 mL of 0.2 M NaOH are added to 50 mL of 0.2 M acetic acid ($K_a = 1.8 \times 10^{-5}$).
(1) What is pH of solution?
(2) Calculate volume of 0.2 M NaOH required to make the pH of solution 4.74.
61. Calculate the ratio of pH of a solution containing 1 mole of CH_3COONa + 1 mole of HCl per litre and of other solution containing 1 mole CH_3COONa + 1 mole of acetic acid per litre.
62. A 0.1 M solution of weak acid HA is 1% dissociated at 25°C. What is its K_a ? If this solution is w.r.t. NaA 0.2 M, what will be the new degree of dissociation of HA and pH?
63. The $[\text{H}^+]$ in 0.2 M solution of formic acid is 6.4×10^{-3} mol litre⁻¹. To this solution formate is added so as to adjust the conc. of sodium formate to one mol per litre. What will be pH of this solution? K_a for HCOOH is 2.4×10^{-4} and degree of dissociation of HCOONa is 0.75.
64. What is the pH of a solution when 0.20 mole of HCl is added to one litre solution containing.
(a) 1 M each of acetic acid and acetate ion?
(b) 0.1 M each of acetic acid and acetate ion?
Given K_a for acetic acid is 1.8×10^{-5} .
65. Calculate the composition of an acidic buffer solution made up of HA and NaA of total molarity 0.29 having pH = 4.4 and $K_a = 1.8 \times 10^{-5}$.
66. Calculate the amount of NH_3 and NH_4Cl required to prepare a buffer solution of pH 9.0 when total concentration of buffering reagents is 0.6 mol litre⁻¹.
(Roorkee 1997)
67. Calculate the pH of a solution obtained by mixing 100 mL of 0.3 M HCl with 100 mL of 0.4 M NH_3 . pK_a for $\text{NH}_4^+ = 9.2552$.
68. Calculate the pH of a solution obtained by mixing 50 mL of 0.2 M NH_4Cl and 75 mL of 0.1 M NaOH. pK_a for $\text{NH}_4^+ = 9.2552$.
69. A certain buffer solution contains equal concentration of X^- and HX. K_b for X^- is 10^{-10} . Calculate pH of buffer.
70. Two buffer, (X) and (Y) of pH 4.0 and 6.0 respectively are prepared from acid HA and the salt NaA. Both the buffers are 0.50 M in HA. What would be the pH of the solution obtained by mixing equal volumes of the two buffers? ($K_{\text{HA}} = 1.0 \times 10^{-5}$) (Roorkee 1999)
71. A certain weak acid has $K_a = 1.0 \times 10^{-4}$. Calculate the equilibrium constant for its reaction with a strong base.
(IIT 1991)
72. The pH of blood stream is maintained by a proper balance of H_2CO_3 and NaHCO_3 concentrations. What volume of 5 M NaHCO_3 solution, should be mixed with 10 mL sample of blood which is 2 M in H_2CO_3 in order to maintain a pH of 7.4 K_a for H_2CO_3 in blood is 7.8×10^{-7} ?
(IIT 1993)

73. 0.1 M CH_3COOH solution is titrated against 0.05 M NaOH solution. Calculate pH at 1/4th and 3/4th stages of neutralization of acid. The pH for 0.1 M CH_3COOH is 3.
74. K_{sp} of AgCl is 1.5×10^{-10} at 25°C . Calculate solubility of AgCl in; (a) Pure water, (b) 0.1 M AgNO_3 , (c) 0.1 M NaCl. (Roorkee 1995)
75. How many grams of potassium bromide (molar mass 120) can be added to 0.5 litre of 0.05 M solution of silver nitrate just to start the precipitation of silver bromide? K_{sp} of AgBr is 5.0×10^{-13} .
76. The solubility of AgCl in water at 25°C is 1.79×10^{-3} g/litre. Calculate K_{sp} of AgCl at 25°C .
77. K_{sp} of AgBr is 4×10^{-13} and $[\text{Ag}^+]$ in a solution is 1×10^{-6} mol litre $^{-1}$. What is the $[\text{Br}^-]$ in that solution?
78. The $[\text{Ag}^+]$ ion in a saturated solution of Ag_2CrO_4 at 25°C is 1.5×10^{-4} M. Determine K_{sp} of Ag_2CrO_4 at 25°C .
79. 25 mL of a sample of clear saturated solution of PbI_2 requires 10 mL of a certain AgNO_3 (aq.) for its titration. What is the molarity of this AgNO_3 (aq.)? K_{sp} for $\text{PbI}_2 = 4 \times 10^{-9}$.
80. Equal volumes of 0.02 M CaCl_2 and 0.0004 M Na_2SO_4 are mixed. Will a precipitate form? (K_{sp} of $\text{CaSO}_4 = 2.4 \times 10^{-5}$).
81. What (H_3O^+) must be maintained in a saturated H_2S solution to precipitate Pb^{2+} , but not Zn^{2+} from a solution in which each ion is present at a concentration of 0.01 M? ($K_{sp}\text{H}_2\text{S} = 1.1 \times 10^{-22}$; $K_{sp}\text{ZnS} = 1.0 \times 10^{-21}$) (Roorkee 2000)
82. K_{sp} of PbCl_2 is 10^{-13} . What will be $[\text{Pb}^{2+}]$ in a solution prepared by mixing 100 mL of 0.1 M $\text{Pb}(\text{NO}_3)_2$ and 1 mL of 1 M HCl?
83. K_{sp} of PbBr_2 is 8×10^{-5} . If the salt is 80% dissociated in solution, calculate the solubility of salt in g per litre.
84. The solubility of $\text{Pb}(\text{OH})_2$ in water is 6.7×10^{-6} M. Calculate the solubility of $\text{Pb}(\text{OH})_2$ in a buffer solution of pH = 8. (IIT 1999)
85. The ionisation constant of benzoic acid is 6.46×10^{-5} and K_{sp} for $\text{C}_6\text{H}_5\text{COOAg}$ is 2.5×10^{-13} . How many times is silver benzoate more soluble in a buffer of pH 3.19 as compared to pure water.
86. Calculate the solubility of CaF_2 in a solution buffered at pH = 3.0. K_a for HF is 6.3×10^{-4} and K_{sp} of $\text{CaF}_2 = 3.45 \times 10^{-11}$.
87. Will a precipitate of $\text{Mg}(\text{OH})_2$ be formed in a 0.001 M solution of $\text{Mg}(\text{NO}_3)_2$, if the pH of solution is adjusted to 9? K_{sp} of $\text{Mg}(\text{OH})_2 = 8.9 \times 10^{-12}$.
88. Calculate pH at which $\text{Mg}(\text{OH})_2$ begins to precipitate from a solution containing 0.10 M Mg^{2+} ions. K_{sp} of $\text{Mg}(\text{OH})_2 = 1 \times 10^{-11}$. (Roorkee 1992)
89. Calculate the $[\text{OH}^-]$ of a solution after 100 mL of 0.1 M MgCl_2 is added to 100 mL of 0.2 M NaOH. K_{sp} of $\text{Mg}(\text{OH})_2$ is 1.2×10^{-11} .
90. A sample of AgCl was treated with 5.00 mL of 1.5 M Na_2CO_3 solution to give Ag_2CO_3 . The remaining solution contained 0.0026 g of Cl^- per litre. Calculate the solubility product of AgCl ($K_{sp}\text{Ag}_2\text{CO}_3 = 8.2 \times 10^{-12}$). (IIT 1997)
91. Calculate simultaneous solubility of AgCNS and AgBr in a solution of water K_{sp} of $\text{AgBr} = 5 \times 10^{-13}$ and K_{sp} of $\text{AgCNS} = 1 \times 10^{-12}$. (UPSEAT 1995)
92. A solution contains a mixture of Ag^+ (0.10 M) and Hg_2^{2+} (0.10 M) which are to be separated by selective precipitation. Calculate the maximum concentration of iodide ion at which one of them gets precipitated almost completely. What % of that metal ion is precipitated? (K_{sp} of $\text{AgI} = 8.5 \times 10^{-17}$ and K_{sp} of $\text{Hg}_2\text{I}_2 = 2.5 \times 10^{-26}$)
93. 0.01 mole of AgNO_3 is added to 1 litre of a solution which is 0.1 M in Na_2CrO_4 and 0.005 M in NaIO_3 . Calculate the mole of precipitate formed at equilibrium and the concentrations of Ag^+ , IO_3^- and CrO_4^{2-} . (K_{sp} values of Ag_2CrO_4 and AgIO_3 are 10^{-8} and 10^{-13} respectively). (Roorkee 2001)
94. The K_{sp} of $\text{Ca}(\text{OH})_2$ is 4.42×10^{-5} at 25°C . A 500 mL of saturated solution of $\text{Ca}(\text{OH})_2$ is mixed with equal volume of 0.4 M NaOH. How much $\text{Ca}(\text{OH})_2$ in mg is precipitated? (IIT 1992)
95. A sample of hard water contains 0.005 mole of CaCl_2 per litre. What is the minimum concentration of Na_2SO_4 which must be added for removing Ca^{2+} ions from this water sample? K_{sp} for CaSO_4 is 2.4×10^{-5} at 25°C .
96. 1.75 g of solid NaOH are added to 0.25 dm 3 of 0.1 M NiCl_2 solution. Calculate: (a) mass of $\text{Ni}(\text{OH})_2$ formed; (b) pH of final solution. Given, K_{sp} of $\text{Ni}(\text{OH})_2 = 1.6 \times 10^{-14}$.
97. A mixture of water and AgCl is shaken until a saturated solution is obtained. Now the solution is filtered and 100 mL of clear solution of filtrate is mixed with 100 mL

- of 0.03 M NaBr. Should a precipitate form? K_{sp} of AgCl and AgBr are 1×10^{-10} and 5×10^{-13} .
98. Calculate pH of a saturated solution of $\text{Mg}(\text{OH})_2$. K_{sp} for $\text{Mg}(\text{OH})_2$ is 8.9×10^{-12} .
99. 0.1 milli mole of CdSO_4 are present in 10 mL acid solution of 0.08 N HCl. Now H_2S is passed to precipitate all the Cd^{2+} ions. What would be the pH of solution after filtering off precipitate, boiling of H_2S and making the solution 100 mL by adding H_2O ?
100. Zn salt is mixed with $(\text{NH}_4)_2\text{S}$ of molarity 0.021 M. What mass of Zn^{2+} will remain unprecipitated in 12 mL of the solution? K_{sp} of $\text{ZnS} = 4.51 \times 10^{-24}$.
101. Freshly precipitated Al and Mg hydroxides are stirred vigorously in a buffer solution containing 0.25 M of NH_4Cl and 0.05 M of NH_4OH . Calculate $[\text{Al}^{3+}]$ and $[\text{Mg}^{2+}]$ in solution. K_b for $\text{NH}_4\text{OH} = 1.8 \times 10^{-5}$. K_{sp} of $\text{Al}(\text{OH})_3 = 6 \times 10^{-32}$ and K_{sp} of $\text{Mg}(\text{OH})_2 = 8.9 \times 10^{-12}$.
102. A solution has 0.05 M Mg^{2+} and 0.05 M NH_3 . Calculate the concentration of NH_4Cl required to prevent the formation of $\text{Mg}(\text{OH})_2$ in solution. K_{sp} of $\text{Mg}(\text{OH})_2 = 9.0 \times 10^{-12}$ and ionisation constant of NH_3 is 1.8×10^{-5} . (Roorkee 1993)
103. A particular water sample has 131 ppm CaSO_4 . What fraction of the water must be evaporated in a container before solid CaSO_4 begins to deposit K_{sp} of $\text{CaSO}_4 = 9.0 \times 10^{-6}$?
104. To a solution of 0.1 M Mg^{2+} and 0.8 M NH_4Cl , an equal volume of NH_3 is added which just gives precipitate. Calculate $[\text{NH}_3]$ in solution. K_{sp} of $\text{Mg}(\text{OH})_2 = 1.4 \times 10^{-11}$ and K_b of $\text{NH}_4\text{OH} = 1.8 \times 10^{-5}$.
105. 10 mL of 0.3 M Na_2SO_4 are mixed with 20 mL solution having initially 0.1 M Ca^{2+} and 0.1 M Sr^{2+} in it. What are the final concentrations of Ca^{2+} , Sr^{2+} and SO_4^{2-} in solution? Given K_{sp} of $\text{SrSO}_4 = 7.6 \times 10^{-7}$ and K_{sp} of $\text{CaSO}_4 = 2.4 \times 10^{-5}$.
106. The solubility of CaCO_3 is 7 mg/litre. Calculate the solubility product of BaCO_3 from this information and from the fact that when Na_2CO_3 is added slowly to a solution containing equimolar concentration of Ca^{2+} and Ba^{2+} , no precipitate is formed until 90% of Ba^{2+} has been precipitated as BaCO_3 .
107. The solubility of $\text{Mg}(\text{OH})_2$ is increased by the addition of NH_4^+ ion. Calculate.
- (a) K_C for,

$$\text{Mg}(\text{OH})_2 + 2\text{NH}_4^+ \rightleftharpoons 2\text{NH}_3 + 2\text{H}_2\text{O} + \text{Mg}^{2+}$$
 K_{sp} of $\text{Mg}(\text{OH})_2 = 6 \times 10^{-12}$, K_b of $\text{NH}_3 = 1.8 \times 10^{-5}$
- (b) Find solubility of $\text{Mg}(\text{OH})_2$ in a solution containing 0.5 M NH_4Cl before addition of $\text{Mg}(\text{OH})_2$.
108. The K_{sp} of $\text{Ag}_2\text{C}_2\text{O}_4$ at 25°C is $1.29 \times 10^{-11} \text{ mol}^3 \text{ L}^{-3}$. A solution of $\text{K}_2\text{C}_2\text{O}_4$ containing 0.152 mole in 500 mL water is shaken at 25°C with excess of Ag_2CO_3 till the equilibrium is reached.


$$\text{Ag}_2\text{CO}_3 + \text{K}_2\text{C}_2\text{O}_4 \rightleftharpoons \text{Ag}_2\text{C}_2\text{O}_4 + \text{K}_2\text{CO}_3$$
At equilibrium the solution contains 0.0358 mole of K_2CO_3 . Assuming degree of dissociation of $\text{K}_2\text{C}_2\text{O}_4$ and K_2CO_3 to be same, calculate K_{sp} of Ag_2CO_3 . (IIT 1991)
109. Calculate the solubility of AgCN in a buffer solution of pH = 3. Given K_{sp} of AgCN = 1.2×10^{-16} and K_a for HCN = 4.8×10^{-10} .
110. Determine the concentration of NH_3 solution whose one litre can dissolve 0.10 mole AgCl. K_{sp} of AgCl and K_f of $[\text{Ag}(\text{NH}_3)_2]^+$ are $1.0 \times 10^{-10} \text{ M}^2$ and $1.6 \times 10^7 \text{ M}^{-2}$ respectively. (Roorkee 1999)
111. Predict whether or not AgCl will be precipitated from a solution which is 0.02 M in NaCl and 0.05 M in $[\text{Ag}(\text{CN})_2]^-$. Given $K_{\text{instability constant}}$ for $[\text{Ag}(\text{CN})_2]^- = 4.0 \times 10^{-19}$ and K_{sp} AgCl = 2.8×10^{-10} .
112. Given: $\text{Ag}(\text{NH}_3)_2^+ \rightleftharpoons \text{Ag}^+ + 2\text{NH}_3$, $K_C = 6.2 \times 10^{-8}$ and K_{sp} of AgCl = 1.8×10^{-10} at 298 K. Calculate the concentration of the complex in 1.0 M aqueous ammonia. (IIT 1998)
113. Equal volumes of 0.02 M AgNO_3 and 0.02 M HCN are mixed. What is $[\text{Ag}^+]$ in solution after attaining equilibrium? Given K_a HCN = 6.2×10^{-10} and K_{sp} AgCN = 2.2×10^{-16} .
114. Determine the mole of AgI which may be dissolved in 1.0 litre of 1.0 M CN^- solution. K_{sp} for AgI and K_C for $[\text{Ag}(\text{CN})_2]^-$ are $1.2 \times 10^{-17} \text{ M}^2$ and $7.1 \times 10^{19} \text{ M}^{-2}$ respectively. (Roorkee 1998)
115. 100.0 mL of a clear saturated solution of Ag_2SO_4 is added to 250.0 mL of a clear saturated solution of PbCrO_4 . Will any precipitate form and if so what? Given, K_{sp} values for Ag_2SO_4 , Ag_2CrO_4 , PbCrO_4 and PbSO_4 are 1.4×10^{-5} , 2.4×10^{-12} , 2.8×10^{-13} and 1.6×10^{-8} respectively.
116. 2 M solution of Na_2CO_3 is boiled in a closed container with excess of CaF_2 . Very little amount of CaCO_3 and

- NaF are formed. If the solubility product of CaCO_3 is x and molar solubility of CaF_2 is y , find the molar concentration of F^- in the resulting solution after equilibrium is attained.
117. 25.0 mL clear saturated solution of $\text{PbI}_2(aq.)$ requires 13.3 mL of $\text{AgNO}_3(aq.)$ solution for complete precipitation. What is molarity of AgNO_3 solution? K_{sp} of PbI_2 is 7.1×10^{-9} .
118. 250.0 mL of saturated clear solution of $\text{CaC}_2\text{O}_4(aq.)$ requires 6.3 mL of 0.00102 M $\text{KMnO}_4(aq.)$ in acid medium for complete oxidation of $\text{C}_2\text{O}_4^{2-}$ ions. Calculate the K_{sp} of CaC_2O_4 .
119. K_{sp} for $\text{SrF}_2 = 2.8 \times 10^{-9}$ at 25°C . How much NaF should be added to 100 mL of solution having 0.016 M in Sr^{2+} ions to reduce its concentration to $2.5 \times 10^{-3} \text{ M}$?
120. An aqueous solution of a metal bromide MBr_2 (0.05 M) is saturated with H_2S . What is the minimum pH at which MS will precipitate? K_{sp} for MS = 6.0×10^{-21} . Concentration of saturated $\text{H}_2\text{S} = 0.1 \text{ M}$; $K_1 = 10^{-7}$ and $K_2 = 1.3 \times 10^{-13}$ for H_2S . (IIT 1993)
121. H_2S is bubbled into a 0.02 M NaCN solution which is 0.02 M each in $\text{Ag}(\text{CN})_2^-$ and $\text{Cd}(\text{CN})_4^{2-}$. If K_{sp} of Ag_2S and K_{sp} of CdS are 1.0×10^{-50} and 7.1×10^{-28} and $K_{\text{instability}}$ for $[\text{Ag}(\text{CN})_2^-]$ and $[\text{Cd}(\text{CN})_4^{2-}]$ are 1.0×10^{-20} and 7.8×10^{-18} , which sulphide will precipitate first.
122. Calculate the pH at which an acid indicator with $K_a = 1 \times 10^{-5}$ changes colour when the indicator concentration is $1 \times 10^{-3} \text{ M}$. Also report the pH at which coloured ion is 80% present.
123. An acid type indicator. HIn differs in colour from its conjugate base (In^-). The human eye is sensitive to colour differences only when the ratio $[\text{In}^-]/[\text{HIn}]$ is greater than 10 or smaller than 0.1. What should be the minimum change in the pH of the solution to observe a complete colour change ($K_a = 1.0 \times 10^{-5}$)? (IIT 1997)
124. Bromophenol blue is an indicator with a value of $K_a = 5.84 \times 10^{-5}$. At what pH it will work as an indicator? Also report the % of this indicator in its basic form at a pH of 4.84.
125. Calculate the percentage hydrolysis in 0.003 M aqueous solution of NaOCN. K_a for HOCN = 3.33×10^{-4} . (Roorkee 1996)
126. What is the pH of a 0.5 M aqueous NaCN solution? pK_b of $\text{CN}^- = 4.70$. (IIT 1996)
127. Calculate degree of hydrolysis and pH of 0.2 M solution of NH_4Cl . Given K_b for NH_4OH is 1.8×10^{-5} .
128. Find out the mass of NH_4Cl dissolved in 500 mL to have pH = 4.5 K_b for NH_4OH is 1.8×10^{-5} .
129. (a) K_a for butyric acid is 2.0×10^{-5} . Calculate pH and hydroxyl ion concentration in 0.2 M aqueous solution of sodium butyrate. (Roorkee 1994)
(b) The dissociation constant of a substituted benzoic acid at 25°C is 1×10^{-4} . Calculate the pH of 0.01 M solution of its sodium salt. (IIT 2009)
130. K_a for ascorbic acid (HAsc) is 5×10^{-5} . Calculate the hydrogen ion concentration and percentage of hydrolysis in an aqueous solution in which the concentration of Asc^- ions is 0.02 M . (Roorkee 1997)
131. Calculate the pH at the equivalence point when a solution of 0.1 M acetic acid is titrated with a solution of 0.1 M NaOH. K_a for acid = 1.9×10^{-5} . (Roorkee 1990)
132. 0.1 M NaOH is titrated with 0.1 M HA till the end point. K_a of HA is 5.6×10^{-6} and degree of dissociation is less compared to 1. Calculate the pH of the resulting solution at the end point. (IIT 2004)
133. Calcium lactate is a salt of weak acid and represented as $\text{Ca}(\text{LaC})_2$. A saturated solution of $\text{Ca}(\text{LaC})_2$ contains 0.13 mole of salt in 0.50 litre solution. The pOH of this is 5.60. Assuming complete dissociation of salt, calculate K_a of lactic acid. (Roorkee 1991)
134. Calculate the pH of 0.1 M K_3PO_4 solution. The third dissociation constant of orthophosphoric acid is 1.3×10^{-12} . Assume that the hydrolysis proceeds only in the first step.
135. Equilibrium constant for the acid ionisation of Fe^{3+} to $\text{Fe}(\text{OH})^{2+}$ and H^+ is 6.5×10^{-3} . What is the maximum pH which could be used so that at least 95% of the total Fe^{3+} in a dilute solution exists as Fe^{3+} ?
136. The acid ionisation constant for

$$\text{Zn}^{2+} + \text{H}_2\text{O} \rightleftharpoons \text{Zn}(\text{OH})^+ + \text{H}^+$$
 is 1.0×10^{-9} . Calculate the pH of 0.0010 M solution of ZnCl_2 . Also calculate basic dissociation constant of $\text{Zn}(\text{OH})^+$.
137. The dissociation constants for aniline, acetic acid and water at 25°C are 3.83×10^{-10} , 1.75×10^{-5} and 1.008×10^{-14} respectively. Calculate degree of hydrolysis of aniline acetate in a deci normal solution. Also report the pH.
138. Calculate the pH of an aqueous solution of 1.0 M ammonium formate assuming complete dissociation. (pK_a of formic acid = 3.8 and pK_b of ammonia = 4.8) (IIT 1995)

139. Calculate pH of the following mixtures. Given that $K_a = 1.8 \times 10^{-5}$ and $K_b = 1.8 \times 10^{-5}$.
- 50 mL of 0.10 M NaOH + 50 mL of 0.05 M CH_3COOH .
 - 50 mL of 0.05 M NaOH + 50 mL of 0.10 M CH_3COOH .
 - 50 mL of 0.10 M NaOH + 50 mL of 0.10 M CH_3COOH .
 - 50 mL of 0.10 M NH_4OH + 50 mL of 0.05 M HCl.
 - 50 mL of 0.05 M NH_4OH + 50 mL of 0.10 M HCl.
 - 50 mL of 0.10 M NH_4OH + 50 mL of 0.10 M HCl.
 - 50 mL of 0.05 M NH_4OH + 50 mL of 0.05 M CH_3COOH .
140. The vapour pressure of 0.01 molal solution of weak base BOH in water at 20°C is 17.536 mm. Calculate K_b for base. Aqueous tension at 20°C is 17.540 mm. Assume molality and molarity same.
141. A 0.01 M aqueous solution of weak acid HA has an osmotic pressure 0.293 atm 25°C. Another 0.01 M aqueous solution of other weak acid HB has an osmotic pressure of 0.345 atm under the same conditions. Calculate equilibrium constants of two acids for their dissociation.
142. The salt $\text{Zn}(\text{OH})_2$ is involved in the following two equilibria :
- $$\text{Zn}(\text{OH})_2(s) \rightleftharpoons \text{Zn}^{2+}(aq.) + 2\text{OH}^-(aq.)$$
- $$K_{sp} = 1.2 \times 10^{-17}$$
- $$\text{Zn}(\text{OH})_2(s) + 2\text{OH}^- \rightleftharpoons \text{Zn}(\text{OH})_4^{2-}(aq.) \quad K_f = 0.12$$
- Calculate the $[\text{OH}^-]$ at which solubility of $\text{Zn}(\text{OH})_2$ be a minimum. Also find the solubility of $\text{Zn}(\text{OH})_2$ at this pH.
143. A 500 mL sample of an equilibrium mixture of gaseous N_2O_4 and NO_2 at 25°C and 753 mm of Hg was allowed to react with enough water to make 250.0 mL of solution at 25°C. Assume that all the dissolved N_2O_4 is converted to NO_2 which disproportionates in water yielding a solution of nitrous acid and nitric acid. Assume further that disproportionation reaction goes to completion and that none of the nitrous acid disproportionates. The equilibrium constant (K_p) for $\text{N}_2\text{O}_4(g) \rightleftharpoons 2\text{NO}_2(g)$ is 0.113 at 25°C. K_a for HNO_2 is 4.5×10^{-4} at 25°C.
- Write balanced equation for disproportionation.
 - What is molar concentration of NO_2^- and pH of the solution?
 - What is osmotic pressure of solution?
 - How many grams of lime (CaO) would be required to neutralize the solution?
144. K_a for HCN and CH_3COOH are 4.9×10^{-10} and 1.8×10^{-5} respectively. Calculate the equilibrium constant for the reaction:
- $$\text{CH}_3\text{COOH} + \text{NaCN} \rightleftharpoons \text{CH}_3\text{COONa} + \text{HCN}$$
145. It is found that 0.1 M solution of three sodium salts NaX, NaY and NaZ have pH 7.0, 9.0 and 11.0 respectively. Arrange the acids (HX, HY and HZ) in order of increasing acidic character. Where possible calculate dissociation constant of acid.
146. Calculate the pH of 0.05 M $\text{KHC}_8\text{H}_4\text{O}_4$
- $$\text{H}_2\text{C}_8\text{H}_4\text{O}_4 + \text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{O}^+ + \text{HC}_8\text{H}_4\text{O}_4^-; \text{p}K_{a1} = 2.94$$
- $$\text{HC}_8\text{H}_4\text{O}_4^- + \text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{O}^+ + \text{C}_8\text{H}_4\text{O}_4^{2-}; \text{p}K_{a2} = 5.44$$
147. A buffer solution of 0.080 M Na_2HPO_4 and 0.020 M Na_3PO_4 is prepared. The electrolytic oxidation of 1.0 m mole RNHOH is carried out in 100 mL buffer to give.
- $$\text{RNHOH} + \text{H}_2\text{O} \longrightarrow \text{RNO}_2 + 4\text{H}^+ + 4e^-$$
- Calculate approximate pH of the solution after oxidation is complete. $\text{p}K_{a1}$, $\text{p}K_{a2}$ and $\text{p}K_{a3}$ of H_3PO_4 are 2.12, 7.20 and 12.0 respectively.
148. Calculate the difference in pH for 1/3 and 2/3 stages of neutralisation of 0.10 M CH_3COOH and with 0.10 M NaOH.
149. The molar conductivity of a solution of a weak acid HX (0.01 M) is 10 times smaller than the molar conductivity of a solution of a weak acid HY (0.1 M). If $\lambda_{X^-}^0 \approx \lambda_{Y^-}^0$, the difference in their $\text{p}K_a$ values $\text{p}K_a(\text{HX}) - \text{p}K_a(\text{HY})$, is (consider degree of ionization of both acids to be $\ll 1$) [JEE (Advanced II) 2015]

SOLUTIONS (Numerical Problems)



where α is degree of dissociation of acid, if C mol/litre is concentration of acid, then

$$[\text{H}^+] = C\alpha; [\text{CH}_3\text{COO}^-] = C\alpha; [\text{CH}_3\text{COOH}] = C(1-\alpha)$$



$$\text{Also } K_a = \frac{[\text{H}^+][\text{CH}_3\text{COO}^-]}{[\text{CH}_3\text{COOH}]} = \frac{C\alpha \cdot C\alpha}{C(1-\alpha)} = C\alpha^2$$

$\therefore K_a$ is small. $\therefore \alpha$ will also be small and thus $1-\alpha \approx 1$

$$\text{or } 1.8 \times 10^{-5} = C \times \left(\frac{1}{100}\right)^2 \therefore C = 0.18$$

$\therefore$ 1 litre solution contains = 0.18 mole of CH_3COOH

$\therefore$ 1 litre solution contains = $0.18 \times 60 = 10.8 \text{ g CH}_3\text{COOH}$



$$\text{Also } K_b = \frac{C\alpha^2}{(1-\alpha)}$$



Assuming $1-\alpha \approx 1$

$$K_b = C\alpha^2$$

$$[\text{N}_2\text{H}_4] = C = \frac{0.16 \times 1000}{32 \times 500} = 0.01$$

$$\text{Given } K_b = 4 \times 10^{-6} \text{ M} \therefore \alpha^2 = \frac{4 \times 10^{-6}}{0.01} = 4 \times 10^{-4}$$

$$\alpha = 2 \times 10^{-2} \text{ i.e., } \alpha = 0.02 \text{ or } 2\%$$





$$\text{Also, } C = \frac{0.1}{2} = 5 \times 10^{-2} \text{ mol litre}^{-1}; K_a = 1.4 \times 10^{-5}$$

$$\therefore K_a = \frac{C\alpha^2}{(1-\alpha)} = C\alpha^2 \quad (\because 1-\alpha \approx 1)$$

$$\therefore \alpha = \sqrt{\left(\frac{K_a}{C}\right)} = \sqrt{\left(\frac{1.4 \times 10^{-5}}{5 \times 10^{-2}}\right)} = 1.67 \times 10^{-2} \text{ or } 1.67\%$$

$$4. \quad [\text{HSaC}] = \frac{\text{mole}}{\text{litre}} = \frac{4 \times 10^{-4}}{200/1000} = 2 \times 10^{-3} \text{ M}$$

The dissociation of HSaC takes place in presence of $[\text{H}^+] = 10^{-3}$



In presence of H^+ the dissociation of HSaC is almost negligible because of common ion effect. Thus, at equilibrium

$$[\text{HSaC}] = 2 \times 10^{-3}; [\text{H}^+] = 10^{-3}$$

$$\therefore K_a = \frac{[\text{H}^+][\text{SaC}^-]}{[\text{HSaC}]} \therefore 2 \times 10^{-12} = \frac{[10^{-3}][\text{SaC}^-]}{2 \times 10^{-3}}$$

$$\therefore [\text{SaC}^-] = 4 \times 10^{-12} \text{ M}$$



$$5. \quad [\text{HC}_9\text{H}_7\text{O}_4] = \frac{0.32 \times 2 \times 1000}{180 \times 250} = 0.014 \text{ M}$$

$$\therefore \alpha = \sqrt{\frac{K_a}{C}} = \sqrt{\frac{2.75 \times 10^{-9}}{0.014}} = 4.43 \times 10^{-4}$$

$$\therefore [\text{H}^+] = C \cdot \alpha = 0.014 \times 4.43 \times 10^{-4} = 6.21 \times 10^{-6} \text{ M}$$



$$[\text{C}_9\text{H}_7\text{O}_4^-] = C \cdot \alpha = 6.21 \times 10^{-6} \text{ M}$$

$$[\text{OH}^-] = \frac{10^{-14}}{[\text{H}^+]} = \frac{10^{-14}}{6.21 \times 10^{-6}} = 1.61 \times 10^{-9} \text{ M}$$



$$K_{(\text{base})\text{NH}_3} = \frac{K_f}{K_b} = \frac{K_w}{K_{\text{acid}}(\text{NH}_4^+)} \quad (\because K_{\text{acid}} \times K_{\text{base}} = K_w)$$

$$\text{or } \frac{K_f}{3.4 \times 10^{-10}} = \frac{10^{-14}}{5.6 \times 10^{-10}} \therefore K_f = 6.07 \times 10^{-5}$$



$$\text{Given, } [\text{H}^+] = C \cdot \alpha = 1.5 \times 10^{-3} \text{ mole litre}^{-1}$$

$$\therefore K_a = \frac{(C\alpha)(C\alpha)}{C(1-\alpha)} = \frac{C\alpha^2}{(1-\alpha)}$$

$$2.6 \times 10^{-3} = \frac{1.5 \times 10^{-3} \times \alpha}{(1-\alpha)} \therefore \alpha = 0.634$$

$$\text{Now, } C \cdot \alpha = 1.50 \times 10^{-3}$$

$$\therefore C = \frac{1.50 \times 10^{-3}}{0.634} = 2.37 \times 10^{-3} \text{ M}$$


Note: Since K_a is of the order of 10^{-3} M and thus it is not advisable to use $K_a = C\alpha^2$. Because $(1-\alpha)$ is not equal to 1 since α is not small.

$$8. \text{ Given, } \frac{\text{mass of NH}_3}{\text{mass of solution}} = \frac{10}{100}$$

$$\therefore 100 \text{ g solution contains } 10 \text{ g NH}_3$$

$$\therefore M_{\text{NH}_3} = (10 \times 1000) / [(17 \times (100/0.99))] = 5.82$$

$$(\because V = \text{mass} / \text{density})$$




$$\therefore [\text{OH}^-] = C \cdot \alpha = C \sqrt{(K_b/C)} = \sqrt{(K_b \cdot C)}$$

$$[\because C = 5.82 \text{ M and } K_b = K_w / K_a = 10^{-14} / (5 \times 10^{-10}) = 2 \times 10^{-5}]$$

$$\therefore [\text{OH}^-] = \sqrt{[2 \times 10^{-5} \times 5.82]} = 1.07 \times 10^{-2} \text{ M}$$

$$\therefore [\text{H}^+] = 10^{-14} / 1.07 \times 10^{-2} = 0.9268 \times 10^{-12} \text{ M}$$

$$\therefore \text{pH} = -\log[\text{H}^+] = -\log 0.9268 \times 10^{-12} = 12.0330$$



$$\therefore [\text{NH}_2^-] = \sqrt{K} = \sqrt{10^{-30}} = 10^{-15} \text{ M}$$

$$\text{Number of amide ions in } 10^3 \text{ cm}^3 = 10^{-15} \times 6 \times 10^{23}$$

$$\therefore \text{Number of amide ions in one cm}^3 = \frac{10^{-15} \times 6 \times 10^{23}}{10^3} = 6 \times 10^5 \text{ ions}$$

10. Given density of formic acid = 1.22 g/cm^3

$$\therefore \text{Mass of formic acid in 1 litre solution} = 1.22 \times 10^3 \text{ g}$$

$$\text{Thus, } [\text{HCOOH}] = \frac{1.22 \times 10^3}{46} = 26.5 \text{ M}$$




$$\text{Since in case of auto ionisation } [\text{HCOOH}_2^+] = [\text{HCOO}^-]$$

$$\text{and } [\text{HCOO}^-][\text{HCOOH}_2^+] = 10^{-6} \therefore [\text{HCOO}^-] = 10^{-3}$$

Now % dissociation of

$$\text{HCOOH} = \frac{[\text{HCOO}^-] \times 100}{[\text{HCOOH}]} = \frac{10^{-3}}{26.5} \times 100$$

$$\left[\alpha = \frac{C\alpha}{C(1-\alpha)} \right] = 0.004\%$$



Similarly, ΔS° for the change = $-76.53 \text{ JK}^{-1} \text{ mol}^{-1}$

or for the change :



$$\text{and } \Delta S^\circ = -76.53 \text{ JK}^{-1} \text{ mol}^{-1}$$

Now we have $\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$


$$\therefore \Delta G^\circ = 4.4 - (-76.53 \times 10^{-3}) \times 298 = 27.21 \text{ KJ mol}^{-1}$$

$$\text{Also, } \Delta G^\circ = -2.303 RT \log K_b$$

$$27.21 = -2.303 \times 8.314 \times 10^{-3} \times 298 \log K_b$$

$$\therefore K_b = 1.7 \times 10^{-5}$$

12. For a weak monoprotic acid



$$[\text{H}^+] = C\alpha \quad \dots(i)$$

$$\text{and } K_a = \frac{C\alpha^2}{(1-\alpha)} \quad \dots(ii)$$

By substituting value of C from Eq. (ii) in Eq. (i)

$$[\text{H}^+] = \frac{K_a \cdot (1-\alpha) \cdot \alpha}{\alpha^2} = \frac{K_a(1-\alpha)}{\alpha} \quad \dots(iii)$$


$$\text{Thus, } -\log[\text{H}^+] = -[\log K_a + \log(1-\alpha) - \log \alpha]$$

$$\text{or } \text{pH} = \log \frac{1}{K_a} + \log \frac{\alpha}{(1-\alpha)} \quad \dots(iv)$$

$$\text{Also, } \text{pH} = \text{p}K_a - \log \left[\frac{1-\alpha}{\alpha} \right] \text{ or } \log \frac{1-\alpha}{\alpha} = [\text{p}K_a - \text{pH}]$$

$$\frac{1-\alpha}{\alpha} = 10^{(\text{p}K_a - \text{pH})} \text{ or } \frac{1}{\alpha} - 1 = 10^{(\text{p}K_a - \text{pH})}$$

$$\text{or } \alpha = \frac{1}{1 + 10^{(\text{p}K_a - \text{pH})}}$$




$$\begin{array}{ccc} 1 & 0 & 0 \\ (1-\alpha) & \alpha & \alpha \end{array}$$

Given, $\text{pH} = 11$

$$\therefore [\text{H}^+] = 10^{-11} \therefore [\text{OH}^-] = 10^{-3} = C\alpha$$

Since, $C = 0.05$





$$\therefore \alpha = \frac{10^{-3}}{C} = \frac{10^{-3}}{0.05} = 2 \times 10^{-2} \text{ or } 2\%$$



$$\therefore [\text{OH}^-]^2 = 4 \times 10^{-10} \therefore [\text{OH}^-] = 2 \times 10^{-5}$$

$$\therefore \text{pOH} = 4.6989$$

$$\therefore \text{pH} = 9.3010$$



$$\begin{array}{ccc} 0.1 & 0 & 0 \\ 0.1-C & C & C \end{array}$$

$$K_1 = \frac{[\text{H}^+][\text{H}_2\text{PO}_4^-]}{[\text{H}_3\text{PO}_4]} = \frac{C \cdot C}{(0.1-C)}$$


$$7.5 \times 10^{-3} = \frac{C^2}{(0.1-C)} \therefore C = 0.024$$

$$\therefore [\text{H}^+] = 0.024 \text{ M} \therefore \text{pH} = 1.6198$$

$$[\text{H}_2\text{PO}_4^-] = 0.024 \text{ M}$$

$$[\text{H}_3\text{PO}_4] = 0.1 - 0.024 = 0.076 \text{ M}$$

The value of K_1 is much larger than K_2 and K_3 . Also dissociation of II and III steps occurs in presence of H^+ furnished in I step and thus, dissociation of II and III steps is further suppressed due to common ion effect.



$$\begin{array}{ccc} 0.024 & 0.024 & 0 \\ (0.024-y) & (0.024+y) & y \end{array}$$


The dissociation of H_2PO_4^- occurs in presence of $[\text{H}^+]$ furnished in step I.

$$\text{Thus, } K_2 = \frac{[\text{H}^+][\text{HPO}_4^{2-}]}{[\text{H}_2\text{PO}_4^-]} \text{ or } 6.2 \times 10^{-8} = \frac{(0.024+y)y}{(0.024-y)}$$

$\therefore y$ is small $\therefore 0.024 - y = 0.024$ and neglecting y^2 .

$$\therefore 6.2 \times 10^{-8} = \frac{0.024y}{0.024} \therefore y = 6.2 \times 10^{-8}$$

$$\text{or } [\text{HPO}_4^{2-}] = K_2 = 6.2 \times 10^{-8} \text{ (Insignificant)}$$



$$\begin{array}{ccc} (6.2 \times 10^{-8} - x) & (0.024 + x) & x \end{array}$$

$$\therefore K_3 = \frac{[H^+][PO_4^{3-}]}{[HPO_4^{2-}]} = \frac{(0.024+x) \cdot x}{(6.2 \times 10^{-8} - x)}$$

Again neglecting x^2 and assuming, $6.2 \times 10^{-8} - x = 6.2 \times 10^{-8}$

$$\therefore 3.6 \times 10^{-13} = \frac{0.024x}{6.2 \times 10^{-8}}$$

$$\therefore x = \frac{3.6 \times 10^{-13} \times 6.2 \times 10^{-8}}{0.024} = 9.3 \times 10^{-19}$$

(Insignificant)

Note: For weak polyprotic acid having no other electrolyte, the anion concentration produced in II step of dissociation is always equal to K_2 if concentration is reasonable.

16. **Case I:** $[OH^-] = [NH_2C_2H_4NH_3]^+$

$$= C \times \sqrt{\frac{K_{b1}}{C}} = \sqrt{0.15 \times 8.5 \times 10^{-5}}$$

$$= 3.57 \times 10^{-3} M$$

Case II: $NH_2C_2H_4NH_3^+ + H_2O \rightleftharpoons$

$$\begin{array}{ccc} [NH_3C_2H_4NH_3]^{2+} & + & OH^- \\ 3.57 \times 10^{-3} & 0 & 3.57 \times 10^{-3} \\ [3.57 \times 10^{-3} - X] & X & (X + 3.57 \times 10^{-3}) \end{array}$$

$$\therefore 2.7 \times 10^{-8} = \frac{X \cdot (X + 3.57 \times 10^{-3})}{(3.57 \times 10^{-3} - X)}$$


Neglecting X^2 , also $3.57 \times 10^{-3} - X = 3.57 \times 10^{-3}$; (X is very small)

$$2.7 \times 10^{-8} = \frac{3.57 \times 10^{-3} \cdot X}{3.57 \times 10^{-3}}$$

$$\therefore X = 2.7 \times 10^{-8}$$

$$\therefore [NH_2C_2H_4NH_3]^{2+} = 2.7 \times 10^{-8} M = K_{b2}$$

17. Strong acids ionise completely at normal dilutions.




$$\begin{array}{ccc} \text{Conc. before ionisation} & 10^{-3} N & 0 & 0 \\ \text{Conc. after ionisation} & 0 & 10^{-3} & 10^{-3} \end{array}$$

$$\therefore [H^+] = 10^{-3} \text{ mol/litre or Eq./litre } (\because H^+ \text{ is monovalent})$$

$$\therefore pH = -\log [H^+] = -\log 10^{-3}$$

$$pH = 3$$




$$\begin{array}{ccc} \text{Conc. before ionisation} & 10^{-3} M & 0 & 0 \\ \text{Conc. after ionisation} & 0 & 2 \times 10^{-3} & 10^{-3} \end{array}$$

$$\therefore \text{Mole ratio of } H_2SO_4 : H^+ : SO_4^{2-} :: 1 : 2 : 1$$

$$\therefore [H^+] = 2 \times 10^{-3} M$$

$$\therefore pH = -\log [H^+] = -\log 2 \times 10^{-3}$$

$$\therefore pH = 2.6989$$



$$\begin{array}{ccc} \text{Conc. before ionisation} & 10^{-3} N & 0 & 0 \\ \text{Conc. after ionisation} & 0 & 10^{-3} & 10^{-3} \end{array}$$

$\therefore$ Equal equivalent of a substance gives equal equivalent of its components.

$$\therefore [H^+] = 10^{-3} M$$


$$\therefore pH = -\log [H^+] \therefore pH = 3$$



$$\begin{array}{ccc} \text{Conc. before ionisation} & 10^{-2} N & 0 & 0 \\ \text{Conc. after ionisation} & 0 & 10^{-2} & 10^{-2} \end{array}$$

$$\therefore [H^+] = 10^{-2} M$$

$$\therefore pH = -\log [H^+] \therefore pH = 2$$



$$\begin{array}{ccc} \text{Conc. before ionisation} & 10^{-8} N & 0 & 0 \\ \text{Conc. after ionisation} & 0 & 10^{-8} & 10^{-8} \end{array}$$


$\therefore [H^+] = 10^{-8} M$ but $pH = 8$ is not possible because it is acid. Now $[H^+] = 10^{-7} M$ are already present in solution and since $10^{-8} < 10^{-7}$ and thus, it should not be neglected.

$$\therefore [H^+] = 10^{-8} + 10^{-7} = 10^{-7} (1.1) M = 1.1 \times 10^{-7} M$$

$$\therefore pH = 6.9586$$

Solution II The above solution lacks with discrepancy that dissociation of H_2O , a weak electrolyte is also suppressed in presence of HCl due to common ion effect and thus, $[H^+]_{H_2O} \neq 10^{-7}$ but will be lesser than this.

Therefore, dissociation of H_2O in presence of $10^{-8} H^+$.



$$\therefore K_w = (10^{-8} + a)a \quad \therefore a = 0.95 \times 10^{-7}$$

$$\therefore [H^+] = 10^{-8} + 0.95 \times 10^{-7} = 10^{-7} \times 1.05 = 1.05 \times 10^{-7}$$

$$\therefore pH = 6.9788$$



$$\begin{array}{ccc} \text{Conc. before dissociation} & 10^2 & 0 & 0 \\ \text{Conc. after dissociation} & 0 & 10^2 & 10^2 \end{array}$$

$$\therefore [H^+] = 10^2 M \quad \therefore pH = -2$$

Students are often under the illusion that it is impossible to have a negative pH. There is no theoretical basis for this. A negative pH only means that the hydrogen ion concentration is greater than 1 M. However in actual practice, a negative pH is uncommon because of two reasons. First, even strong acids (say 100% H_2SO_4) becomes partially dissociated at high concentrations. The second reason has to do with activity.

Sorensens originally intended pH to be related to $[H^+]$, but his fundamental method of measurement—the hydrogen electrode—is now known to depend on thermodynamics activities rather than $[H^+]$, i.e., on a_{H^+} and $a_{H^+} = [H^+] f_{H^+}$. In dilute solutions activity coefficient, f_{H^+} is near enough to unity and thus, $a_{H^+} = [H^+]$. At high concentrations, the activity coefficient is less than unity. Thus, pH defined by

$-\log [H^+]$ is not only of little theoretical significance, but in fact cannot be measured directly. It has therefore, come to be accepted that $pH = -\log_{10} a_{H^+}$ (this is what a pH meter reading is a measure of), i.e., pH of $10^2 M$ HCl cannot be calculated until f_{H^+} is known. Nevertheless, there is mathematically no basis for not having a negative pH.

18. (a) $[H^+] = 0.05 = 5 \times 10^{-2}$

$$\therefore pH = -\log[H^+] = -\log 5 \times 10^{-2}$$

$$pH = 1.3010$$


(b) $[H^+] = 5.0 M$ but $pH \neq -\log 5 \neq -0.6989$

See Problem 17 part 6.

(c) $[H^+] = 10^{-8}$ $\therefore pH \neq -\log 10^{-8} \neq 8$

See Problem 17 part 5.

19. (a) $0.001 N$ NaOH:




$$\therefore [OH^-] = 10^{-3} M \quad (\because N_{\text{NaOH}} = M_{\text{NaOH}})$$

$$\therefore pOH = -\log[OH^-] = -\log 10^{-3} = 3$$


$$\therefore pH = 14 - pOH = 14 - 3 = 11 \quad \therefore pH = 11$$

(b) $0.01 N$ Ca(OH)_2 :



$$\therefore [OH^-] = 10^{-2} M \quad \therefore pOH = 2 \quad \therefore pH = 12$$

(c) $0.01 M$ Ca(OH)_2 :



$$\therefore [OH^-] = 2 \times 10^{-2} M \quad \therefore pOH = 1.6989$$

$$\therefore pH = 14 - 1.6989 = 12.3010$$


(d) $10^{-8} M$ NaOH: $\text{NaOH} \longrightarrow \text{Na}^+ + \text{OH}^-$



$$\therefore [OH^-] = 10^{-8} M$$

Now proceed for OH^- as in problem 17 part 5.


(e) $10^2 M$ NaOH:



$$[OH^-] = 10^2 M$$

Now proceed as in problem 17 part 6.

(f) $0.0008 M$ Mg(OH)_2 :



$$\therefore [OH^-] = 16 \times 10^{-4} M \quad \therefore pOH = 2.7958$$

$$\therefore pH = 11.2041$$

20. (a) $[OH^-] = 0.05 M = 5 \times 10^{-2}$

$$\therefore pOH = -\log[OH^-] \quad \therefore pOH = 1.3010$$

$$\therefore pH = 12.6989$$



(b) $[OH^-] = 5$

Proceed as in problem 17 part 6.

(c) $[OH^-] = 10^{-8}$

Proceed as in problem 17 part 5.

21. (a) $0.002 N$ CH_3COOH : Acetic acid is weak electrolyte and partially dissociated.




$$\therefore [H^+] = C\alpha = 2 \times 10^{-3} \times \frac{2.3}{100} = 4.6 \times 10^{-5} M$$

$$\therefore pH = -\log[H^+] = -\log 4.6 \times 10^{-5}$$

$$pH = 4.3372$$

(b) $0.002 N$ NH_4OH : NH_4OH is weak base and partially dissociated.



$$\therefore [OH^-] = C\alpha = 2 \times 10^{-3} \times \frac{2.3}{100} = 4.6 \times 10^{-5} M$$

$$\therefore pOH = -\log[OH^-] = -\log 4.6 \times 10^{-5}$$

$$\therefore pOH = 4.3372 \quad \therefore pH = 14 - 4.3372$$

$$pH = 9.6628$$

22. Concentration of SO_2 in air is 10 ppm or 10 mole in 10^6 mole air or 10^{-5} mole SO_2 per mole of air. The concentration of SO_2 in air being substantial and since rain water is falling from enormously great height so, each drop of rain water will get saturated with SO_2 before it reaches earth.

Now the given concentration or solubility of SO_2 at 298 K is 1.3653 M. This value of solubility corresponds when $P_{\text{SO}_2} = 1 \text{ atm}$.

Thus according to Henry's law,

$[\text{SO}_2]$ dissolved in water $\propto P_{\text{SO}_2}$ in gas phase

Thus solubility of SO_2 at the condition of P_{SO_2} of 10^{-5} atm as $P \propto$ mole and therefore since solubility is reported at 1 atm

$$\therefore [\text{SO}_2] \text{ dissolved at pressure } 1 \text{ atm} = 1.3653 M$$

$$\therefore [\text{SO}_2]_2 \text{ dissolved at pressure } 10^{-5} \text{ atm}$$

$$= 1.3653 \times 10^{-5} M = C$$

$$\text{i.e., } [\text{H}_2\text{SO}_3] = [\text{SO}_2] = 1.3653 \times 10^{-5} M$$

Now, $\text{SO}_2 + \text{H}_2\text{O} \rightleftharpoons \text{H}^+ + \text{HSO}_3^-$



$$\therefore K_a = \frac{C\alpha \cdot C\alpha}{C(1 - \alpha)} = \frac{C\alpha^2}{(1 - \alpha)}$$

or $0.012 = \frac{1.3653 \times 10^{-5} \times \alpha^2}{(1-\alpha)}$
 $(\because pK_b = 1.92 \text{ and thus } K_b = 0.012)$
 $1.3653 \times 10^{-5} \alpha^2 + 0.012\alpha - 0.012 = 0$
 $\alpha = \frac{-0.012 \pm \sqrt{1.44 \times 10^{-5} + 4 \times 1.44 \times 10^{-5}}}{2}$
 $= 1$
 $\therefore [H^+] = C \times \alpha = 1.3653 \times 10^{-5}$
 $\therefore \text{pH} = 4.8648$

23. $\text{CHCl}_2\text{COOH} \rightleftharpoons \text{CHCl}_2\text{COO}^- + \text{H}^+$

C	0	0.01
$C(1-\alpha)$	$C\alpha$	$C\alpha + 0.01$

 $\therefore K_a = \frac{C\alpha \times (C\alpha + 0.01)}{C(1-\alpha)} = \frac{\alpha(0.01\alpha + 0.01)}{(1-\alpha)} = 5 \times 10^{-2}$
or $\frac{0.01\alpha(1+\alpha)}{(1-\alpha)} = 5 \times 10^{-2}$ or $\alpha^2 + 6\alpha - 5 = 0$
 $\therefore \alpha = 0.7416$
 $\therefore [\text{CHCl}_2\text{COO}^-] = 0.01 \times 0.7416 = 7.416 \times 10^{-3} M$
 $[H^+] = 7.416 \times 10^{-3} + 0.01 = 0.0174 M$

24. pH will be decided by $[H^+]$ furnished by HCl and CHCl_2COOH

$\text{CHCl}_2\text{COOH} \rightleftharpoons \text{CHCl}_2\text{COO}^- + \text{H}^+$

Initial conc.	0.09	0	0.09 (From HCl)
Final conc.	$(0.09 - x)$	x	$(0.09 + x)$

 $\therefore [H^+] = 0.09 + x$
but $\text{pH} = 1$, $\therefore [H^+] = 10^{-1} = 0.1$
 $\therefore 0.09 + x = 0.1$ $\therefore x = 0.01$
 K_a for CHCl_2COOH can be given as
 $K_a = \frac{[H^+][\text{CHCl}_2\text{COO}^-]}{[\text{CHCl}_2\text{COOH}]} = \frac{0.1 \times 0.01}{(0.09 - 0.01)} = 1.25 \times 10^{-2}$

25. (i) $\text{pH} = 3$ or $-\log[H^+] = 3$ $\therefore [H^+] = 10^{-3} M$
(ii) $\text{pH} = 4.75$ $\therefore -\log[H^+] = 4.75$
 $\therefore [H^+] = 1.7782 \times 10^{-5} M$

26. Diethyl amine is base and give OH^- as,
 $(\text{C}_2\text{H}_5)_2\text{NH} + \text{H}_2\text{O} \rightleftharpoons (\text{C}_2\text{H}_5)_2\text{NH}_2^+ + \text{OH}^-$

Initial conc.	1	0	0
Equilibrium	$(1-\alpha)$	α	α

 $\therefore [\text{OH}^-] = C\alpha$
where C is conc. of base and $C = 0.05 M$
 $\therefore \text{pH} = 12$ $\therefore \text{pOH} = 2$
or $[\text{OH}^-] = 10^{-2} M$ $\therefore C\alpha = 10^{-2}$
or $0.05 \times \alpha = 10^{-2}$ ($\because C = 0.05$)
 $\therefore \alpha = 0.2$
Now for a base, $K_b = \frac{C\alpha^2}{(1-\alpha)} = \frac{0.05 \times (0.2)^2}{(1-0.2)}$
 $= \frac{0.05 \times 0.04}{0.8} = 2.5 \times 10^{-3}$

Note: Do not use $K_b = C\alpha^2$ since $\alpha = 0.2$ and $1-\alpha = 0.8$.

27. K_w for H_2O at $25^\circ\text{C} = 10^{-14}$
 $\therefore [H^+][\text{OH}^-] = 10^{-14}$ ($\because K_w = [H^+][\text{OH}^-]$)
 $\therefore [H^+] = 10^{-7} M$ $\therefore \text{pH} = 7$
Now K_w for H_2O at $60^\circ\text{C} = 9.62 \times 10^{-14}$
 $\therefore [H^+][\text{OH}^-] = 9.62 \times 10^{-14}$
For pure water $[H^+] = [\text{OH}^-]$
 $\therefore [H^+]^2 = 9.62 \times 10^{-14}$
 $\therefore [H^+] = \sqrt{9.62 \times 10^{-14}} = 3.1 \times 10^{-7} M$
 $\therefore \text{pH} = -\log H^+ = -\log 3.1 \times 10^{-7}$
 $\text{pH} = 6.51$

Thus, pH of water becomes 6.51 at 60°C but the nature is neutral since calculation for pure water has been made, i.e. pH scale at 60°C becomes in between 0 to 13.02.

28. $\text{H}_2\text{O} \rightleftharpoons \text{H}^+ + \text{OH}^-$ (molarity of water = 55.6 M)
 $\therefore K_b = \frac{[H^+][\text{OH}^-]}{[\text{H}_2\text{O}]} = \frac{K_w}{[\text{H}_2\text{O}]} = \frac{10^{-14}}{55.6} = 1.8 \times 10^{-16}$
Also, $\text{H}_2\text{O} + \text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{O}^+ + \text{OH}^-$
 $K_{a-p} = \frac{[\text{H}_3\text{O}^+][\text{OH}^-]}{[\text{H}_2\text{O}]^2} = \frac{K_w}{[\text{H}_2\text{O}]^2}$
 $= \frac{10^{-14}}{(55.6)^2} = 3.24 \times 10^{-18}$

29. At 25°C , K_w for $\text{H}_2\text{O} = 10^{-14}$. Thus $[H^+] = [\text{OH}^-] = 10^{-7}$ or pH of water = 7. Also as the temperature decreases K_w for H_2O decrease (i.e., $K_w < 10^{-14}$) because of low dissociation of water ($\text{H}_2\text{O} \rightleftharpoons \text{H}^+ + \text{OH}^-$; $\Delta H = +ve$) following Le Chatelier's principle. This leads to lower concentrations of H^+ and OH^- ions. Thus pH of water is more at 4°C than at 25°C , however H_2O remains neutral.

30. At 25°C ; $[H^+] = 10^{-7}$ $\therefore K_w = 10^{-14}$
At 35°C ; $[H^+] = 10^{-6}$ $\therefore K_w = 10^{-12}$
Now using $2.303 \log_{10} \frac{K_{w_2}}{K_{w_1}} = \frac{\Delta H}{R} \left[\frac{T_2 - T_1}{T_1 \times T_2} \right]$
 $2.303 \log_{10} \frac{10^{-12}}{10^{-14}} = \frac{\Delta H}{2} \left[\frac{10}{298 \times 308} \right]$
 $\therefore \Delta H = 84551.4 \text{ cal/mol} = 84.551 \text{ kcal/mol}$
Thus, $\text{H}_2\text{O} \rightleftharpoons \text{H}^+ + \text{OH}^-$; $\Delta H = 84.551 \text{ kcal/mol}$
 $\therefore \text{H}^+ + \text{OH}^- \rightleftharpoons \text{H}_2\text{O}$; $\Delta H = -84.551 \text{ kcal/mol}$


31. $\text{HCl}_1 = 10^{-5} M$ since $\text{pH} = 5$
Meq. of HCl_1 in 1 mL = $10^{-5} \times 1$
Meq. of HCl_{II} in 1000 mL = $N \times 1000$
Since II is prepared by diluting I and Meq. does not change on dilution.

i.e., Meq. of HCl (concentrated) = Meq. of HCl (dilute)

$$\therefore 10^{-5} \times 1 = N \times 1000 \quad \therefore N_{\text{HCl}} = 10^{-8}$$

Now, proceed as in problem 17 part 5.


$$\therefore \text{pH} = 6.9788$$



1 mole (=27.6 g) of B_2H_6 = 2 mole H_3BO_3

$$\therefore 2.0 \text{ g of } \text{B}_2\text{H}_6 = \frac{2 \times 2}{27.6} = 0.145 \text{ mole } \text{H}_3\text{BO}_3$$

$$\therefore [\text{H}_3\text{BO}_3] = \frac{0.145 \times 1000}{100} = 1.45 \text{ M}$$



$$K_a = \frac{C\alpha^2}{1-\alpha} \quad (\because 1-\alpha \approx 1)$$

$$\text{or } 7.3 \times 10^{-10} = 1.45 \times \alpha^2 \quad \therefore \alpha = 2.24 \times 10^{-5}$$

$$\therefore [\text{H}^+] = C\alpha = 1.45 \times 2.24 \times 10^{-5} = 3.25 \times 10^{-5}$$

$$\therefore \text{pH} = 4.4881$$



$$\therefore P_{\text{HCl dry}} = \frac{740 - 23.7}{760} \text{ atm} \quad V = \frac{100}{1000} \text{ litre}$$

$$T = 25 + 273 = 298 \text{ K}$$

$$\therefore \text{Mole of HCl, i.e., } n_{\text{HCl}} = \frac{PV}{RT} = \frac{(740 - 23.7) \times 100}{760 \times 1000 \times 0.0821 \times 298}$$


$$\therefore n = 3.85 \times 10^{-3}$$

$$\text{Now, Molarity of HCl} = \frac{n}{V_{\text{solution}}} = \frac{3.85 \times 10^{-3}}{1} \text{ (V in litre)}$$


$$\therefore [\text{H}^+] = 3.85 \times 10^{-3} \text{ M}$$

$$\therefore \text{pH} = -\log \text{H}^+ = -\log 3.85 \times 10^{-3}$$

$$\text{pH} = 2.4142$$



$$\therefore [\text{H}^+] = 6.61 \times 10^{-5} \text{ mol litre}^{-1}$$




$$K_1 = \frac{[\text{H}^+][\text{HCO}_3^-]}{[\text{H}_2\text{CO}_3]}$$

$$\text{or } 4.45 \times 10^{-7} = \frac{[6.61 \times 10^{-5}][\text{HCO}_3^-]}{[0.01]}$$

$$\text{or } [\text{HCO}_3^-] = 6.73 \times 10^{-5} \text{ mol litre}^{-1}$$


Again for dissociation of HCO_3^- , we have



$$K_2 = \frac{[\text{H}^+][\text{CO}_3^{2-}]}{[\text{HCO}_3^-]}$$

$$\text{or } 4.69 \times 10^{-11} = \frac{[6.61 \times 10^{-5}][\text{CO}_3^{2-}]}{[6.73 \times 10^{-5}]}$$

$$\therefore [\text{CO}_3^{2-}] = 4.78 \times 10^{-11} \text{ mol litre}^{-1}$$



Meq. before reaction	50 × 0.6	50 × 0.3		
	= 30	= 15	0	0
Meq. after reaction	15	0	15	15

For monovalent electrolytes

$$\text{Molarity} = \text{Normality} = \frac{\text{milli equivalent}}{\text{total volume}}$$

$$\therefore [\text{Cl}^-] = \frac{15 + 15}{100} = 0.3 \text{ M (Cl}^-\text{ proved by HCl and NaCl)}$$



$$[\text{Na}^+] = \frac{15}{100} = 0.15 \text{ M}$$

$$[\text{H}^+] = \frac{15}{100} = 0.15 \text{ M}$$

$$[\text{OH}^-] = \frac{10^{-14}}{[\text{H}^+]} = \frac{10^{-14}}{0.15} = 6.6 \times 10^{-14} \text{ M}$$

$$\text{Also } \text{pH} = -\log [\text{H}^+] = -\log 0.15$$

$$\text{pH} = 0.8239$$



$$\therefore \text{Total Meq. of } \text{H}^+ \text{ in solution} = 1 + 16 = 17$$

$$\therefore [\text{H}^+] = \frac{17}{50} = 3.4 \times 10^{-1} \quad \left(\because [\text{H}^+] = \frac{\text{Meq.}}{V_{\text{in mL}}} \right)$$

$$\therefore \text{pH} = -\log [\text{H}^+] = -\log 0.34$$

$$\text{pH} = 0.4685$$



Conc. before dissociation	1	0	0
Conc. after dissociation	1 - α	α	α

$$[\text{H}^+] = C\alpha = C \sqrt{\left(\frac{K_a}{C} \right)} = \sqrt{(K_a C)} = \sqrt{(1.8 \times 10^{-5} \times 1)}$$

$$= \sqrt{(1.8 \times 10^{-5})} = 4.24 \times 10^{-3} \text{ M}$$

$$\therefore \text{pH} = -\log \text{H}^+ = -\log (4.24 \times 10^{-3})$$

$$\text{pH} = 2.3724$$

Case II: New pH is given as = 2.3724 × 2 = 4.7448

Let new conc. be C_1 and degree of dissociation be α_1

$$\therefore -\log [\text{H}^+] = 4.7448 \quad \therefore [\text{H}^+] = 1.8 \times 10^{-5}$$

$$\text{or } C_1 \alpha_1 = 1.8 \times 10^{-5}$$

$$\text{Now again } K_a = \frac{[\text{H}^+][\text{CH}_3\text{COO}^-]}{[\text{CH}_3\text{COOH}]}$$

$$K_a = \frac{C_1 \alpha_1 \times C_1 \alpha_1}{C_1 (1 - \alpha_1)} = \frac{C_1 \alpha_1 \cdot \alpha_1}{(1 - \alpha_1)}$$

$$\therefore 1.8 \times 10^{-5} = \frac{1.8 \times 10^{-5} \times \alpha_1}{(1 - \alpha_1)}$$

$$\alpha_1 = 0.5$$

$$\text{Now } C_1 \alpha_1 = 1.8 \times 10^{-5}$$

$$\therefore C_1 = \frac{1.8 \times 10^{-5}}{\alpha_1} = \frac{1.8 \times 10^{-5}}{0.5} = 3.6 \times 10^{-5} \text{ M}$$


Let 1 litre of conc. solution be diluted to V litre.

Ionic Equilibrium

Eq. of dilute solution = Eq. of concentration solution
 $3.6 \times 10^{-5} \times V = 1 \times 1$

$$\therefore V = \frac{1}{3.6 \times 10^{-5}} = 2.77 \times 10^4 \text{ litre}$$



Note : In II case α_1 comes 0.5 by $K_a = \frac{C_1 \alpha_1^2}{(1 - \alpha_1)}$ and thus, it is not advisable to assume $(1 - \alpha_1) = 1$.



$$\therefore [\text{H}^+] \text{ left from HCl} = \frac{0.1}{109.9} = 9.099 \times 10^{-4} \text{ M}$$

$$\therefore \text{pH} = -\log \text{H}^+ = -\log 9.099 \times 10^{-4}$$



$$\text{pH} = 3.0409$$



$$\therefore [\text{OH}^-] \text{ left from NaOH} = \frac{1}{500} = 2 \times 10^{-3} \text{ M}$$

$$\therefore \text{pOH} = -\log \text{OH}^- = -\log 2 \times 10^{-3}$$

$$\therefore \text{pOH} = 2.6989 \therefore \text{pH} = 11.3010$$



$$\therefore K_a = \frac{[\text{CH}_3\text{COO}^-][\text{H}^+]}{[\text{CH}_3\text{COOH}]} = \frac{x(0.1+x)}{(0.1-x)}$$

Due to common ion effect dissociation of CH_3COOH is very small in presence of HCl. Therefore, $(0.1+x) = 0.1$ and $(0.1-x) = 0.1$.

$$\therefore K_a = \frac{x \times 0.1}{0.1} \therefore x = K_a = 1.75 \times 10^{-5}$$


Thus, degree of dissociation

$$\alpha = \frac{x}{0.1} = \frac{1.75 \times 10^{-5}}{0.1} = 1.75 \times 10^{-4} = 0.000175 = 0.0175\%$$

$$\text{Also, } [\text{H}^+] = 0.1 + x = 0.1 \quad (x \ll 0.1)$$

$$\therefore \text{pH} = -\log [\text{H}^+] = -\log [0.1] = 1$$

(ii) Eq. of NaOH or mole of NaOH added = $\frac{6}{40} = 0.15$




Thus, the solution will act as acidic buffer having
 $[\text{CH}_3\text{COOH}] = \frac{0.05}{1000}$ and $[\text{CH}_3\text{COONa}] = \frac{0.05}{1000}$

$$\text{Thus, } \text{pH} = -\log K_a + \log \frac{[\text{Conjugate base}]}{[\text{Acid}]}$$

$$= -\log 1.75 \times 10^{-5} + \log \frac{[0.05/1000]}{[0.05/1000]}$$

$$\text{pH} = 4.757$$



$$\text{Thus, } K_b = \frac{[\text{C}_6\text{H}_5\text{NH}_3^+][\text{OH}^-]}{[\text{C}_6\text{H}_5\text{NH}_2]} \quad \dots(1)$$

$$\text{Also } K_b \text{ for } \text{C}_6\text{H}_5\text{NH}_2 = \frac{K_w}{K_a \text{ for } \text{C}_6\text{H}_5\text{NH}_3^+}$$

$$= \frac{1 \times 10^{-14}}{2.4 \times 10^{-5}} \quad \dots(2)$$

Since dissociation of $\text{C}_6\text{H}_5\text{NH}_2$ occurs in presence of NaOH and thus dissociation of $\text{C}_6\text{H}_5\text{NH}_2$ will suppress.



$$\text{Thus, } [\text{OH}^-] = ?; [\text{C}_6\text{H}_5\text{NH}_2] = 0.24; [\text{C}_6\text{H}_5\text{NH}_3^+] = 10^{-8}$$

$$\text{Therefore, } \frac{1 \times 10^{-14}}{2.4 \times 10^{-5}} = \frac{10^{-8} \times [\text{OH}^-]}{0.24}$$

$$\therefore [\text{OH}^-] = \frac{0.24 \times 10^{-14}}{2.4 \times 10^{-5} \times 10^{-8}} = 0.01$$

$$\therefore [\text{NaOH}] = 0.01 \text{ M}$$

42. In this problem both the acids contribute for $[\text{H}^+]$ due to appreciable dissociation. Thus,



Because $[\text{H}^+]$ will remain common in solution. Thus,

$$K_{\text{HCOOH}} = \frac{[\text{H}^+][\text{HCOO}^-]}{[\text{HCOOH}]} = 1.8 \times 10^{-4} \quad \dots(1)$$

$$K_{\text{HOCN}} = \frac{[\text{H}^+][\text{OCN}^-]}{[\text{HOCN}]} = 3.3 \times 10^{-4} \quad \dots(2)$$

$$\text{or } K_{\text{HCOOH}} = \frac{(x+y)x}{0.1} = 1.8 \times 10^{-4} \quad \dots(3)$$

$$K_{\text{HOCN}} = \frac{(x+y)y}{0.1} = 3.3 \times 10^{-4} \quad \dots(4)$$

$$\text{Thus, by Eqs. (3) and (4) } \frac{x}{y} = \frac{1.8}{3.3}$$

$$\text{or } y = 1.83x \quad \dots(5)$$

$$\text{From Eq. (3) } (x+1.83x) \cdot x = 1.8 \times 10^{-4} \therefore x = 2.52 \times 10^{-3}$$

$$\text{Therefore, } y = 4.61 \times 10^{-3}$$

$$\text{Thus, } [\text{H}^+] = x + y = 2.52 \times 10^{-3} + 4.61 \times 10^{-3}$$

$$= 7.13 \times 10^{-3} \text{ M}$$

43. $\text{HCOOH} \rightleftharpoons \text{H}^+ + \text{HCOO}^-$
 $\text{CH}_3\text{COOH} \rightleftharpoons \text{H}^+ + \text{CH}_3\text{COO}^-$
 Since both have same pH and therefore,
 $[\text{H}^+]$ by $\text{HCOOH} = [\text{H}^+]$ by CH_3COOH

$$\frac{C_1 \alpha_1}{\sqrt{K_{a1} C_1}} = \frac{C_2 \alpha_2}{\sqrt{K_{a2} C_2}}$$

$$\therefore \sqrt{(2.4 \times 10^{-4} \times 0.5)} = \sqrt{(1.8 \times 10^{-5} \times C)}$$

44. $C_{\text{CH}_3\text{COOH}} = 6.666 \text{ M}$
 $\text{HA} \rightleftharpoons \text{H}^+ + \text{A}^-; \quad K_a = 1.38 \times 10^{-4}$
 $\text{HB} \rightleftharpoons \text{H}^+ + \text{B}^-; \quad K_a = 1.05 \times 10^{-10}$

It is thus evident on account of low K_a values for HB, $[\text{H}^+]$ from HB is appreciably small in comparison to $[\text{H}^+]$ from HA and thus may be neglected.

For HA: $K_a = \frac{[\text{H}^+][\text{A}^-]}{[\text{HA}]}$

$$1.38 \times 10^{-4} = \frac{[\text{H}^+]^2}{[\text{HA}]}$$

$$[\text{H}^+]^2 = 1.38 \times 10^{-4} \times [\text{HA}]$$

$$= 1.38 \times 10^{-4} \times 0.03$$

Therefore, $[\text{H}^+] = [\text{A}^-] = 2.04 \times 10^{-3} \text{ M}$

Now for HB: $K_a = \frac{[\text{H}^+][\text{B}^-]}{[\text{HB}]}$

$$1.05 \times 10^{-10} = \frac{2.04 \times 10^{-3} \times [\text{B}^-]}{0.1}$$

(Since $[\text{H}^+]$ is provided by acid HA and HB is almost undissociated)

or $[\text{B}^-] = 5.15 \times 10^{-9} \text{ M}$

45. $\text{CH}_3\text{COOH} \rightleftharpoons \text{CH}_3\text{COO}^- + \text{H}^+$
 $\text{C}_6\text{H}_5\text{COOH} \rightleftharpoons \text{C}_6\text{H}_5\text{COO}^- + \text{H}^+$
 $[\text{H}^+] = \sqrt{K_{a1} \cdot C_1 + K_{a2} \cdot C_2}$
 $= \sqrt{1.8 \times 10^{-5} \times 0.02 + 6.4 \times 10^{-5} \times 0.01} = 1 \times 10^{-3}$
 $\therefore \text{pH} = 3$

Also $K_a = \frac{[\text{CH}_3\text{COO}^-][\text{H}^+]}{[\text{CH}_3\text{COOH}]}$

$$\therefore 1.8 \times 10^{-5} = \frac{[\text{CH}_3\text{COO}^-][1 \times 10^{-3}]}{0.02}$$

$$\therefore [\text{CH}_3\text{COO}^-] = 3.6 \times 10^{-4}$$

Similarly, $[\text{C}_6\text{H}_5\text{COO}^-] = 6.4 \times 10^{-4}$

46. Given, $[\text{HCOOH}] = 0.015 \text{ M}$
 $[\text{HCl}] = 0.02 \text{ M}$
 $\therefore [\text{H}^+] \text{ in solution} = 0.02 \text{ M}$

The dissociation of HCOOH is suppressed due to common ion effect in presence of HCl. The $[\text{H}^+]$ is provided by HCl in solution.



$$K_a = \frac{[\text{H}^+][\text{HCOO}^-]}{[\text{HCOOH}]}$$

$$1.8 \times 10^{-4} = \frac{[0.02][\text{HCOO}^-]}{[0.015]}$$

$$\therefore [\text{HCOO}^-] = 1.35 \times 10^{-4} \text{ M}$$

47. $\text{H}_2\text{S} \rightleftharpoons \text{H}^+ + \text{HS}^-; \quad K_{a1} = 10^{-7}$
 $\text{HS}^- \rightleftharpoons \text{H}^+ + \text{S}^{2-}; \quad K_{a2} = 1.3 \times 10^{-13}$
 $\text{HCl} \longrightarrow \text{H}^+ + \text{Cl}^-$

Due to common ion effect the dissociation of H_2S is suppressed and the $[\text{H}^+]$ in solution is due to HCl.

$$\therefore K_{a1} = \frac{[\text{H}^+][\text{HS}^-]}{[\text{H}_2\text{S}]}$$

$$10^{-7} = \frac{[0.3][\text{HS}^-]}{[0.1]} \quad [\because [\text{H}^+] \text{ from HCl} = 0.3]$$

$$\therefore [\text{HS}^-] = \frac{10^{-7} \times 0.1}{0.3} = 3.3 \times 10^{-8} \text{ M}$$

Further $K_{a2} = \frac{[\text{H}^+][\text{S}^{2-}]}{[\text{HS}^-]}$

$$1.3 \times 10^{-13} = \frac{[0.3][\text{S}^{2-}]}{3.3 \times 10^{-8}}$$

$$\therefore [\text{S}^{2-}] = \frac{1.3 \times 10^{-13} \times 3.3 \times 10^{-8}}{0.3} = 1.43 \times 10^{-20} \text{ M}$$

48. $\therefore$ Relative strength of weak acids = $\sqrt{\left(\frac{K_{a1} \times C_1}{K_{a2} \times C_2}\right)}$

Assume C_1 and C_2 are same (Although not given)

$$\therefore \text{Relative strength} = \sqrt{\left(\frac{K_{a1}}{K_{a2}}\right)} = \sqrt{\left(\frac{2.1 \times 10^{-4}}{1.1 \times 10^{-5}}\right)}$$

Relative strength for HCOOH to $\text{CH}_3\text{COOH} = 4.36:1$

49. (a) We have $\text{pH} = -\log K_a + \log \frac{[\text{Conjugate base}]}{[\text{Acid}]}$

$$\therefore [\text{Conjugate base}] = \frac{3 \times 1000}{82 \times 100} \text{ M and } [\text{Acid}] = \frac{2 \times 1000}{60 \times 100} \text{ M}$$

$$\therefore \text{pH} = -\log 1.8 \times 10^{-5} + \log \frac{\frac{3 \times 1000}{82 \times 100}}{\frac{2 \times 1000}{60 \times 100}} \quad \therefore \text{pH} = 4.7851$$

(b) $\text{pOH} = -\log K_b + \log \frac{[\text{Conjugate acid}]}{[\text{Base}]}$

$$\therefore \text{Total volume after mixing} = 250 + 5 = 255 \text{ mL}$$

$$\text{Meq. of conjugate acid} = 250 \times 0.1 = 25$$


$$\text{Meq. of base} = 5 \times 0.1 = 0.5$$

$$\therefore [\text{Conjugate acid}] = \frac{25}{255} \text{ and } [\text{Base}] = \frac{0.5}{255}$$

$$\therefore \text{pOH} = -\log 1.8 \times 10^{-5} + \log \frac{25/255}{0.5/255}$$

$$\text{pOH} = 6.4437 \quad \therefore \text{pH} = 14 - \text{pOH} = 7.5563$$

$$\begin{aligned} \text{(c) } \text{pH} &= -\log K_a + \log \frac{[\text{Anion}]}{[\text{Acid}]} \\ &= -\log 3.6 \times 10^{-4} + \log \frac{0.35/500}{0.25/500} \\ \text{pH} &= 3.5898 \end{aligned}$$



$$\begin{aligned} \therefore \text{pH} &= -\log K_a + \log \frac{[\text{CO}_3^{2-}]}{[\text{HCO}_3^-]} \\ \text{pH} &= -\log K_a + \log \frac{133}{150} = -\log 5.63 \times 10^{-11} + \log \frac{133}{150} \\ &= 10.249 - 0.052 \\ \text{pH} &= 10.197 \end{aligned}$$

51. [Pyridinium chloride] = $(0.15/500) \times 1000 = 0.3 \text{ M}$
[Pyridine] = 0.2 M

$\therefore$ A mixture of pyridine and its salt pyridinium chloride forms a basic buffer and therefore,

$$\begin{aligned} \text{pOH} &= -\log K_b + \log \frac{[\text{Conjugate acid}]}{[\text{Base}]} \\ \text{or } \text{pOH} &= -\log 1.5 \times 10^{-9} + \log (0.30/0.20) \\ &= -\log 1.5 + 9 \log 10 + \log 1.5 = 9 \\ \therefore [\text{OH}^-] &= 10^{-9} \end{aligned}$$

and $[\text{H}^+] = 10^{-5}$ So $\text{pH} = 5$

52. $\text{pOH} = -\log K_b + \log \frac{[\text{Conjugate acid}]}{[\text{Base}]}$

or $\text{pOH} = -\log K_b + \log \frac{[\text{NH}_4^+]}{[\text{NH}_4\text{OH}]}$

$\therefore [\text{NH}_4^+]$ is obtained from salt $(\text{NH}_4)_2\text{SO}_4$

$\therefore \text{pH} = 9.35 \therefore \text{pOH} = 14 - 9.35 = 4.65$

$\therefore$ Millimole of NH_4OH in solution = $0.2 \times 500 = 100$

Let millimole of NH_4^+ added in solution = a

$$\begin{aligned} \therefore [\text{NH}_4^+] &= \frac{a}{500}; \quad [\text{NH}_4\text{OH}] = \frac{100}{500} \\ 4.65 &= -\log 1.78 \times 10^{-5} + \log \frac{a/500}{100/500} \end{aligned}$$

$$4.65 = 4.7496 + \log \frac{a}{100} \therefore a = 79.51$$

$\therefore$ Millimole of $(\text{NH}_4)_2\text{SO}_4$ added = $\frac{a}{2} = \frac{79.51}{2} = 39.755$

$\therefore \frac{w}{132} \times 1000 = 39.755 \therefore w(\text{NH}_4)_2\text{SO}_4 = 5.248 \text{ g}$

53. Let V mL of 0.1 M HCOONa be mixed to 50 mL of 0.05 M HCOOH .

$\therefore$ [Molarity] = $\frac{\text{Total millimole}}{\text{Total volume}}$

$\therefore$ In mixture $[\text{HCOONa}] = \frac{0.1 \times V}{(V + 50)}$


$$[\text{HCOOH}] = \frac{50 \times 0.05}{V + 50}$$

$\therefore \text{pH} = -\log K_a + \log \frac{[\text{Conjugate base}]}{[\text{Acid}]}$

$\therefore 4.0 = 3.80 + \log \frac{(0.1 \times V)/(V + 50)}{2.5/(V + 50)}$

$\therefore V = 39.62 \text{ mL}$

54. $\text{NaCN} + \text{HCl}$ is not a buffer but if HCl is in less mass then, it gives a buffer as it produces HCN .




This is buffer of $\text{HCN} + \text{NaCN}$

Let a mole of HCl be used for this purpose

$$\begin{aligned} \therefore \text{pH} &= -\log K_a + \log \frac{0.01 - a}{a} \\ 8.5 &= -\log 4.1 \times 10^{-10} + \log \frac{0.01 - a}{a} \end{aligned}$$


$\therefore a = 8.85 \times 10^{-3} \text{ mol of HCl}$



Millimole before reaction	a	$1.0 \times 5 = 0.5$	0	0
Millimole after reaction	$(a - 0.5)$	0	0.5	0.5

$\therefore \text{pH} = 10.04 \quad \therefore \text{pOH} = -\log K_b + \log \frac{[\text{B}^+]}{[\text{BOH}]} \dots (1)$

$\therefore \text{pOH} = 3.96 \quad \therefore 3.96 = -\log K_b + \log \frac{0.5}{(a - 0.5)} \dots (2)$



Millimole before reaction	a	$0.1 \times 20 = 2$	0	0
Millimole after reaction	$(a - 0.2)$	0	2	2

$\therefore \text{pH} = 9.14 \quad \therefore \text{pOH} = -\log K_b + \log \frac{[\text{B}^+]}{[\text{BOH}]} \dots (3)$


$\therefore \text{pOH} = 4.86 \quad \therefore 4.86 = -\log K_b + \log \frac{2}{a - 2} \dots (4)$

Solving Eqs. (2) and (4), $K_b = 1.81 \times 10^{-5}$

56. For neutralization:

Total Meq. of acid = Meq. of base = $26.6 \times 0.1 = 2.66$

Now for partial neutralization of acid



The resultant mixture acts as a buffer and $[\text{HA}]$ and $[\text{BA}]$ may be placed in terms of Meq. since volume of mixture is constant.


$\therefore \text{pH} = -\log K_a + \log \frac{[\text{Conjugate base}]}{[\text{Acid}]}$

or $5 = -\log K_a + \log \frac{[1.2]}{[1.46]}$


$K_a = 8.219 \times 10^{-6}$

57. For complete neutralization,

Meq. of acid = Meq. of $\text{NaOH} = 36.12 \times 0.1 = 3.612$



Now 1.806 Meq. of HCl (18.06×0.1) are added to this solution containing 3.612 Meq. of NaA.





The solution has HA and NaA and thus acts as buffer

$$\therefore \text{pH} = -\log K_a + \log \frac{1.806}{1.806}$$

$$4.92 = -\log K_a$$

$$\therefore K_a = 1.2 \times 10^{-5}$$



For the dissociation of acid

$$K_a = 5 \times 10^{-2} = \frac{[\text{CHCl}_2 \text{COO}^-][\text{H}^+]}{[\text{CHCl}_2 \cdot \text{COOH}]}$$

$$\text{or } 0.05 = \frac{[0.1+x][x]}{[0.2-x]}$$

$$\therefore x = 0.05 \text{ or } [\text{H}^+] = 0.05$$

59. Initial pH of solution when,



$$[\text{NH}_3] = \frac{0.1}{1} \text{ and } [\text{NH}_4\text{Cl}] = \frac{0.1}{1}$$

$$\text{pOH} = -\log 1.8 \times 10^{-5} + \log \frac{[\text{Conjugate acid}]}{[\text{Base}]}$$

$$= -\log 1.8 \times 10^{-5} + \log \frac{0.1}{0.1}$$

$$\text{pOH} = 4.7447 \therefore \text{pH} = 9.2553$$

(i) Now 0.02 mole of HCl are added, then



$\therefore$ Volume = 1 litre

$$\therefore [\text{NH}_4\text{OH}] = \frac{0.08}{1} \text{ and } [\text{NH}_4\text{Cl}] = \frac{0.12}{1}$$



$$\therefore \text{pOH}_1 = -\log 1.8 \times 10^{-5} + \log \frac{0.12}{0.08}$$

$$\therefore \text{pOH}_1 = 4.9208 \therefore \text{pH}_1 = 9.0792$$

$$\text{Change in pH} = \text{pH} - \text{pH}_1 = 9.2553 - 9.0792 = +0.1761$$

$\therefore$ Change in pH = 0.1761 unit, i.e., pH decreases

(ii) Now 0.02 mole of NaOH are added




$$\therefore \text{pOH}_2 = -\log 1.8 \times 10^{-5} + \log \frac{0.08}{0.12}$$

$$\text{pOH}_2 = 4.5686 \therefore \text{pH}_2 = 9.4314$$

$$\text{Change in pH} = \text{pH} - \text{pH}_2 = 9.2553 - 9.4314 = -0.1761$$

$\therefore$ Change in pH = 0.1761 unit, i.e., pH increases





$$\therefore [\text{Molarity}] = \frac{\text{Millimole}}{\text{Total volume}}$$

$$\therefore [\text{CH}_3\text{COOH}] = \frac{6}{70} \quad [\text{CH}_3\text{COONa}] = \frac{4}{70}$$

$$\therefore \text{pH} = -\log 1.8 \times 10^{-5} + \log \frac{4/70}{6/70}$$

$$\text{pH} = 4.5686$$

(2) Let V mL of 0.2 M NaOH is required to make pH 4.74 then,



$$\therefore [\text{Acid}] = \frac{10 - 0.2V}{50 + V}; \quad [\text{Conjugate base}] = \frac{0.2V}{50 + V}$$

$$\therefore 4.74 = -\log 1.8 \times 10^{-5} + \log \frac{(0.2V)/(50+V)}{(10-0.2V)/(50+V)}$$

$$(\text{take } \log 1.8 \times 10^{-5} = 4.7447 = 4.74)$$

$$\therefore V = 25 \text{ mL}$$

61. Case I : pH when 1 mole CH_3COONa and 1 mole HCl are present.



$$\therefore [\text{CH}_3\text{COOH}] = 1 \text{ M}$$

$$\therefore [\text{H}^+] = C \cdot \alpha = C \cdot \sqrt{\frac{K_a}{C}} = \sqrt{K_a \cdot C} = \sqrt{K_a} \therefore C = 1$$

$$\therefore \text{pH}_1 = \frac{1}{2} \log K_a$$

Case II : pH when 1 mole CH_3COONa and 1 mole of CH_3COOH ; a buffer solution

$$\therefore \text{pH}_2 = -\log K_a + \log \frac{[\text{Conjugate base}]}{[\text{Acid}]} \quad \left| \begin{array}{l} \therefore [\text{Salt}] = [\text{Anion}] \\ = 1 \text{ M} \\ \therefore [\text{Acid}] = 1 \text{ M} \end{array} \right.$$

$$\text{pH}_2 = -\log K_a$$

$$\therefore \frac{\text{pH}_1}{\text{pH}_2} = \frac{1}{2}$$


62. For weak acid $\text{HA} : \alpha_{\text{HA}} = \frac{1}{100} = 0.01, [\text{HA}] = 0.1 \text{ M}$

$$\therefore K_a = C\alpha^2 = 0.1 \times (0.01)^2 = 10^{-5}$$

Now 0.2 M NaA, a salt of HA, is added to it resulting a buffer solution of $[\text{HA}] = 0.1 \text{ M}$ and $[\text{NaA}] = 0.2 \text{ M}$

$$\therefore \text{pH} = -\log 10^{-5} + \log \frac{0.2}{0.1}$$

$$\text{pH} = 5.3010$$



$\therefore [\text{A}^-]$ is provided by NaA since dissociation of HA in presence of NaA is suppressed due to a common ion effect

$$\therefore K_a = \frac{[\text{H}^+][\text{A}^-]}{[\text{HA}]} = \frac{(C \cdot \alpha) \times 0.2}{C(1 - \alpha)} = 10^{-5}$$

$$\therefore \alpha = 5 \times 10^{-5}$$


63. In 0.2 M HCOOH
- $[H^+] = 6.4 \times 10^{-3}$

$$C\alpha = 6.4 \times 10^{-3} \quad \therefore \alpha = 3.2 \times 10^{-2}$$

Now sodium formate is added and the dissociation will further be suppressed and therefore, new degree of dissociation (α_1) for HCOOH in presence of HCOONa is so small that it may be neglected, i.e.,

$\therefore$ [HCOOH] after dissociation = [HCOOH] before dissociation

$$\therefore [\text{HCOOH}] = 0.2$$



Conc. before dissociation	1	0	0
Conc. after dissociation	(1 - 0.75)	0.75	0.75


$$\text{So } [\text{HCOOH}] = 0.2 \therefore [\text{HCOO}^-] = 0.75$$

$$\therefore \text{pH} = -\log 2.4 \times 10^{-4} + \log \frac{0.75}{0.2} = 4.19$$

64. (a) Initially [Acetic acid] = 1 M

$$[\text{Acetate}] = 1 \text{ M}$$

Now 0.2 mole of HCl are added to it.



Mole before reaction	0.2	1	1	0
Mole after reaction	0	0.8	1.2	0.2


$$\therefore \text{Now } [\text{CH}_3\text{COOH}] = 1.2; [\text{CH}_3\text{COO}^-] = 0.8$$

$$\therefore \text{pH} = -\log 1.8 \times 10^{-5} + \log \frac{0.8}{1.2} = 4.5686$$

- (b) In II case initially [Acetic acid] = 0.1 M

$$[\text{Acetate}] = 0.1 \text{ M}$$

Now 0.2 mole of HCl are added to it



Mole of before reaction	0.2	0.1	0.1	0
Mole after reaction	0.1	0	0.2	0.1

$$\therefore [H^+] \text{ from free HCl} = 0.1 = 10^{-1} \therefore \text{pH} = 1$$

Note: CH_3COOH no doubt gives H^+ but being weak acid as well as in presence of HCl does not dissociate appreciably and thus, H^+ from CH_3COOH may be neglected.

$$65. \quad \text{pH} = -\log K_a + \log \frac{[\text{Conjugate base}]}{[\text{Acid}]}$$

Let a mol litre⁻¹ be concentration of salt, then concentration of acid = $(0.29 - a)$

$$4.4 = -\log 1.8 \times 10^{-5} + \log \frac{a}{(0.29 - a)} \therefore a = 0.09$$

$$\therefore [\text{Salt}] = 0.09 \text{ M}$$

$$[\text{Acid}] = 0.29 - 0.09 = 0.20 \text{ M}$$

$$66. \quad \text{pOH} = -\log K_b + \log \frac{[\text{Conjugate acid}]}{[\text{Base}]}$$

$$5 = 4.7 + \log \frac{a}{b}$$

$$\frac{a}{b} = 2 \therefore a = 2b$$

Given

$$a + b = 0.6$$

$$2b + b = 0.6$$


$$3b = 0.6$$

$\therefore$

$$\text{or } b = 0.2 \text{ mole} \quad \text{or } 0.2 \times 17 = 3.4 \text{ g/L}$$

$$\therefore a = 0.4 \text{ mole} \quad \text{or } 0.4 \times 53.5 = 21.4 \text{ g/L}$$

$$\text{Thus, } [\text{Salt}] = 0.4 \text{ M} \quad \text{and} \quad [\text{Base}] = 0.2 \text{ M}$$




mm at $t = 0$	30	40	0
mm after reaction	0	10	30

$$\therefore \text{pOH} = \text{p}K_b + \log \frac{[\text{NH}_4^+]}{[\text{NH}_3]} \quad (\text{p}K_a + \text{p}K_b = 14)$$

$$= 4.7448 + \log \frac{30}{10} = 5.2218$$

$$\therefore \text{pH} = 14 - 5.2218 = 8.7782$$



50 × 0.2	75 × 0.1	0	0
= 10	7.5	0	7.5
2.5	0	7.5	7.5

$$(\text{p}K_a + \text{p}K_b = 14)$$

$$\therefore \text{pOH} = 4.7448 + \log \frac{2.5}{7.5} = 4.2676$$

$$\therefore \text{pH} = 9.7324$$

$$69. \quad K_{b(x^{-1})} = 10^{-10}$$

Also for conjugate acid - base pair

$$K_{a(\text{HX})} \times K_{b(x^{-1})} = 10^{-14} \therefore K_{a(\text{HX})} = 10^{-4}$$

Now

$$\frac{[\text{HX}]}{(\text{acid})} = \frac{[X^-]}{(\text{anion})}$$

$$\therefore \text{pH} = -\log K_a + \log \frac{[\text{Conjugate base}]}{[\text{Acid}]} = -\log 10^{-4}$$

$$\text{pH} = 4$$

70. pH of buffer is given by:

$$\text{pH} = -\log K_a + \log \frac{[\text{Conjugate base}]}{[\text{Acid}]}$$

$$\text{Case I: } 4 = -\log 1.0 \times 10^{-5} + \log \frac{[\text{Conjugate base}]}{(0.5)}$$

$$\therefore \log \frac{[\text{Conjugate base}]}{0.5} = -1$$

$$[\text{Conjugate base}] = 0.1 \times 0.5 = 0.05 \text{ M}$$

$$\text{Case II: } 6 = -\log 1.0 \times 10^{-5} + \log \frac{[\text{Conjugate base}]}{0.5}$$

$$\therefore \log \frac{[\text{Conjugate base}]}{0.5} = 1$$

$$\therefore [\text{Conjugate base}] = [\text{Salt}] = 10 \times 0.5 = 5 \text{ M}$$

Now the two buffer [(I. NaA = 0.05 M and HA = 0.5 M) and (II. NaA = 5 M and HA = 0.5 M)] are mixed in equal proportion.


Thus, new conc. of NaA is mixed buffer

$$= \frac{0.05 \times V + 5 \times V}{2V} = \frac{5.05}{2}$$

$$\text{New conc. of HA in mixed buffer} = \frac{0.5 \times V + 0.5 \times V}{2V} = 0.5 \text{ M}$$

$$\text{Thus, } \text{pH} = -\log 1.0 \times 10^{-5} + \log \frac{[5.05/2]}{[0.5]}$$

$$\text{pH} = 5 + 0.7033 = 5.7033$$




$$\therefore K = \frac{[\text{A}^-]}{[\text{HA}][\text{OH}^-]} \quad \dots(1)$$

Also for weak acid HA: $\text{HA} \rightleftharpoons \text{H}^+ + \text{A}^-$

$$K_a = \frac{[\text{H}^+][\text{A}^-]}{[\text{HA}]} \quad \dots(2)$$

By Eqs. (2) and (1), $\frac{K_a}{K} = K_w$

$$\therefore K = \frac{K_a}{K_w} = \frac{10^{-4}}{10^{-14}} = 10^{10}$$



Volume of blood = 10 mL

$[\text{NaHCO}_3] = 5M$

Let volume of NaHCO_3 used = V mL


$$\therefore [\text{H}_2\text{CO}_3] \text{ in mixture} = \frac{2 \times 10}{(V+10)}$$

$$[\text{NaHCO}_3] \text{ in mixture} = \frac{(5 \times V)}{(V+10)}$$

$$\therefore \text{pH} = \text{p}K_a + \log \frac{[\text{Conjugate base}]}{[\text{Acid}]}$$


$$7.4 = -\log 7.8 \times 10^{-7} + \log \frac{(5 \times V)/(V+10)}{(2 \times 10)/(V+10)}$$

$$\therefore V = 78.36 \text{ mL}$$



$$C\alpha = 10^{-3} \quad \therefore \alpha = \frac{10^{-3}}{0.1} = 10^{-2}$$

Now $K_a = C\alpha^2 = 0.1 \times (10^{-2})^2 = 10^{-5}$




Before addition NaOH 0.1

After addition of NaOH $0.1 \times \frac{3}{4}$ $0.1 \times \frac{1}{4}$

when $\frac{1}{4}$ acid neutralizes

$$\therefore \text{pH} = -\log K_a + \log \frac{[\text{Conjugate base}]}{[\text{Acid}]}$$

$$\text{pH} = -\log 10^{-5} + \log \frac{0.1/4}{0.3/4} \quad \therefore \text{pH} = 4.5228$$




After addition of NaOH $0.1 \times 1/4$ $0.1 \times 3/4$

For $3/4$ neutralization

$$\therefore \text{pH} = -\log 10^{-5} + \log \frac{0.3/4}{0.1/4} \quad \therefore \text{pH} = 5.4771$$





Let solubility of AgCl be s mol litre⁻¹



$$K_{sp} = [\text{Ag}^+][\text{Cl}^-] = s \times s$$

$$\therefore s = \sqrt{K_{sp}} = \sqrt{1.5 \times 10^{-10}} \\ = 1.224 \times 10^{-5} \text{ mol litre}^{-1}$$





$$\therefore K_{sp} = [\text{Ag}^+][\text{Cl}^-] = (0.1+s)(s)$$

($\because s \ll 0.1$, presence of common ion decreases solubility)


$$\therefore s(0.1) = 1.5 \times 10^{-10}$$

$$\therefore s = 1.5 \times 10^{-9} \text{ mol litre}^{-1}$$

(c) Solubility of AgCl is more in (aq.) NaCl than AgNO_3 due to complex formation.



$$\text{In saturated solution } K_{sp} = [\text{Ag}^+][\text{Cl}^-] \quad \dots (i)$$



$$\text{In NaCl}(aq.) \quad K_f = \frac{[\text{AgCl}_2]^-}{[\text{Ag}^+][\text{Cl}^-]^2} \quad \dots (ii)$$

$$\text{By (i) and (ii)} \quad K_{sp} \times K_f = \frac{[\text{AgCl}_2]^-}{[\text{Cl}^-]} \quad \dots (iii)$$

Solubility of AgCl in pure water:

$$K_{sp} = [\text{Ag}^+][\text{Cl}^-] = [\text{Ag}^+]^2$$

$$\therefore [\text{Ag}^+] = \sqrt{K_{sp}} = \sqrt{1.0 \times 10^{-10}} \\ = 10^{-5} \text{ M}$$

In 0.01 M NaCl : By (iii)

$$[\text{AgCl}_2]^- = K_{sp} \times K_f \times [\text{Cl}^-] \\ = 1 \times 10^{-10} \times 3 \times 10^5 \times 0.01 \\ = 3 \times 10^{-7} \text{ M}$$



Let a mole of KBr be added into 1 litre of 0.05 M AgNO_3 to bring in precipitation.

$$\text{Thus, } [\text{KBr}] = \frac{a}{1}; [\text{AgNO}_3] = 0.05$$

$$\text{or } [\text{Br}^-] = a \text{ and } [\text{Ag}^+] = 0.05$$

$$\therefore [\text{Ag}^+][\text{Br}^-] = K_{sp}$$

$$0.05 \times a = 5.0 \times 10^{-13}$$

$$\therefore a = 5 \times 10^{-11} \text{ M} = 5 \times 10^{-11} \times 120$$

$$= 6.00 \times 10^{-9} \text{ g / 1000 mL}$$

Thus, mass of KBr needed for precipitation of AgBr from 500 mL 0.05 M are $3.0 \times 10^{-9} \text{ g}$



$$= \frac{1.79 \times 10^{-3}}{143.5} \text{ mol litre}^{-1}$$

$$= 1.247 \times 10^{-5} \text{ mol litre}^{-1}$$

$$\begin{aligned}\therefore K_{sp} &= [\text{Ag}^+][\text{Cl}^-] = s \times s \\ &= [1.247 \times 10^{-5}][1.247 \times 10^{-5}] \\ K_{sp} &= 1.55 \times 10^{-10} \text{ mol}^2 \text{ litre}^{-2}\end{aligned}$$




$$K_{sp} = [\text{Ag}^+][\text{Br}^-]$$

$$4 \times 10^{-13} = [1 \times 10^{-6}][\text{Br}^-]$$

$$\therefore [\text{Br}^-] = \frac{4 \times 10^{-13}}{1 \times 10^{-6}} = 4 \times 10^{-7} \text{ mol litre}^{-1}$$


78. For saturated solution of Ag_2CrO_4 , if solubility is s mol litre⁻¹. Then,



$$K_{sp} = (2s)^2(s) = 4s^3 \quad [\because [\text{Ag}^+] = 2s = 1.5 \times 10^{-4}]$$

$$= 4 \times (0.75 \times 10^{-4})^3 \quad \therefore s = 0.75 \times 10^{-4}$$

$$K_{sp} = 1.688 \times 10^{-12} \text{ mol}^3 \text{ litre}^{-3}$$




where s is solubility of PbI_2

$$\therefore K_{sp} = 4s^3 \quad \therefore s^3 = \frac{K_{sp}}{4}$$

$$\text{or } s = \sqrt[3]{\frac{K_{sp}}{4}} = \sqrt[3]{\frac{4 \times 10^{-9}}{4}} = 10^{-3}$$


$$\therefore [\text{I}^-] = 2s = 2 \times 10^{-3} \text{ mol/litre}$$



$$\text{Meq. of Ag}^+ = \text{Meq. of I}^-$$

$$N \times 10 = 2 \times 10^{-3} \times 25$$

$$\therefore N \text{ or } M \text{ of AgNO}_3 = 5 \times 10^{-3}$$



Millimole added 0.02V 0.0004V 0 0

Suppose V mL of both are mixed

$$\therefore [\text{Ca}^{2+}] = \frac{0.02V}{2V} \quad [\text{SO}_4^{2-}] = \frac{0.0004V}{2V}$$

$$\text{Thus, } [\text{Ca}^{2+}][\text{SO}_4^{2-}] \text{ in solution} < K_{sp}$$

$$2 \times 10^{-6} < 2.4 \times 10^{-5}$$


$\therefore \text{CaSO}_4$ will not precipitate.

81. For ZnS not to be precipitated from a solution of Zn^{2+} and Pb^{2+}

$$[\text{Zn}^{2+}][\text{S}^{2-}] < K_{sp} \text{ of ZnS}$$

$$[10^{-2}][\text{S}^{2-}] < 1.0 \times 10^{-21}$$

or the maximum $[\text{S}^{2-}] = 10^{-19}$ at which ZnS will begin to precipitate or upto this concentration, no precipitation will occur.




$$\therefore [\text{H}^+]^2[\text{S}^{2-}] = 1.1 \times 10^{-22}$$

$$\therefore [\text{H}^+]^2[10^{-19}] = 1.1 \times 10^{-22}$$

$$\therefore [\text{H}^+]^2 = 11 \times 10^{-4} \quad \therefore [\text{H}^+] = 3.3 \times 10^{-2} M$$

Thus, if $[\text{H}^+] = 3.3 \times 10^{-2}$ or slightly higher, the precipitation of ZnS will not take place and only PbS will precipitate.



Millimole added	100×0.1	1×1		
	= 10	= 1	0	0
Millimole left	9.5	0	0.5	1

$$\therefore \text{Concentration in } M = \frac{\text{mmole}}{\text{Total volume}}$$

$$\therefore [\text{Pb}^{2+}] = \frac{9.5 + 0.5}{101}$$

Now if PbCl_2 is precipitated, then contribution 0.5 of $[\text{Pb}^{2+}]$ from PbCl_2 should be left.

To see precipitation, ionic concentration product $> K_{sp}$


Ionic conc. product

$$= [\text{Pb}^{2+}][\text{Cl}^-]^2 = \left[\frac{10}{101} \right] \left[\frac{1}{101} \right]^2 = 9.70 \times 10^{-6}$$

which is greater than K_{sp} of PbCl_2 and thus, precipitation of PbCl_2 occurs.

$$\therefore [\text{Pb}^{2+}] = \frac{9.5}{101} = 9.4 \times 10^{-2} \text{ mol litre}^{-1}$$

83. Let solubility of PbBr_2 be s mol litre⁻¹



$\therefore$ Ionisation of $\text{PbBr}_2(s) = 80\%$

$$\therefore K_{sp} = [\text{Pb}^{2+}][\text{Br}^-]^2$$

$$8 \times 10^{-5} = \left[\frac{s \times 80}{100} \right] \left[\frac{2s \times 80}{100} \right]^2$$

$$\therefore s = 0.034 \text{ mol litre}^{-1} (\because \text{M. mass of PbBr}_2 = 367)$$

$$s = 0.034 \times 367 \text{ litre}^{-1}$$

$$s = 12.48 \text{ g litre}^{-1}$$

84. K_{sp} of $\text{Pb}(\text{OH})_2 = 4s^3 = 4 \times (6.7 \times 10^{-6})^3 = 1.203 \times 10^{-15}$

The buffer contains $\text{pH} = 8 \therefore \text{pOH} = 6$ or $[\text{OH}^-] = 10^{-6}$

Now left solubility of $\text{Pb}(\text{OH})_2$ be s mol litre⁻¹ in it.

$$\text{Thus, } [\text{Pb}^{2+}][\text{OH}^-]^2 = K_{sp}$$

$$[\text{Pb}^{2+}][10^{-6}]^2 = 1.203 \times 10^{-15}$$

$$\left[\begin{array}{l} \text{Buffer has pH} = 8; \therefore \text{pOH} = 6 \\ \text{and } [\text{OH}^-] = 10^{-6} \end{array} \right]$$

$$\therefore [\text{Pb}^{2+}] = \frac{1.203 \times 10^{-15}}{10^{-12}} = 1.203 \times 10^{-3} \text{ mol litre}^{-1}$$




$$= 2.5 \times 10^{-13}$$

In pure water $\text{S}^2 = 25 \times 10^{-14}$

$$\text{S} = 5 \times 10^{-7}$$

In pH = 3.19 ; pOH = 10.81 or $[\text{OH}^-] = 1.55 \times 10^{-11}$,
 $\text{C}_6\text{H}_5\text{COO}^-$ undergoes hydrolysis to form $\text{C}_6\text{H}_5\text{COOH}$.



$$\begin{aligned} \therefore K_H &= \frac{K_w}{K_a} = \frac{[\text{C}_6\text{H}_5\text{COOH}][\text{OH}^-]}{[\text{C}_6\text{H}_5\text{COO}^-]} \\ \therefore \frac{[\text{C}_6\text{H}_5\text{COOH}]}{[\text{C}_6\text{H}_5\text{COO}^-]} &= \frac{K_w}{K_a[\text{OH}^-]} \\ &= \frac{10^{-14}}{6.46 \times 10^{-5} \times 1.55 \times 10^{-11}} = 10 \end{aligned}$$

$[\text{Ag}^+]$ dissolved in pH 3.19 = $[\text{C}_6\text{H}_5\text{COO}^-]$ left after hydrolysis + $[\text{C}_6\text{H}_5\text{COOH}]$ formed due to hydrolysis

$$= [\text{C}_6\text{H}_5\text{COO}^-] + 10[\text{C}_6\text{H}_5\text{COO}^-] = 11[\text{C}_6\text{H}_5\text{COO}^-]$$

$$\therefore [\text{C}_6\text{H}_5\text{COO}^-] = \frac{[\text{Ag}^+]}{11}$$

$$[\text{Ag}^+][\text{C}_6\text{H}_5\text{COO}^-] = [\text{Ag}^+] \frac{[\text{Ag}^+]}{11} = 2.5 \times 10^{-13}$$

$$\therefore [\text{Ag}^+] = 1.658 \times 10^{-6}$$

i.e., solubility is $\frac{1.658 \times 10^{-6}}{5 \times 10^{-7}} = 3.316$ times greater than pure water.

$$86. [\text{Ca}^{2+}][\text{F}^-]^2 = 3.45 \times 10^{-11}$$

The F^- reacts with H^+ (pH = 3.0) to produce HF

$$K_{a\text{HF}} = \frac{[\text{H}^+][\text{F}^-]}{[\text{HF}]}$$

$$\text{or } 6.3 \times 10^{-4} = \frac{10^{-3} \times [\text{F}^-]}{[\text{HF}]}$$

$$\therefore \text{HF} = 1.58 \times [\text{F}^-]$$

Also the solution contains $[\text{HF}] + [\text{F}^-] = 2 \times [\text{Ca}^{2+}]$

$$\text{or } 1.58 \times [\text{F}^-] + [\text{F}^-] = 2 \times [\text{Ca}^{2+}]$$

$$\therefore [\text{F}^-] = \frac{2}{2.58} \times [\text{Ca}^{2+}] = 0.775 [\text{Ca}^{2+}]$$

Let solubility of CaF_2 be S mol litre $^{-1}$ $\therefore [\text{Ca}^{2+}] = S$

$$\therefore [\text{F}^-] = 0.775 \times S$$

$$\text{Thus } S \times (0.775 \times S)^2 = 3.45 \times 10^{-11}$$

$$\therefore S = 3.86 \times 10^{-4} \text{ M}$$

$$87. \text{pH} = 9 \therefore [\text{H}^+] = 10^{-9} \text{ M or } [\text{OH}^-] = 10^{-5} \text{ M}$$

Now if $\text{Mg}(\text{NO}_3)_2$ is present in a solution of $[\text{OH}^-] = 10^{-5} \text{ M}$, then,

Product of ionic conc.

$$\begin{aligned} &= [\text{Mg}^{2+}][\text{OH}^-]^2 = [0.001][10^{-5}]^2 \\ &= 10^{-13} \text{ lesser than } K_{sp} \text{ of } \text{Mg}(\text{OH})_2 \quad \text{i.e.,} \end{aligned}$$

$$8.9 \times 10^{-12}$$

$\therefore \text{Mg}(\text{OH})_2$ will not precipitate.


88. When $\text{Mg}(\text{OH})_2$ starts precipitation, then,

$$[\text{Mg}^{2+}][\text{OH}^-]^2 = K_{sp} \text{ of } \text{Mg}(\text{OH})_2$$

$$[0.1][\text{OH}^-]^2 = 1 \times 10^{-11}$$

$$\therefore [\text{OH}^-] = 10^{-5} \text{ M} \quad \therefore \text{pOH} = 5$$

$$\therefore \text{pH} = 14 - \text{pOH} \quad \therefore \text{pH} = 14 - 5 = 9$$



mm before reaction	10	20	0	0
	0	0	10	20

Thus, 10 mmole of $\text{Mg}(\text{OH})_2$ are formed. The product of


$$[\text{Mg}^{2+}][\text{OH}^-]^2 \text{ is therefore } \left[\frac{10}{200}\right] \times \left[\frac{20}{200}\right]^2 = 5 \times 10^{-4}$$

which is more than K_{sp} of $\text{Mg}(\text{OH})_2$. Now solubility (s) of $\text{Mg}(\text{OH})_2$ can be derived by

$$K_{sp} = 4s^3$$

$$\therefore s = \sqrt[3]{\frac{K_{sp}}{4}} = \sqrt[3]{\frac{1.2 \times 10^{-11}}{4}} = 1.4 \times 10^{-4}$$

$$\therefore [\text{OH}^-] = 2s = 2.8 \times 10^{-4}$$



mm added	7.5	excess	0	0
mm left	(7.5 - a)	excess	2a	a

$$\text{Given } [\text{Cl}^-] = \frac{0.0026}{35.5} = 7.32 \times 10^{-5}$$

$$\text{Also, conc. of } \text{Cl}^- \text{ formed} = \frac{\text{Millimole}}{\text{Volume in mL}} = \frac{2a}{5}$$

$$\therefore \frac{2a}{5} = \frac{0.0026}{35.5} \quad \therefore a = 1.83 \times 10^{-4} \text{ millimole}$$

$$\therefore \text{mmole of } \text{Na}_2\text{CO}_3 \text{ left in 5 mL} = 7.5 - 1.83 \times 10^{-4} = 7.5$$

$$\text{or } [\text{CO}_3^{2-}] = \frac{7.5}{5}$$

$$\text{Now } K_{sp\text{Ag}_2\text{CO}_3} = [\text{Ag}^+]^2[\text{CO}_3^{2-}]$$

$$\therefore [\text{Ag}^+]^2 = \frac{8.2 \times 10^{-12}}{7.5/5} = 5.46 \times 10^{-12}$$

$$\therefore [\text{Ag}^+] = 2.34 \times 10^{-6}$$


$$\therefore K_{sp} \text{ of } \text{AgCl} = [\text{Ag}^+][\text{Cl}^-] = 2.34 \times 10^{-6} \times \frac{0.0026}{35.5}$$

$$K_{sp} = 1.71 \times 10^{-19}$$

91. Let solubility of AgCNS and AgBr in a solution be a and b mol litre $^{-1}$ respectively.



$$[\text{Ion}] \text{ furnished on dissolution} \quad \begin{array}{cc} a & a \end{array}$$



$$[\text{Ion}] \text{ furnished on dissolution} \quad \begin{array}{cc} b & b \end{array}$$

$$\therefore [\text{Ag}^+] = a + b; \quad [\text{CNS}^-] = a; \quad [\text{Br}^-] = b$$

$$\therefore \text{For } \text{AgCNS } K_{sp\text{AgCNS}} = [\text{Ag}^+][\text{CNS}^-]$$

$$1 \times 10^{-12} = (a + b)(a) \quad \dots(1)$$

$$\text{For } \text{AgBr } K_{sp\text{AgBr}} = [\text{Ag}^+][\text{Br}^-]$$

$$5 \times 10^{-13} = (a + b)(b) \quad \dots(2)$$

By Eqs. (1) and (2),

$$\therefore \frac{a}{b} = \frac{10^{-12}}{5 \times 10^{-13}} = 2 \quad \text{or } a = 2b$$

$$\therefore \text{By Eq. (1), } (2b + b)(2b) = 1 \times 10^{-12} \therefore 6b^2 = 1 \times 10^{-12}$$

$$b = 4 \times 10^{-7} \text{ mol litre}^{-1}$$

$$\text{By Eq. (1), } (a + a/2)(a) = 1 \times 10^{-12}$$

$$\therefore a = 8.16 \times 10^{-7} \text{ mol litre}^{-1}$$

92. The $[I^-]$ needed for precipitation of Ag^+ and Hg_2^{2+} are derived as:

$$\text{For AgI: } [Ag^+][I^-] = K_{spAgI}$$

$$(0.1)[I^-] = 8.5 \times 10^{-17}$$

$$\therefore [I^-] = 8.5 \times 10^{-16} M \quad \dots(1)$$

$$\text{For } Hg_2I_2: [Hg_2^{2+}][I^-]^2 = 2.5 \times 10^{-26}$$

$$(0.1)[I^-]^2 = 2.5 \times 10^{-26}$$

$$\therefore [I^-] = 5 \times 10^{-13} M \quad \dots(2)$$

Since, $[I^-]$ required for precipitation of AgI is less and thus AgI begins to precipitate first. Also it will continue upto addition of $[I^-] = 5 \times 10^{-13}$ when Hg_2I_2 begins to precipitate and thus,

$$\text{Maximum } [I^-] \text{ for AgI precipitation} = 5 \times 10^{-13} M$$

Now at this concentration of I^- , $[Ag^+]$ left in solution is

$$[Ag^+]_{\text{left}}[I^-] = K_{spAgI}$$

$$\therefore [Ag^+]_{\text{left}} = \frac{8.5 \times 10^{-17}}{5.0 \times 10^{-13}} = 1.7 \times 10^{-4} M$$

$$\therefore 0.1 M Ag^+ \text{ will be left} = 1.7 \times 10^{-4} M Ag^+ \text{ in solution}$$

$$\therefore 100 \quad " \quad " \quad = 0.17\% M Ag^+$$

$$\therefore \% \text{ of Ag precipitated} = 99.83\%$$

93. The K_{sp} values of Ag_2CrO_4 and $AgIO_3$ reveals that CrO_4^{2-} and IO_3^- will be precipitated on addition of $AgNO_3$ as:

$$[Ag^+][IO_3^-] = 10^{-13}$$


$$[Ag^+]_{\text{needed}} = \frac{10^{-13}}{[0.005]} = 2 \times 10^{-11}$$

$$[Ag^+]^2[CrO_4^{2-}] = 10^{-8}$$


$$[Ag^+]_{\text{needed}} = \sqrt{\frac{10^{-8}}{0.1}} = 3.16 \times 10^{-4}$$

Thus, $AgIO_3$ will be precipitated first.

Now, in order to precipitate $AgIO_3$, one can show:



The left mole of $AgNO_3$ are now used to precipitate Ag_2CrO_4




Thus, $[CrO_4^{2-}]$ left in solution = 0.0975

Now, solution has $AgIO_3(s) + Ag_2CrO_4(s) + CrO_4^{2-}$ ions

$$\therefore [Ag^+]_{\text{left}} = \frac{K_{spAg_2CrO_4}}{[CrO_4^{2-}]} = \sqrt{\frac{10^{-8}}{0.0975}} = 3.2 \times 10^{-4} M$$

$$\therefore [IO_3^-]_{\text{left}} = \frac{K_{spAgIO_3}}{[Ag^+]} = \frac{10^{-13}}{3.2 \times 10^{-4}} = 3.2 \times 10^{-10} M$$

94. 500 mL of 0.4 M NaOH are mixed with 500 mL of $Ca(OH)_2$, a saturated solution having $Ca(OH)_2$ solubility as $s M$.



$$K_{sp} = s \times (2s)^2 = 4s^3$$

$$\text{Then, } 4s^3 = 4.42 \times 10^{-5}$$

$$\therefore s = \sqrt[3]{\frac{4.42 \times 10^{-5}}{4}} = 0.0223 M$$

Now $Ca(OH)_2 + NaOH$ are mixed

$\therefore$ Solution has Ca^{2+} and OH^- out of which some Ca^{2+} are precipitated

$$\text{On mixing, } [Ca^{2+}] = \frac{0.0223 \times 500}{1000}$$

$$= 0.01115 = 111.5 \times 10^{-4} M$$

$$[OH^-] = \frac{0.0223 \times 2 \times 500}{1000} + \frac{500 \times 0.4}{1000}$$

$$\text{(from } Ca(OH)_2 \text{) (from NaOH)}$$

$$= 0.2223 M$$

$$\therefore [Ca^{2+}][OH^-]^2 = K_{sp}$$

$$[Ca^{2+}]_{\text{left}}[0.2223]^2 = 4.42 \times 10^{-5}$$

$$[Ca^{2+}]_{\text{left}} = \frac{4.42 \times 10^{-5}}{[0.2223]^2} = 8.94 \times 10^{-4} \text{ mol litre}^{-1}$$

$\therefore$ Mole of $Ca(OH)_2$ precipitated = Mole of $[Ca^{2+}]$ precipitated

$$= 111.5 \times 10^{-4} - 8.94 \times 10^{-4} = 102.46 \times 10^{-4}$$

$\therefore$ Mass of $Ca(OH)_2$ precipitated from $Ca(OH)_2$ solution

$$= 102.46 \times 10^{-4} \times 74 = 7582.04 \times 10^{-4} g = 758.2 \text{ mg}$$

95. For $CaSO_4 \rightleftharpoons Ca^{2+} + SO_4^{2-}$

$$[Ca^{2+}][SO_4^{2-}] = K_{sp}$$

Let $[SO_4^{2-}] = a$, just sufficient to precipitate $CaSO_4$ from a solution having $[Ca^{2+}] = 0.005 M$

$$\text{Then, } [0.005][a] = 2.4 \times 10^{-5} \therefore a = \frac{2.4 \times 10^{-5}}{0.005}$$

$$[SO_4^{2-}] = 4.8 \times 10^{-3} \text{ mol litre}^{-1}$$

96. (a) $2NaOH + NiCl_2 \longrightarrow Ni(OH)_2 + 2NaCl$

$$\text{Mole before reaction } \frac{1.75}{40} \quad 0.25 \times 0.1$$

$$= 0.0438 \quad = 0.025$$

$$\text{Mole left } 0 \quad \left(0.025 - \frac{0.0438}{2}\right) \quad \frac{0.0438}{2} \quad 0.0438$$

$$\therefore \text{Mole of Ni(OH)}_2 \text{ formed} = \frac{0.0438}{2}$$

$$\therefore \text{mass of Ni(OH)}_2 \text{ formed} = \frac{0.0438}{2} \times 92.6 = 2.0279 \text{ g}$$

(b) Also for Ni(OH)_2 solution, $\text{Ni(OH)}_2 \rightleftharpoons \text{Ni}^{2+} + 2\text{OH}^-$

$$[\text{Ni}^{2+}][\text{OH}^-]^2 = K_{sp}$$

$$\left[\frac{0.025 - \frac{0.0438}{2}}{0.25} \right] [\text{OH}^-]^2 = 1.6 \times 10^{-14}$$

$$\therefore [\text{OH}^-] = 11.35 \times 10^{-7} \therefore \text{pOH} = 5.94 \therefore \text{pH} = 8.06$$

Note: Ni(OH)_2 formed will be precipitated out since maximum solubility of Ni(OH)_2 is $1.58 \times 10^{-5} \text{ M}$.

$$97. [\text{Ag}^+] \text{ in solution} = \sqrt{K_{sp} \text{AgCl}} = \sqrt{1 \times 10^{-10}} = 10^{-5} \text{ M}$$

Now the solution after mixing with NaBr has


$$[\text{Ag}^+] = \frac{10^{-5} \times 100}{200}$$

$$\text{and } [\text{Br}^-] = \frac{100 \times 0.03}{200}$$

$$\text{Thus, } [\text{Ag}^+][\text{Br}^-] = \left[\frac{10^{-5} \times 100}{200} \right] \times \left[\frac{100 \times 0.03}{200} \right] = 7.5 \times 10^{-8}$$

The product of ionic concentration is greater than K_{sp} and thus AgBr will be precipitated.

98. Let solubility of Mg(OH)_2 be $s \text{ mol litre}^{-1}$



$$\therefore [\text{Mg}^{2+}][\text{OH}^-]^2 = K_{sp}$$

$$4s^3 = 8.9 \times 10^{-12}$$

$$\therefore s = 1.305 \times 10^{-4} \text{ M}; \text{OH}^- = 2 \times 1.305 \times 10^{-4} \text{ M}$$

$$\therefore \text{pOH} = 3.5832$$

$$\text{pH} = 10.4168$$

99. $\text{CdSO}_4 + \text{HCl} + \text{H}_2\text{S} \longrightarrow \text{CdS} + \text{H}_2\text{SO}_4$

$$\begin{array}{ccc} \text{Millimole added} & 0.1 & 10 \times 0.08 \\ & & = 0.8 \end{array}$$

$$\begin{array}{ccc} \text{Millimole after reaction} & 0 & 0.8 \end{array}$$

$$\therefore \text{Millimole of H}^+ \text{ left} = 0.8 + 0.1 \times 2 = 1.0$$

(from HCl) (from H_2SO_4)

$$\text{Total volume} = 100 \text{ mL}$$

$$\therefore [\text{H}^+] = \frac{1}{100} = 10^{-2} \text{ M}$$

$$\therefore \text{pH} = 2$$

100. $[(\text{NH}_4)_2\text{S}] = 0.021 \text{ M}$

$$\therefore [\text{S}^{2-}] = 0.021 \text{ M}$$

$\therefore$ At equilibrium, $[\text{Zn}^{2+}][\text{S}^{2-}] = K_{sp}$ of ZnS

$$\therefore [\text{Zn}^{2+}] = \frac{4.51 \times 10^{-24}}{0.021} = 2.15 \times 10^{-22} \text{ M}$$

$$\therefore [\text{Zn}^{2+}] \text{ left in solution} = 2.15 \times 10^{-22} \times 65 \text{ g / litre}$$

$$\begin{aligned} &= \frac{2.15 \times 10^{-22} \times 65 \times 12}{1000} \text{ g / 12 mL} \\ &= 1.677 \times 10^{-22} \text{ g / 12 mL} \end{aligned}$$

101. For buffer solution,

$$\text{NH}_4\text{Cl} = 0.25 \text{ M and } \text{NH}_4\text{OH} = 0.05 \text{ M}$$

$$\text{pOH} = -\log K_b + \log \frac{[\text{Salt}]}{[\text{Base}]}$$

$$\therefore \text{pOH} = -\log 1.8 \times 10^{-5} + \log \frac{0.25}{0.05}$$

$$\therefore [\text{OH}^-] = 3.6 \times 10^{-6} \text{ M}$$

Now Al(OH)_3 and Mg(OH)_2 are stirred vigorously in it,

$$\therefore [\text{Al}^{3+}][\text{OH}^-]^3 = K_{sp} \text{ of } \text{Al(OH)}_3$$

$$[\text{Al}^{3+}][3.6 \times 10^{-6}]^3 = 6 \times 10^{-32}$$

$$\therefore [\text{Al}^{3+}] = 1.28 \times 10^{-15} \text{ M}$$

$$\text{Also } [\text{Mg}^{2+}][\text{OH}^-]^2 = K_{sp} \text{ Mg(OH)}_2$$

$$[\text{Mg}^{2+}][3.6 \times 10^{-6}]^2 = 8.9 \times 10^{-12}$$

$$\therefore [\text{Mg}^{2+}] = 0.686 \text{ M}$$



102. The minimum $[\text{OH}^-]$ at which there will be no precipitation of Mg(OH)_2 can be obtained by

$$K_{sp} = [\text{Mg}^{2+}][\text{OH}^-]^2$$

$$9.0 \times 10^{-12} = [0.05][\text{OH}^-]^2$$

$$\therefore [\text{OH}^-] = 1.34 \times 10^{-5} \text{ M}$$

Thus, a solution having $[\text{OH}^-] = 1.34 \times 10^{-5} \text{ M}$ will not show precipitation of Mg(OH)_2 in 0.05 M Mg^{2+} solution. These hydroxyl ions are to be derived by a buffer of NH_4Cl and NH_4OH , i.e.,



$$\text{For } \text{NH}_4\text{OH} \quad K_b = \frac{[\text{NH}_4^+][\text{OH}^-]}{[\text{NH}_4\text{OH}]}$$

In presence of NH_4Cl ; all the $[\text{NH}_4^+]$ are provided by NH_4Cl since common ion effect decreases dissociation of NH_4OH .

$$\therefore 1.8 \times 10^{-5} = \frac{[\text{NH}_4^+][1.34 \times 10^{-5}]}{[0.05]}$$

$$\therefore [\text{NH}_4^+] = 0.067 \text{ M or } [\text{NH}_4\text{Cl}] = 0.067 \text{ M}$$

103. Maximum solubility of CaSO_4 in water

$$S = \sqrt{K_{sp}} = 3 \times 10^{-3} \text{ mol litre}^{-1}$$

Let V litre of sample is taken, then CaSO_4 present

$$= \frac{131 \times V \times 10^{-3}}{10^6} \text{ g}$$

$$[\therefore \text{ppm} = \text{g of CaSO}_4 \text{ in } 10^6 \text{ g of sample}]$$

$$= 131 \times 10^{-3} V \text{ g} = \frac{131 \times 10^{-3} \times V}{136} \text{ mole in } V \text{ litre}$$

if water is evaporated on heating so that just deposition of CaSO_4 occurs. Let V_1 litre of water is left, then

$\frac{131 \times 10^{-3} \times V}{136}$ mole are present in V_1 litre solution are equal to $3 \times 10^{-3} \times V_1$ mole.

$$\therefore \frac{131 \times 10^{-3} \times V}{136} = 3 \times 10^{-3} \times V_1 \quad \therefore V_1 = 0.32V$$

Thus, volume evaporated = $V - 0.32V = 0.68V$ or **68% of water should be evaporated.**

104. Suppose V mL of solution contains $0.1 M \text{Mg}^{2+}$ and $0.8 M \text{NH}_4\text{Cl}$

Now V mL of ' a ' molarity NH_3 is added which just gives precipitate of $\text{Mg}(\text{OH})_2$, then

$$[\text{Mg}^{2+}][\text{OH}^-]^2 = K_{sp} \text{Mg}(\text{OH})_2$$

$$\left[\frac{0.1V}{2V}\right][\text{OH}^-]^2 = 1.4 \times 10^{-11}$$

$$\left([\text{Mg}^{2+}] = \frac{\text{Millimole}}{\text{Total volume}}\right)$$

$$\therefore [\text{OH}^-] = 1.67 \times 10^{-5} M$$

Now if the $[\text{OH}^-] = 1.67 \times 10^{-5}$, on addition of NH_3 in NH_4Cl , then $\text{Mg}(\text{OH})_2$ will precipitate. For buffer solution of NH_3 and NH_4Cl .

$$\therefore -\log[\text{OH}^-] = -\log K_b + \log \frac{[\text{NH}_4\text{Cl}]}{[\text{NH}_3]}$$

$$-\log 1.67 \times 10^{-5} = -\log 1.8 \times 10^{-5} + \log \frac{(0.8 \times V)/2V}{(a \times V)/2V}$$

$$\therefore a = 0.7421 M$$

$$\therefore [\text{NH}_3] \text{ in solution} = \frac{0.7421 \times V}{2V} = 0.3710 M$$

105. $[\text{Ca}^{2+}][\text{SO}_4^{2-}] = 2.4 \times 10^{-5}$

$$[\text{Sr}^{2+}][\text{SO}_4^{2-}] = 7.6 \times 10^{-7}$$

$$\therefore \frac{[\text{Sr}^{2+}]}{[\text{Ca}^{2+}]} = 0.03167$$

$$\text{Initial conc. of } \text{SO}_4^{2-} = \frac{10 \times 0.3}{30} = 0.1 M$$

$$\text{Initial conc. of } \text{Sr}^{2+} = \text{Initial conc. of } \text{Ca}^{2+} = \frac{0.1 \times 20}{30} = 0.0667 M$$

$$\text{Let final conc. of } \text{Ca}^{2+} = X, \therefore [\text{Sr}^{2+}] = 0.03167 X$$

$$\therefore \text{Change in conc. of } \text{SO}_4^{2-} = \text{change in conc. of } \text{Ca}^{2+} \text{ and } \text{Sr}^{2+}$$

$$= 0.0667 - X + 0.0667 - 0.03167 X$$

$$= 0.1334 - 1.03167 X$$

$$\therefore \text{Final } [\text{SO}_4^{2-}] = 0.1 - [0.1334 - 1.03167 X]$$

$$[\text{Using } [\text{Ca}^{2+}][\text{SO}_4^{2-}] = K_{sp} \text{ at equilibrium}]$$

$$\therefore X \cdot [0.1 - (0.1334 - 1.03167 X)] = 2.4 \times 10^{-5}$$

$$\therefore 1.03167 X^2 - 0.0334 X = 2.4 \times 10^{-5}$$

$$\therefore X = 3.3 \times 10^{-2} M \text{ or } [\text{Ca}^{2+}] = 3.3 \times 10^{-2} M$$

$$\therefore [\text{Sr}^{2+}] = 1.05 \times 10^{-3} M \therefore [\text{SO}_4^{2-}] = 7.17 \times 10^{-4} M$$

$$106. K_{sp} \text{ of } \text{CaCO}_3 = \left(\frac{7 \times 10^{-3}}{100}\right)^2 = 49 \times 10^{-10}$$

When only Ba^{2+} is 90% precipitated then only CaCO_3 starts precipitation then if solution contains a mol litre⁻¹ of Ca^{2+} and Ba^{2+} each,

$$[\text{Ca}^{2+}][\text{CO}_3^{2-}] = 49 \times 10^{-10}$$

$$[\text{CO}_3^{2-}] = \frac{49 \times 10^{-10}}{a}$$

Now for BaCO_3 ;

$$K_{sp} = [\text{Ba}^{2+}][\text{CO}_3^{2-}] = \frac{a \times 10}{100} \times \frac{49 \times 10^{-10}}{a}$$

$$= 4.9 \times 10^{-10}$$

107. (a) For $\text{Mg}(\text{OH})_2 + 2\text{NH}_4^+ \rightleftharpoons 2\text{NH}_3 + 2\text{H}_2\text{O} + \text{Mg}^{2+}$

$$\therefore K_C = \frac{[\text{NH}_3]^2 [\text{Mg}^{2+}]}{[\text{NH}_4^+]^2} = \frac{[\text{NH}_4\text{OH}]^2 [\text{Mg}^{2+}]}{[\text{NH}_4^+]^2} \quad \dots(1)$$


$$\text{Also } \text{NH}_4\text{OH} \rightleftharpoons \text{NH}_4^+ + \text{OH}^-$$

$$K_b = \frac{[\text{NH}_4^+][\text{OH}^-]}{[\text{NH}_4\text{OH}]} \quad \dots(2)$$

$$\therefore K_C \times K_b^2 = [\text{Mg}^{2+}][\text{OH}^-]^2 = K_{sp} \text{ of } \text{Mg}(\text{OH})_2$$

$$\therefore K_C = \frac{K_{sp}}{K_b^2} = \frac{6 \times 10^{-12}}{(1.8 \times 10^{-5})^2} = 1.85 \times 10^{-2}$$

(b) Let a mol litre⁻¹ of $\text{Mg}(\text{OH})_2$ be dissolved in presence of $0.5 M \text{NH}_4\text{Cl}$



Mole before reaction	0.5	0	0
Mole after reaction	$(0.5 - 2a)$	$2a$	a

$$\therefore K_C = \frac{a \times (2a)^2}{(0.5 - 2a)^2}$$

$$1.85 \times 10^{-2} = \frac{4a^3}{(0.5 - 2a)^2} \quad \dots(3)$$

Solving cubic equation,

$$\therefore a = 0.081 M$$

Note: Eq. (3) is cubic equation and its solution is not included in our subject. However if $2a$ is neglected in comparison to 0.5 , then ' a ' comes equal to $0.1049 M$.

108. $\text{Ag}_2\text{CO}_3 + \text{K}_2\text{C}_2\text{O}_4 \rightleftharpoons \text{Ag}_2\text{C}_2\text{O}_4 + \text{K}_2\text{CO}_3$

Mole before reaction	excess	0.1520	
Mole at equilibrium	$(0.1520 - 0.0358)$	0.0358	0.0358
		0.1162	

At equilibrium,

$$\therefore [\text{C}_2\text{O}_4^{2-}] = \frac{0.1162}{0.5} \quad [\text{CO}_3^{2-}] = \frac{0.0358}{0.5}$$

$$= 0.2324 M; \quad = 0.0716 M$$

$$\therefore [\text{Ag}^+]^2 [\text{C}_2\text{O}_4^{2-}] = \left[\frac{2 \times 0.0358}{0.5} \right]^2 \left[\frac{0.1520}{0.5} \right] = 6.23 \times 10^{-3}$$

$$\text{Thus, } [\text{Ag}^+]^2 [\text{C}_2\text{O}_4^{2-}] > K_{sp} \text{ of } \text{Ag}_2\text{C}_2\text{O}_4 \\ (1.29 \times 10^{-11})$$

$\therefore \text{Ag}_2\text{C}_2\text{O}_4$ will precipitate out.

$\therefore \text{Ag}_2\text{CO}_3$ is solid and $\text{Ag}_2\text{C}_2\text{O}_4$ is almost precipitated out. For $\text{Ag}_2\text{C}_2\text{O}_4$ precipitation.

$$[\text{Ag}^+]^2 [\text{C}_2\text{O}_4^{2-}] = K_{sp, \text{Ag}_2\text{C}_2\text{O}_4}$$


$$[\text{Ag}^+]^2 [0.2324] = 1.29 \times 10^{-11}$$

$$\therefore [\text{Ag}^+]^2 = \frac{1.29 \times 10^{-11}}{0.2324} = 5.55 \times 10^{-11}$$


Now for Ag_2CO_3

$$K_{sp} = [\text{Ag}^+]^2 [\text{CO}_3^{2-}] = [5.55 \times 10^{-11}] [0.0716] \\ = 3.97 \times 10^{-12} \text{ mol}^3 \text{ litre}^{-3}$$

109. Let solubility of AgCN be a mol litre⁻¹



However the CN^- formed will react with H^+ to form HCN



$$\therefore [\text{Ag}^+] = a \text{ and } [\text{HCN}] = a$$

Since HCN is weak acid and has low degree of dissociation. Also its dissociation is suppressed in presence of $[\text{H}^+]$.

Thus



$$\therefore \frac{[\text{CN}^-][\text{H}^+]}{[\text{HCN}]} = K_a$$

$$\text{or } [\text{CN}^-] = \frac{K_a [\text{HCN}]}{[\text{H}^+]} = \frac{a \times (4.8 \times 10^{-10})}{10^{-3}}$$

Now for $\text{AgCN}(s) + aq. \rightleftharpoons \text{Ag}^+ + \text{CN}^-$


$$K_{sp} = [\text{Ag}^+][\text{CN}^-]$$

$$1.2 \times 10^{-16} = \frac{a \times a \times 4.8 \times 10^{-10}}{10^{-3}}$$

$$\therefore a^2 = \frac{1.2 \times 10^{-16} \times 10^{-3}}{4.8 \times 10^{-10}}$$

$$\therefore a = 1.58 \times 10^{-5} \text{ mol litre}^{-1}$$

110. $\text{AgCl} + 2\text{NH}_3 \rightleftharpoons \text{Ag}(\text{NH}_3)_2^+ + \text{Cl}^-$



$$K_{sp} = [\text{Ag}^+][\text{Cl}^-] \quad \dots(1)$$

Given $\text{Ag}^+ + 2\text{NH}_3 \rightleftharpoons \text{Ag}(\text{NH}_3)_2^+$

$$K_f = \frac{[\text{Ag}(\text{NH}_3)_2^+]}{[\text{Ag}^+][\text{NH}_3]^2} \quad \dots(2)$$

$$\text{By Eqs. (1) and (2), } K_{sp} \times K_f = \frac{[\text{Ag}(\text{NH}_3)_2^+][\text{Cl}^-]}{[\text{NH}_3]^2}$$

$$\text{or } 1 \times 10^{-10} \times 1.6 \times 10^7 = \frac{a \times a}{[\text{NH}_3]^2}$$

Given solubility of $\text{AgCl} = 0.1M$

$$\therefore a = 0.1M \text{ for } \text{Ag}(\text{NH}_3)_2^+ \text{ and } \text{Cl}^-$$

$$\therefore [\text{NH}_3]^2 = \frac{0.1 \times 0.1}{1.6 \times 10^{-3}} = 6.25 \therefore [\text{NH}_3] = 2.5M$$

Also $0.2M \text{ NH}_3$ is needed to dissolve $0.1M \text{ Ag}^+$ ions, thus

$$[\text{NH}_3] = 2.5 \times 0.2 = 2.7M$$

111. $[\text{Ag}(\text{CN})_2]^- \rightleftharpoons \text{Ag}^+ + 2\text{CN}^-$

$$K = \frac{[\text{Ag}^+][\text{CN}^-]^2}{[\text{Ag}(\text{CN})_2]^-} = 4 \times 10^{-19} \quad \dots(1)$$

$$K_{sp} \text{AgCl} = [\text{Ag}^+][\text{Cl}^-] = 2.8 \times 10^{-10} \quad \dots(2)$$

$$\text{By Eq. (1) } 4 \times 10^{-19} = \frac{x \cdot 4x^2}{0.05}$$


$$\therefore x = [\text{Ag}^+] = 1.70 \times 10^{-7}$$

$$\therefore [\text{Ag}^+][\text{Cl}^-] = 1.70 \times 10^{-7} \times 0.02 = 3.41 \times 10^{-9} > K_{sp}$$

Thus, precipitation of AgCl will take place.

112. $\text{Ag}(\text{NH}_3)_2^+(aq.) \rightleftharpoons \text{Ag}^+(aq.) + 2\text{NH}_3(aq.)$

$$K_C = \frac{[\text{NH}_3]^2 [\text{Ag}^+]}{[\text{Ag}(\text{NH}_3)_2^+]} \quad \dots(1)$$



$$K_{sp} = [\text{Ag}^+][\text{Cl}^-] \quad \dots(2)$$

In case of simultaneous solubility Ag^+ remains same in solution.

Given $[\text{NH}_3] = 2a = 1M$; also $[\text{Ag}(\text{NH}_3)_2^+] = [\text{Cl}^-] = b$ because Ag^+ obtained from AgCl passes in $[\text{Ag}(\text{NH}_3)_2^+]$ state.



Thus, by Eqs. (1) and (2)

$$\frac{K_C}{K_{sp}} = \frac{[\text{NH}_3]^2}{[\text{Cl}^-][\text{Ag}(\text{NH}_3)_2^+]} = \frac{1}{b^2}$$


$$\text{or } b^2 = \frac{1.8 \times 10^{-10}}{62 \times 10^{-8}} = 0.29 \times 10^{-2}$$

$$\therefore b = 0.539 \times 10^{-1} = 0.0539$$

$$\text{or } [\text{Ag}(\text{NH}_3)_2^+] = 0.0539$$



$$\therefore K_{sp} = 2.2 \times 10^{-16} = [\text{Ag}^+][\text{CN}^-]$$



Now, soluble CN^- formed will hydrolyse as
 $\text{CN}^- + \text{H}_2\text{O} \rightleftharpoons \text{HCN} + \text{OH}^-$

$\therefore [\text{Ag}^+]$ of soluble $\text{AgCN} = \text{CN}^-$ left after hydrolysis + HCN formed after hydrolysis.

$$\therefore \left[\frac{2.2 \times 10^{-16}}{[\text{CN}^-]} \right] = [\text{CN}^-] + \frac{[\text{H}^+][\text{CN}^-]}{6.2 \times 10^{-10}}$$


$$= [\text{CN}^-] + \frac{10^{-2}[\text{CN}^-]}{6.2 \times 10^{-10}} \quad \left[\because \text{CN}^- \ll \frac{10^{-2}[\text{CN}^-]}{6.2 \times 10^{-10}} \right]$$

$$\text{or } [\text{CN}^-]^2 = \frac{2.2 \times 10^{-16} \times 6.2 \times 10^{-10}}{10^{-2}}$$

$$\therefore [\text{CN}^-] = 3.7 \times 10^{-12}$$


$$\therefore [\text{Ag}^+] = \frac{K_{sp}}{[\text{CN}^-]} = \frac{2.2 \times 10^{-16}}{3.7 \times 10^{-12}} = 5.96 \times 10^{-5} \text{ M}$$

114. Given, $\text{AgI}(s) \rightleftharpoons \text{Ag}^+(aq.) + \text{I}^-(aq.)$;
 $K_{sp} = [\text{Ag}^+][\text{I}^-] = 1.2 \times 10^{-17} \quad \dots(1)$



$$K_f = \frac{[\text{Ag}(\text{CN})_2]^-}{[\text{Ag}^+][\text{CN}^-]^2} = 7.1 \times 10^{19} \quad \dots(2)$$

Let x mole of AgI be dissolved in CN^- solution then,



By Eqs. (1) and (2), $K_{eq} = K_{sp} \times K_f$

$$K_{eq} = \frac{[\text{Ag}(\text{CN})_2]^- [\text{I}^-]}{[\text{CN}^-]^2} = 1.2 \times 10^{-17} \times 7.1 \times 10^{19}$$

$$K_{eq} = 8.52 \times 10^2 \quad \dots(3)$$


$$\therefore K_{eq} = 8.52 \times 10^2 = \frac{x \cdot x}{(1-2x)^2} = \frac{x^2}{(1-2x)^2}$$

$$\text{or } \frac{x}{1-2x} = 29.2$$

$$\text{Thus, } x = 29.2 - 58.4x \quad \text{or } x = 0.49 \text{ mol}$$

115. For $\text{Ag}_2\text{SO}_4 \rightleftharpoons 2\text{Ag}^+ + \text{SO}_4^{2-}$

$$K_{sp} = 4s^3 \quad \text{or } s = \sqrt[3]{\frac{K_{sp}}{4}} = \sqrt[3]{\frac{1.4 \times 10^{-5}}{4}} = 1.52 \times 10^{-2} \text{ M}$$



$$K_{sp} = s_1^2 \quad \text{or } s_1 = \sqrt{K_{sp}} = \sqrt{2.8 \times 10^{-13}} = 5.29 \times 10^{-7} \text{ M}$$

In solution, conc. of each ion can be given as:

$$\text{Thus, } [\text{Ag}^+] = \frac{2s \times 100}{350} = \frac{2 \times 1.52 \times 100}{350} = 0.869 \times 10^{-2} \text{ M}$$

$$[\text{SO}_4^{2-}] = \frac{s \times 100}{350} = \frac{1.52 \times 10^{-2} \times 100}{350} = 0.43 \times 10^{-2}$$

$$[\text{Pb}^{2+}] = \frac{s_1 \times 250}{350} = \frac{5.29 \times 10^{-7} \times 250}{350} = 3.78 \times 10^{-7}$$


$$[\text{CrO}_4^{2-}] = \frac{s_1 \times 250}{350} = \frac{5.29 \times 10^{-7} \times 250}{350} = 3.78 \times 10^{-7}$$

It is thus evident that

$$[\text{Ag}^+]^2 [\text{CrO}_4^{2-}] = (0.869 \times 10^{-2})^2 \times (3.78 \times 10^{-7})$$

$$= 2.85 \times 10^{-11} > K_{sp} \text{ Ag}_2\text{CrO}_4$$

The product is greater than K_{sp} of Ag_2CrO_4 and thus Ag_2CrO_4 will ppt.




Mole taken	2	0	0
Mole left	$(2-a)$	a	$2a$

where a is very-very small and thus assume that CaCO_3 is in soluble form

$$\text{Now, } K_{sp} \text{ of } \text{CaCO}_3 = x = [\text{Ca}^{2+}][\text{CO}_3^{2-}]$$

$$\text{Also } [\text{CO}_3^{2-}] = 2-a+a=2 \quad \therefore [\text{Ca}^{2+}] = \frac{x}{2}$$



$$K_{sp\text{CaF}_2} = [\text{Ca}^{2+}][\text{F}^-]^2$$

$$K_{sp\text{CaF}_2} = [y][2y]^2 = 4y^3$$


Further for $[\text{F}^-]$, we can have

$$[\text{F}^-] = [\text{F}^-] \text{ from } \text{CaF}_2 + [\text{F}^-] \text{ from } \text{NaF}$$

$$[\text{F}^-] = \sqrt{\frac{K_{sp\text{CaF}_2}}{[\text{Ca}^{2+}]}} + \text{Negligible value}$$

$$\therefore [\text{F}^-] = \sqrt{\frac{4y^3}{x/2}} = \sqrt{\frac{8y^3}{x}}$$

117. For PbI_2 saturated solution,



$$\therefore K_{sp} = 4s^3$$

$$\text{or } s = \sqrt[3]{\frac{K_{sp}}{4}} = \sqrt[3]{\frac{7.1 \times 10^{-9}}{4}} = 1.21 \times 10^{-3} \text{ M}$$

$$\therefore [\text{I}^-] \text{ in solution} = 2 \times 1.21 \times 10^{-3} \text{ M} = 2.42 \times 10^{-3} \text{ M}$$

Now, if AgNO_3 is used to neutralise these I^- ions

Then, Meq. of $\text{AgNO}_3 = \text{Meq. of I}^-$

$$13.3 \times N = 2.42 \times 10^{-3} \times 25$$

$$\therefore N_{\text{AgNO}_3} = 4.55 \times 10^{-3} = M_{\text{AgNO}_3}$$

For AgNO_3 normality = molarity

118. Meq. of $\text{C}_2\text{O}_4^{2-} = \text{Meq. of KMnO}_4$

$$= 0.00102 \times 6.3 \times 5 (\text{Mn}^{7+} + 5e \rightarrow \text{Mn}^{2+}) = 0.032$$

$$\therefore \text{Molarity of } \text{C}_2\text{O}_4^{2-} = \frac{\text{Meq.}}{\text{Volume in mL} \times \text{Valence factor}}$$

$$\text{or } s = \frac{0.032}{250 \times 2} \text{ M} = 6.4 \times 10^{-5} \text{ M} \quad (\text{C}_2^{3+} \rightarrow 2\text{C}^{4+} + 2e)$$

- $\therefore K_{sp} \text{ of } \text{CaC}_2\text{O}_4 = s^2 = (6.4 \times 10^{-5})^2$
 $= 4.09 \times 10^{-9}$
- 119.** Initial $[\text{Sr}^{2+}] = 16 \times 10^{-3} \text{ M}$
 Left $[\text{Sr}^{2+}] = 2.5 \times 10^{-3} \text{ M}$
 $\therefore [\text{Sr}^{2+}] \text{ precipitated} = (16 - 2.5) \times 10^{-3} = 13.5 \times 10^{-3} \text{ M}$
 $\therefore [\text{F}^-] \text{ needed for this precipitation} = 2 \times 13.5 \times 10^{-3} \text{ M}$
 $= 27.0 \times 10^{-3} \text{ M}$
 $(\therefore \text{Sr}^{2+} + 2\text{F}^- \longrightarrow \text{SrF}_2)$
 Also $[\text{Sr}^{2+}][\text{F}^-]^2 = K_{sp, \text{SrF}_2} = 2.8 \times 10^{-9}$
 $\therefore [\text{F}^-]^2 = \frac{2.8 \times 10^{-9}}{2.5 \times 10^{-3}}$
 $\therefore [\text{F}^-] = 1.058 \times 10^{-3} \text{ M}$, i.e., the concentration of F^- which will also appear in solution state.
 Thus, $[\text{F}^-] \text{ needed} = [27.0 + 1.058] \times 10^{-3} \text{ M}$
 $= 28.058 \times 10^{-3} \text{ M}$
 $\therefore \text{NaF needed for 1 litre} = 28.058 \times 10^{-3} \times 42 \text{ g}$
 $\therefore \text{NaF needed for 100 mL} = \frac{28.058 \times 10^{-3} \times 42}{10} \text{ g}$
 $= 0.1178 \text{ g}$

- 120.** Given, $\text{MBr}_2(\text{g}) \rightleftharpoons \text{MBr}_2(\text{aq.}) \longrightarrow \text{M}^{2+} + 2\text{Br}^-$
 $\text{MBr}_2 + \text{H}_2\text{S} \longrightarrow \text{MS} + 2\text{HBr}$
 $K_{sp} \text{ of MS} = [\text{M}^{2+}][\text{S}^{2-}]$
 $6 \times 10^{-21} = [0.05][\text{S}^{2-}]$
 $\therefore [\text{S}^{2-}] = 1.2 \times 10^{-19} \text{ M}$
 Thus, MS will be precipitated if H_2S provides $1.2 \times 10^{-19} \text{ M}$ ions of S^{2-} .
 Now for H_2S $\text{H}_2\text{S} \rightleftharpoons 2\text{H}^+ + \text{S}^{2-}$

$$K_1 \times K_2 = \frac{[\text{H}^+]^2[\text{S}^{2-}]}{[\text{H}_2\text{S}]}$$

$$10^{-7} \times 1.3 \times 10^{-13} = \frac{[\text{H}^+]^2 [1.2 \times 10^{-19}]}{[0.1]}$$

$$\therefore [\text{H}^+] = 1.04 \times 10^{-1} \text{ and } \text{pH} = 0.9826$$

- 121.** $\text{H}_2\text{S} \rightleftharpoons 2\text{H}^+ + \text{S}^{2-}$
 $\text{NaCN} \longrightarrow \text{Na}^+ + \text{CN}^-$
- | | | |
|-----|-----|-----|
| 0.2 | 0 | 0 |
| 0 | 0.2 | 0.2 |
- Also $\text{Ag}(\text{CN})_2^- \rightleftharpoons \text{Ag}^+ + 2\text{CN}^-$
 $K = \frac{[\text{Ag}^+][\text{CN}^-]^2}{[\text{Ag}(\text{CN})_2^-]} = \frac{[\text{Ag}^+] \times (0.2)^2}{0.02} = 1.0 \times 10^{-20}$
 $[\text{Ag}^+] = 5 \times 10^{-21} \text{ M}$

For $\text{Cd}(\text{CN})_4^{2-} \rightleftharpoons \text{Cd}^{2+} + 4\text{CN}^-$
 $K = \frac{[\text{Cd}^{2+}][\text{CN}^-]^4}{[\text{Cd}(\text{CN})_4^{2-}]}$

$$7.8 \times 10^{-18} = \frac{[\text{Cd}^{2+}] \times (0.2)^4}{0.02}$$

$$\therefore [\text{Cd}^{2+}] = 9.75 \times 10^{-17}$$

$$\text{Now, } [\text{Ag}^+]^2 \times [\text{S}^{2-}] = 1.0 \times 10^{-50}$$

$$\therefore [\text{S}^{2-}] \text{ needed to precipitate } \text{Ag}_2\text{S} = \frac{1.0 \times 10^{-50}}{(5 \times 10^{-21})^2}$$


$$= 4 \times 10^{-10} \text{ M}$$

$$[\text{S}^{2-}] \text{ needed to precipitate } \text{CdS} = \frac{7.1 \times 10^{-28}}{9.75 \times 10^{-17}}$$

$$= 7.28 \times 10^{-12} \text{ M}$$

Therefore, CdS will be precipitated first.

- 122.** For indicator dissociation equilibrium; being an acid



$$\therefore K_{\text{In}} = \frac{[\text{H}^+][\text{In}^-]}{[\text{HIn}]}$$

The midpoint of the colour range of an indicator HIn is the point at which $[\text{In}^-] = [\text{HIn}]$.

$$\therefore K_{\text{In}} = [\text{H}^+] = 1 \times 10^{-5}$$

$$\therefore [\text{H}^+] = 1 \times 10^{-5} \text{ or } \text{pH} = 5$$

Thus, at pH = 5 of the solution, the indicator will change its colour.

Again $K_{\text{In}} = \frac{[\text{H}^+][\text{In}^-]}{[\text{HIn}]}$

$$1 \times 10^{-5} = \frac{[\text{H}^+] \times 80 \times 100}{100 \times 20}$$

$$\therefore [\text{H}^+] = 0.25 \times 10^{-5} \therefore \text{pH} = 5.6020$$

- 123.** The two conditions when colour of indicator will be visible are derived by

$$\text{pH} = \text{p}K_a + \log \frac{[\text{In}^-]}{[\text{HIn}]}$$

(i) $\text{pH} = 5 + \log 10 = 6$

(ii) $\text{pH} = 5 + \log 0.1 = 4$

Thus, minimum change in pH = 2

- 124.** $\text{HBPh} \rightleftharpoons \text{H}^+ + \text{BPh}^-$

$$K_a = \frac{[\text{H}^+][\text{BPh}^-]}{[\text{HBPh}]}, \text{ when } \text{BPh}^- = \text{HBPh}, \text{ indicator will work. Thus}$$

$$[\text{H}^+] = 5.84 \times 10^{-5} \therefore \text{pH} = 4.2336$$

Also if $\text{pH} = 4.84$ or $[\text{H}^+] = 1.44 \times 10^{-5}$, then

$$K_a = \frac{[\text{H}^+][\text{BPh}^-]}{[\text{HBPh}]} \text{ or } 5.84 \times 10^{-5} = \frac{1.44 \times 10^{-5} \cdot C\alpha}{C(1-\alpha)}$$

$$\alpha = 0.8 \text{ or } 80\%$$

- 125.** $\text{NaOCN} + \text{H}_2\text{O} \rightleftharpoons \text{NaOH} + \text{HCN}$

$$h = \sqrt{\left(\frac{K_H}{C}\right)} = \sqrt{\left(\frac{K_w}{K_a C}\right)} = \sqrt{\frac{10^{-14}}{3.33 \times 10^{-4} \times 0.003}}$$

- $h = 10^{-4}$
 $\therefore \% \text{ hydrolysis} = 10^{-4} \times 100 = 10^{-2}$
126. $pK_a \text{ for HCN} = 14 - 4.70 = 9.30$
 $\text{NaCN} + \text{H}_2\text{O} \rightleftharpoons \text{NaOH} + \text{HCN}$

1	0	0
1-h	h	h

$$\therefore [\text{OH}^-] = C \cdot h = C \sqrt{\left(\frac{K_H}{C}\right)} = \sqrt{K_H \cdot C} = \sqrt{\left(\frac{K_w}{K_a} \cdot C\right)}$$

or $\text{pOH} = \frac{1}{2} [\text{p}K_w - \log C - \text{p}K_a]$

$$= \frac{1}{2} [14 + 0.3010 - 9.30] = 2.5$$

$$\therefore \text{pH} = 14 - 2.5 = 11.5$$

127. $\text{NH}_4\text{Cl} + \text{H}_2\text{O} \rightleftharpoons \text{NH}_4\text{OH} + \text{HCl}$

Before hydrolysis	1	0	0
After hydrolysis	(1-h)	h	h

where h is degree of hydrolysis

$$h = \sqrt{\left(\frac{K_H}{C}\right)} = \sqrt{\left(\frac{K_w}{K_b \cdot C}\right)}$$

$$= \sqrt{\frac{10^{-14}}{1.8 \times 10^{-5} \times 0.2}} = 5.27 \times 10^{-5}$$

From HCl, a strong acid

$$\therefore [\text{H}^+] = C \cdot h = C \sqrt{\left(\frac{K_H}{C}\right)} = \sqrt{K_H \cdot C} = \sqrt{\left(\frac{K_w}{K_b} \cdot C\right)}$$

$$= \sqrt{\left(\frac{10^{-14} \times 0.2}{1.8 \times 10^{-5}}\right)} = 1.054 \times 10^{-5}$$

$$\therefore \text{pH} = -\log[\text{H}^+] = -\log 1.054 \times 10^{-5} = 4.9771$$

Let conc. of NH_4Cl be C mol litre $^{-1}$

128. $\text{NH}_4\text{Cl} + \text{H}_2\text{O} \rightleftharpoons \text{NH}_4\text{OH} + \text{HCl}$

Before hydrolysis	1	0	0
After hydrolysis	(1-h)	h	h

$$\therefore [\text{H}^+] = \sqrt{K_H \cdot C} = \sqrt{\left(\frac{K_w}{K_b} \cdot C\right)} = \sqrt{\left(\frac{10^{-14} \times C}{1.8 \times 10^{-5}}\right)}$$

$$10^{-4.5} = \sqrt{\left(\frac{10^{-14} \times C}{1.8 \times 10^{-5}}\right)} \therefore C = 1.8 \text{ mol litre}^{-1}$$

$$\therefore \text{Mass of } \text{NH}_4\text{Cl} = 1.8 \times 53.5 \text{ g / litre}^{-1}$$

$$= 1.8 \times 53.5 \times \frac{1}{2} \text{ g / 500 mL} = 48.15 \text{ g}$$

129. (a) $\text{NaBu} + \text{H}_2\text{O} \rightleftharpoons \text{NaOH} + \text{BuH}$

Conc. before hydrolysis	1	0	0
Conc. after hydrolysis	1-h	h	h

$$\therefore [\text{OH}^-] = C \cdot h = C \sqrt{\left(\frac{K_H}{C}\right)} = \sqrt{K_H \cdot C} = \sqrt{\left(\frac{K_w}{K_a} \cdot C\right)}$$

$$[\text{OH}^-] = \frac{10^{-14} \times 0.2}{2 \times 10^{-5}} = \sqrt{10^{-10}} = 10^{-5}$$

$$\therefore \text{pOH} = 5$$



Also $\text{pH} + \text{pOH} = 14 \therefore \text{pH} = 14 - 5 = 9$

(b) [Ans : 8]

130. $\text{Asc}^- + \text{H}_2\text{O} \rightleftharpoons \text{HAsc} + \text{OH}^-$
- $$\therefore [\text{OH}^-] = C \cdot h = C \sqrt{\left(\frac{K_H}{C}\right)} = \sqrt{K_H \cdot C} = \sqrt{\left(\frac{K_w}{K_a} \cdot C\right)}$$
- $$= \sqrt{\left(\frac{10^{-14} \times 0.02}{5 \times 10^{-5}}\right)} = 2 \times 10^{-6}$$
- $$\therefore [\text{H}^+] = \frac{1 \times 10^{-14}}{2 \times 10^{-6}} = 5 \times 10^{-9}$$
- Also $h = \sqrt{\left(\frac{K_H}{C}\right)} = \sqrt{\left(\frac{K_w}{K_a \times C}\right)}$
- $$= \sqrt{\left(\frac{10^{-14}}{5 \times 10^{-5} \times 0.02}\right)} = 10^{-4} \text{ or } 0.01\%$$
131. Let V mL of acid and V mL of NaOH be used. Conc. of both acid and NaOH are same.
- $$\therefore \text{CH}_3\text{COOH} + \text{NaOH} \rightarrow \text{CH}_3\text{COONa} + \text{H}_2\text{O}$$
- | | | | | |
|-----------------------|---------------------------|---------------------------|---------------------------|---------------------------|
| Conc. before reaction | $\frac{0.1 \times V}{2V}$ | $\frac{0.1 \times V}{2V}$ | 0 | 0 |
| Conc. after reaction | 0 | 0 | $\frac{0.1 \times V}{2V}$ | $\frac{0.1 \times V}{2V}$ |
- $$\therefore [\text{CH}_3\text{COONa}] = \frac{0.1}{2} = 0.05 \text{ M}$$
- Now calculate pH by hydrolysis of CH_3COONa
- $$\text{CH}_3\text{COONa} + \text{H}_2\text{O} \rightleftharpoons \text{CH}_3\text{COOH} + \text{NaOH}$$
- $$[\text{OH}^-] = C \cdot h = C \sqrt{\left(\frac{K_H}{C}\right)} = \sqrt{\left(\frac{K_w \times C}{K_a}\right)}$$
- $$= \sqrt{\left(\frac{10^{-14} \times 0.05}{1.9 \times 10^{-5}}\right)} = 5.12 \times 10^{-6}$$
- $$\therefore \text{pOH} = 5.29 \therefore \text{pH} = 8.71$$
132. [Ans : 8.975]
133. $\text{Ca}(\text{LaC})_2 + 2\text{H}_2\text{O} \rightleftharpoons \text{Ca}(\text{OH})_2 + 2\text{HLac}$
- or $2\text{LaC}^- + 2\text{H}_2\text{O} \rightleftharpoons 2\text{OH}^- + 2\text{HLac}$
- | | | | |
|-------------------|-------|---|---|
| Before hydrolysis | 1 | 0 | 0 |
| After hydrolysis | (1-h) | h | h |
- $$\therefore [\text{Ca}(\text{LaC})_2] = \frac{0.13}{0.5} = 0.26 \text{ M}$$
- $$\therefore [\text{LaC}^-] = 0.26 \times 2 = 0.52 \text{ M}$$
- $$\therefore 1 \text{ mole } \text{Ca}(\text{LaC})_2 \text{ gives } 2 \text{ moles } (\text{LaC}^-)$$
- Now $[\text{OH}^-] = C \cdot h = C \sqrt{\left(\frac{K_H}{C}\right)}$
- $$= \sqrt{K_H \cdot C} = \sqrt{\left(\frac{K_w \times C}{K_a}\right)}$$
- where C is conc. of anion which undergoes hydrolysis
- $$\therefore 10^{-5.60} = \sqrt{\left(\frac{10^{-14} \times 0.52}{K_a}\right)} \therefore K_a = 8.25 \times 10^{-4}$$
134. $\text{K}_3\text{PO}_4 + \text{H}_2\text{O} \rightleftharpoons \text{K}_2\text{HPO}_4 + \text{KOH}$
- or $\text{PO}_4^{3-} + \text{H}_2\text{O} \rightleftharpoons \text{HPO}_4^{2-} + \text{OH}^-$
- Since hydrolysis proceeds only in 1 step


$$\therefore [\text{OH}^-] = C \cdot h = C \sqrt{\left(\frac{K_w}{K_a \cdot C}\right)} = \sqrt{\left(\frac{K_w \cdot C}{K_a}\right)}$$

K_a is III dissociation constant of acid H_3PO_4



$$\therefore [\text{OH}^-] = \sqrt{\left(\frac{10^{-14} \times 0.1}{1.3 \times 10^{-12}}\right)} \quad \therefore \text{pOH} = 1.5634$$

$$\therefore \text{pH} = 12.4366$$




$$[\text{H}^+] = Ch$$

and $K_H = \frac{C \cdot h^2}{1-h}$ and $h = \frac{5}{100}$


Thus, $6.5 \times 10^{-3} = \frac{C \times 5 \times 5 \times 100}{100 \times 100 \times 95}$

$$\therefore C = 2.47 \text{ M} \quad \therefore [\text{H}^+] = 2.47 \times \frac{5}{100} \quad \therefore \text{pH} = 0.9083$$



$$\therefore [\text{H}^+] = C \cdot h = C \sqrt{\left(\frac{K_H}{C}\right)} = \sqrt{(K_H \cdot C)}$$

$$= \sqrt{10^{-9} \times 0.001} = 10^{-6} \quad \therefore \text{pH} = 6$$



Before	1	1	0	0
hydrolysis				
After	1-h	1-h	h	h
hydrolysis				

Let concentration of salt be C mol litre⁻¹

$$\therefore K_H = \frac{[\text{Aniline}][\text{Acetic acid}]}{[\text{Aniline}^+][\text{Acetate}^-]} = \frac{C \cdot h \cdot C \cdot h}{C \cdot (1-h) \cdot C \cdot (1-h)}$$

$$\therefore K_H = \frac{h^2}{(1-h)^2} \quad \therefore \frac{h}{1-h} = \sqrt{(K_H)}$$

$$\frac{h}{1-h} = \sqrt{\left(\frac{K_w}{K_a \cdot K_b}\right)} = \sqrt{\left(\frac{1.008 \times 10^{-14}}{1.75 \times 10^{-5} \times 3.83 \times 10^{-10}}\right)}$$

$$\therefore h = 54.95\%$$

Note: If $K_H = h^2$ is used assuming $1-h \approx 1$ the value of h comes greater than 1 which is not possible and thus, it is always advised in all such cases, not to assume $1-h \approx 1$ when h values are higher. Also the term dissociation constant of water is used here for ionic product of water. See the value.


138. The pH of salt HCOONH_4 (a salt of weak acid + weak base) is given by:



$$\text{pH} = \frac{1}{2} [\log K_b - \log K_a - \log K_w]$$

$$\therefore \text{pH} = \frac{1}{2} [\text{p}K_a + \text{p}K_w - \text{p}K_b]$$

$$\text{pH} = \frac{1}{2} [3.8 + 14 - 4.8] = 6.5$$



mM before	50 × 0.1	50 × 0.05		
reaction	= 5	= 2.5	0	0
mM after	2.5	0	2.5	2.5
reaction				

$\therefore$ Strong alkali is left free and thus, it will decide pH of solution.

$$\therefore [\text{NaOH}] = \frac{2.5}{100} = 2.5 \times 10^{-2} \text{ M}$$

$$\therefore [\text{OH}^-] = 2.5 \times 10^{-2} \text{ M} \quad \therefore \text{pOH} = 1.6021$$

$$\therefore \text{pH} = 12.3979$$



mM before	50 × 0.05	50 × 0.1		
reaction	= 2.5	= 2.5	0	0
mM after	0	2.5	2.5	2.5
reaction				

The solution consists CH_3COOH and CH_3COONa and thus, is a buffer

$$\therefore \text{pH} = -\log K_a + \log \frac{[\text{CH}_3\text{COONa}]}{[\text{CH}_3\text{COOH}]}$$

$$\therefore [\text{CH}_3\text{COOH}] = \frac{2.5}{100} \text{ M}; \quad [\text{CH}_3\text{COONa}] = \frac{2.5}{100} \text{ M}$$


$$\therefore \text{pH} = -\log 1.8 \times 10^{-5} + \log \frac{2.5/100}{2.5/100} \quad \therefore \text{pH} = 4.7447$$



mM before	50 × 0.1	50 × 0.1		
reaction	= 5	= 5	0	0
mM after	0	0	5	5
reaction				

$\therefore \text{CH}_3\text{COONa}$ is left with concentration = $\frac{5}{100} \text{ M}$ and

thus, pH is decided by salt hydrolysis




Before hydrolysis	1	0	0
After hydrolysis	1-h	h	h

$$\therefore [\text{OH}^-] = Ch = C \sqrt{\left(\frac{K_H}{C}\right)} = \sqrt{\left(\frac{K_w \cdot C}{K_a}\right)}$$

$$= \sqrt{\left(\frac{10^{-14}}{1.8 \times 10^{-5}} \times \frac{5}{100}\right)}$$

$$= 5.27 \times 10^{-6} \text{ M}$$

$$\therefore \text{pH} = 8.7218$$



mM before reaction	50 × 0.1	50 × 0.05		
	= 5	= 2.5	0	0
mM after reaction	2.5	0	2.5	2.5


The solution contains NH_4OH and NH_4Cl and thus, acts as buffer

$$\text{Thus, } \text{pOH} = -\log K_b + \log \frac{[\text{NH}_4\text{Cl}]}{[\text{NH}_4\text{OH}]}$$

$$\therefore [\text{NH}_4\text{Cl}] = \frac{2.5}{100} \text{ M}; \quad [\text{NH}_4\text{OH}] = \frac{2.5}{100}$$


$$\therefore \text{pOH} = -\log 1.8 \times 10^{-5} + \log \frac{2.5/100}{2.5/100}$$

$$pOH = 4.7447 \therefore pH = 9.2553$$



$\therefore$ Strong acid is left and therefore, pH is decided by HCl.

$$[HCl] = [H^+] = \frac{2.5}{100} M \therefore pH = 1.6021$$



$\therefore$ Salt is left and thus, pH will be decided by salt hydrolysis.


$$[NH_4Cl] = \frac{5}{100} = 5 \times 10^{-2} M$$

Since NH_4Cl is salt of strong acid and weak base and therefore,

$$pH = \frac{1}{2} [\log K_b - \log C - \log K_w]$$


$$= \frac{1}{2} [\log 1.8 \times 10^{-5} - \log 5 \times 10^{-2} - \log 1 \times 10^{-14}]$$

$$\therefore pH = 5.2782$$




$\therefore$ Salt is left and thus, pH will be decided by salt hydrolysis.

$$[CH_3COONH_4] = \frac{2.5}{100} = 2.5 \times 10^{-2} M$$



$$CH_3COOH = Ch$$

Now for dissociation of CH_3COOH



$$\therefore K_a = \frac{[CH_3COO^-][H^+]}{[CH_3COOH]}$$

$\therefore \alpha$ is small and thus, $[CH_3COOH] = Ch$

Also $[CH_3COO^-]$ is provided from CH_3COONH_4 left unhydrolysed in solution, i.e., $[CH_3COO^-] = C(1-h)$

$$\therefore K_a = \frac{C(1-h)[H^+]}{Ch} \quad \therefore h \ll 1$$

$$\text{or } [H^+] = hK_a \quad \therefore 1-h=1$$

$$= \sqrt{\left(\frac{K_w}{K_a \cdot K_b}\right)} \times K_a = \sqrt{\left(\frac{K_w K_a}{K_b}\right)}$$

$$\therefore pH = +\frac{1}{2} [\log K_b - \log K_w - \log K_a]$$

$$= \frac{1}{2} [pK_w + pK_a - pK_b]$$

$$= +\frac{1}{2} [\log 1.8 \times 10^{-5} - \log 10^{-14} - \log 1.8 \times 10^{-5}]$$

$$pH = 7$$


140. We have from Raoult's law

$$\frac{P^0 - P_s}{P_s} = \frac{w \times M}{m \times W}$$

$$\therefore \frac{w}{m \times W} = \frac{17.540 - 17.536}{17.536 \times 18} = 1.267 \times 10^{-5}$$

$$\therefore \text{Molality} = \frac{w}{m \times W} \times 1000 = 1.267 \times 10^{-5} \times 10^3$$

$$= 1.267 \times 10^{-2} = \text{Molarity} = \text{Conc. of } [BOH]$$




Molarity is also given as 1×10^{-2}

$$\therefore \frac{\text{Exp. value of molarity}}{\text{Normal value of molarity}} = 1 + \alpha$$

$$\therefore \frac{1.267 \times 10^{-2}}{1 \times 10^{-2}} = 1 + \alpha \quad \therefore \alpha = 0.267$$

$$\text{Now, } K_b = \frac{C\alpha^2}{(1-\alpha)} = \frac{0.01 \times 0.267 \times 0.267}{(1-0.267)} = 9.74 \times 10^{-4}$$

141.





$$\therefore \pi V = nST(1+\alpha) \quad \text{or } 1+\alpha = \frac{\pi \times V}{n \times S \times T}$$

$$\therefore 1+\alpha = \frac{0.293}{0.01 \times 0.0821 \times 298} \quad \left(\frac{n}{V} = 0.01\right)$$

$$\therefore \alpha = 1.197 - 1 = 0.197$$


$$\therefore K_a \text{ for } HA = \frac{C\alpha^2}{(1-\alpha)} = \frac{0.01 \times (0.197)^2}{(1-0.197)} = 4.83 \times 10^{-4}$$

Similarly, K_a for $HB = 2.85 \times 10^{-3}$



$$K_f = \frac{[Zn(OH)_4]^{2-}}{[OH^-]^2}$$

Let the solubility be s in change (i) and thus solubility will also be s in change (ii) By coupling (i) and (ii),



$$\therefore K_{sp} \times K_f = s \times s$$

$$\text{or } s = \sqrt{K_{sp} \times K_f} = \sqrt{1.2 \times 10^{-17} \times 0.12} = 1.2 \times 10^{-9} M$$

Thus total solubility, will be $2s = 2 \times 1.2 \times 10^{-9}$

$$= 2.4 \times 10^{-9} M$$

This solubility is minimum when $[Zn^{2+}][OH^-]^2$

$$= 1.2 \times 10^{-17}$$


$$1.2 \times 10^{-9} [OH^-]^2 = 1.2 \times 10^{-17}$$

$$\therefore [OH^-] = 10^{-4} M$$

Note: If only I change than solubility of

$$Zn(OH)_2 = \sqrt[3]{\frac{K_{sp}}{4}} = \sqrt[3]{\frac{1.2 \times 10^{-17}}{4}} = 1.44 \times 10^{-6} m$$

and at this conc. $[\text{OH}^-] = 2 \times 144 \times 10^{-6} \text{ M} = 2.88 \times 10^{-6} \text{ M}$



Let a mole of N_2O_4 and b mole of NO_2 were present in equilibrium mixture

$$\therefore (a+b) = \frac{PV}{RT} = \frac{753 \times 0.5}{760 \times 0.0821 \times 298} = 0.020 \quad \dots(1)$$

$$K_p = \frac{(n_{\text{NO}_2})^2}{(n_{\text{N}_2\text{O}_4})} \times \left[\frac{P}{[n_{\text{NO}_2} + n_{\text{N}_2\text{O}_4}]} \right]^1$$

$$\therefore 0.113 = \frac{b^2}{a} \times \left[\frac{753}{760 \times (a+b)} \right]$$

$$\therefore \frac{b^2}{a(a+b)} = 0.114 \quad \dots(ii)$$

By Eqs. (i) and (ii)

$$\therefore \frac{b^2}{a} = 0.114 \times 0.020 = 2.3 \times 10^{-3} \quad \dots(iii)$$

$$b^2 = 2.3 \times 10^{-3} a = 2.3 \times 10^{-3} \times (0.02 - b)$$


$$\text{or } b^2 + 0.0023b - 4.6 \times 10^{-5} = 0$$

$$\therefore b = \frac{-0.0023 \pm \sqrt{(0.0023)^2 + 4 \times 4.6 \times 10^{-5}}}{2 \times 1}$$

$$= \frac{-0.0023 \pm \sqrt{1.90 \times 10^{-4}}}{2}$$


$$b = 5.73 \times 10^{-3}$$

$$\therefore \text{By Eq. (i)} \quad a = 0.014$$



$$\text{Total NO}_2 \text{ mole} = 5.73 \times 10^{-3} + 2 \times \text{moles of N}_2\text{O}_4$$

$$\begin{aligned} &\text{from NO}_2 \\ &= 5.73 \times 10^{-3} + 2 \times 0.014 \\ &= 0.0337 \end{aligned}$$



$$\therefore [\text{HNO}_2] = \frac{0.0337 \times 1000}{2 \times 250} = 0.0674 \text{ M};$$

$$[\text{H}^+] = \frac{0.0337 \times 1000}{2 \times 250} = 0.0674 \text{ M}$$

Due to common ion effect (H_3O^+ furnished by HNO_3), the dissociation of HNO_2 is suppressed.

$$K_a = \frac{[\text{H}^+][\text{NO}_2^-]}{[\text{HNO}_2]} = \frac{0.0674 \times [\text{NO}_2^-]}{0.0674} = 4.5 \times 10^{-4}$$

$$\therefore [\text{NO}_2^-] = 4.5 \times 10^{-4}$$

$$\text{Also } \text{pH} = -\log[\text{H}^+] = -\log 0.0674 = 1.17$$

(c) $\pi = \text{C.S.T.} \times (\text{No. of particles present in solution})$

$$= 0.0674 \times 0.0821 \times 298 \times 3$$

(2NO_2 furnishes three particles)


$$= 4.95 \text{ atm}$$

(d) Eq. of CaO required = Eq. of HNO_2 + Eq. of HNO_3

$$= \frac{0.0337}{2} \times 1 + \frac{0.0337}{2} \times 1 = 0.0337$$

$$\therefore \text{Mole of CaO required} = \frac{0.0337}{2} \text{ or } \frac{w}{56} = \frac{0.0337}{2}$$

$$\therefore w_{\text{CaO}} = 0.924 \text{ g}$$



$$K_{eq} = \frac{[\text{CH}_3\text{COO}^-][\text{HCN}]}{[\text{CH}_3\text{COOH}][\text{CN}^-]} \quad \dots(i)$$




For CH_3COOH and HCN in same solution

$$K_1 = \frac{[\text{CH}_3\text{COO}^-][\text{H}^+]}{[\text{CH}_3\text{COOH}]} \quad \dots(ii)$$

$$K_2 = \frac{[\text{CN}^-][\text{H}^+]}{[\text{HCN}]} \quad \dots(iii)$$

By Eqs. (ii) and (iii)

$$K_{eq} = \frac{K_1}{K_2} = \frac{1.8 \times 10^{-5}}{4.9 \times 10^{-10}} = 3.674 \times 10^4$$



Thus acidic character order for acids is: $\text{HX} > \text{HY} > \text{HZ}$

Also for NaY : $[\text{OH}^-] = 10^{-5}$ $\therefore \text{pH} = 9 \therefore \text{pOH} = 5$

$$C_h = 10^{-5}$$


$$C \cdot \sqrt{\frac{K_H}{C}} = 10^{-5} \quad \text{or} \quad \sqrt{\frac{K_w \times C}{K_a}} = 10^{-5}$$

$$\therefore \frac{10^{-14} \times 0.1}{K_a} = 10^{-10} \quad \therefore K_a \text{ for HY} = 10^{-5}$$

Similarly, K_a for HZ = 10^{-9}


146. The conjugate base (here $\text{HC}_8\text{H}_4\text{O}_4^-$) of polybasic acid (here $\text{H}_2\text{C}_8\text{H}_4\text{O}_4$), is amphoteric and can act as either an acid or a base because it can donate its remaining acidic hydrogen atom or accept an acidic hydrogen atom and revert to the original acid.

Acid:



$$K_{a_2} = \frac{[\text{H}_3\text{O}^+][\text{C}_8\text{H}_4\text{O}_4^{2-}]}{[\text{HC}_8\text{H}_4\text{O}_4^-]} \quad \dots(i)$$

Hydrolysis:



$$K_H = \frac{[\text{H}_2\text{C}_8\text{H}_4\text{O}_4][\text{OH}^-]}{[\text{HC}_8\text{H}_4\text{O}_4^-]} \quad \dots(ii)$$

Also for $\text{H}_2\text{C}_8\text{H}_4\text{O}_4$



$$K_{a_1} = \frac{[\text{H}^+][\text{HC}_8\text{H}_4\text{O}_4^-]}{[\text{H}_2\text{C}_8\text{H}_4\text{O}_4]} \quad \dots(iii)$$

By Eqs. (ii) and (iii)

$$K_H \times K_{a_1} = [\text{H}^+][\text{OH}^-] = K_w \quad \dots(\text{iv})$$

Assuming $\text{HC}_8\text{H}_4\text{O}_4^-$ having low degree of dissociation as well as low degree of hydrolysis, acidic nature of $\text{HC}_8\text{H}_4\text{O}_4^-$ and hydrolysis of $\text{HC}_8\text{H}_4\text{O}_4^-$ producing $\text{C}_8\text{H}_4\text{O}_4^{2-}$ and $\text{H}_2\text{C}_8\text{H}_6\text{O}_4$ of almost equal conc.

$$[\text{H}_2\text{C}_8\text{H}_6\text{O}_4] = [\text{C}_8\text{H}_4\text{O}_4^{2-}] \quad \dots(\text{v})$$

$$\text{From Eqs. (i), (ii) and (v)} \quad \frac{K_H}{K_{a_2}} = \frac{[\text{OH}^-]}{[\text{H}^+]} \quad \dots(\text{vi})$$

$$\text{Also} \quad K_w = [\text{H}^+][\text{OH}^-] \quad \dots(\text{vii})$$

By Eqs. (vi) and (vii)

$$\therefore K_w = \frac{[\text{H}^+] \times K_H \times [\text{H}^+]}{K_{a_2}} \quad \dots(\text{viii})$$

$$\text{By Eqs. (iv) and (viii)} \quad K_H \times K_{a_1} = \frac{[\text{H}^+]^2 \times K_H}{K_{a_2}}$$

$$\text{or } [\text{H}^+]^2 = K_{a_1} \times K_{a_2} \quad \text{or } [\text{H}^+] = \sqrt{K_{a_1} \times K_{a_2}}$$

$$\text{pH} = \frac{1}{2}[\text{p}K_{a_1} + \text{p}K_{a_2}]$$

$$\frac{1}{2}[2.94 + 5.44] = 4.19$$

Note: 1. The pH is independent of salt concentration in case of acidic salt solution.


2. All polyprotic acids (except H_2SO_4) are weak acids and the relation $\text{pH} = \frac{1}{2}[\text{p}K_{a_1} + \text{p}K_{a_2}]$ holds good.

3. The relation $\text{pH} = \frac{1}{2}[\text{p}K_{a_1} + \text{p}K_{a_2}]$ is derived by using reasonable assumptions and doing some intricate algebra. The assumptions include that if $\text{p}K_{a_2}$ is high HA^- is weak acid, the solution will have high pH. If $\text{p}K_{a_1}$ is large HA^- is likely to act as 'strong' weak base. Thus, if both $\text{p}K_{a_1}$ and $\text{p}K_{a_2}$ are large, pH will be high.

147. $[\text{H}^+]$ formed due to electrolyte oxidation = 4 m moles


$$\begin{array}{ccc} \text{Na}_2\text{HPO}_4 & + & \text{Na}_3\text{PO}_4 \\ \text{Initial mm} & 0.08 \times 100 & 0.02 \times 100 \\ & = 8 & 2 \end{array}$$

$[\text{H}^+]$ will be used by PO_4^{3-} to give H_2PO_4^-



Thus, a new buffer containing Na_2HPO_4 (8 mm) and H_2PO_4^- (2 mm) will remain in solution

$$\therefore \text{pH} = \text{p}K_{a_{\text{H}_2\text{PO}_4^-}} + \log \frac{[\text{Na}_2\text{HPO}_4]}{[\text{H}_2\text{PO}_4^-]} = 7.2 + \log \frac{8}{2} = 7.81$$



$$\text{at } 1/3 \text{ neutralisation } \frac{2}{3} \quad \frac{1}{3}$$

$$\therefore [\text{Conjugate base}] = [\text{CH}_3\text{COO}^-] = \frac{1}{3}$$

$$[\text{Acid}] = [\text{CH}_3\text{COOH}] = \frac{2}{3}$$

$$\therefore \text{pH}_1 = -\log K_a + \log \frac{1/3}{2/3} = -\log K_a - \log 2$$

$$\text{at } 2/3 \text{ neutralisation } [\text{conjugate base}] = \frac{2}{3}$$

$$[\text{Acid}] = \frac{1}{3}$$

$$\therefore \text{pH}_2 = -\log K_a + \log \frac{2/3}{1/3}$$

$$= -\log K_a + \log 2$$

$$\therefore \Delta\text{pH} = \text{pH}_1 - \text{pH}_2 = \log \frac{1}{2} - \log 2$$


$$= -2 \log 2$$

149. (3)

$$\Lambda_m(\text{HX}) = \frac{x}{10} \quad \Lambda_m(\text{HY}) = x. \text{ Also } \alpha = \frac{\Lambda_m}{\Lambda_0}. \text{ Let } \alpha_1$$

and α_2 be degree of dissociation of 0.01 M HX and 0.1 M HY

$$\therefore \frac{\alpha_1}{\alpha_2} = \frac{\Lambda_m(\text{HX})/\Lambda_m^0(\text{HX})}{\Lambda_m(\text{HY})/\Lambda_m^0(\text{HY})} = \frac{(x/10)/\Lambda_m^0(\text{HX})}{x/\Lambda_m^0(\text{HY})} = \frac{1}{10}$$



$$0.01 \quad \quad \quad - \quad \quad -$$

$$0.01(1-\alpha_1) \quad 0.01\alpha_1 \quad 0.01\alpha_1$$

$$K_{a_1} = 0.01\alpha_1^2 \quad (\because \alpha_1 \ll 1)$$



$$0.1 \quad \quad \quad - \quad \quad -$$

$$0.1(1-\alpha_2) \quad 0.1\alpha_2 \quad 0.1\alpha_2$$

$$K_{a_2} = 0.1\alpha_2^2 \quad (\because \alpha_2 \ll 1)$$

$$\frac{K_{a_1}}{K_{a_2}} = \frac{0.01\alpha_1^2}{0.1\alpha_2^2} = \frac{1}{10} \cdot \frac{\alpha_1^2}{\alpha_2^2} = \frac{1}{10} \times \left(\frac{1}{10}\right)^2 = \frac{1}{1000}$$

$$\log K_{a_1} - \log K_{a_2} = -3$$

$$\text{p}K_{a_1} - \text{p}K_{a_2} = 3$$

● SINGLE INTEGER ANSWER PROBLEMS ●

- 500 mL of HCl of pH 4.3010 is allowed to react with 500 mL of pH 9.6990 NaOH solution at 300 K. If temperature remains constant, approximate pH of resulting solution is....
- 300 mL of $5 \times 10^{-2} M$ HCl is allowed to react with 200 mL of $5 \times 10^{-2} M$ NaOH at 300 K. The pH of resulting solution at 300 K is.....
- 500 mL of 0.01 M CH_3COOH + 500 mL of 0.02 M CH_3COONa gives a pH equal to 5.3010. The pK_b of CH_3COO^- is.....
- K_b for MeOH an indicator is 10^{-9} . The pH at which it shows is a colour change is.....
- K_{sp} of $M(\text{OH})_2$ is 5×10^{-16} at 25°C . The pH of its saturated solution at 25°C is
- The equilibrium constant of a strong acid (HA) with weak base BOH is 10^{11} . The pH of 0.10 M solution of BA is.....
- 10 mL of 0.25 M H_2SO_4 is completely neutralised by 0.125 M solution of NH_3 . The pH of the solution at the equivalence point is....., if K_b for $\text{NH}_3 = 10^{-5}$.
- 100 mL of 0.20 M weak acid HA is completely neutralised by 0.20 M NaOH. K_b for A^- is 10^{-3} , the pOH at the equivalence point is.....
- One litre of 1 M solution of an acid HA ($K_a = 10^{-4}$ at 25°C) has pH = 2. It is diluted by water so that new pH becomes double. The solution was diluted to 5×10^a mL. The value of a is.....
- K of pure water is 10^{-12} at 60°C . The pH of pure water at 60°C is.....
- An aqueous solution of 0.24 M aniline ($K_b = 4.166 \times 10^{-10}$) is mixed with NaOH solution to maintain anilinium ion concentration to $1 \times 10^{-8} M$. The pOH of NaOH solution used was
- A certain buffer solution equals concentration of X^- and HX. K_b for X^- is 10^{-10} . The pH of buffer is.....
- Two weak acids HA and HB have same pH when their concentration ratio is 3 : 1. The ratio of the dissociation constants of HB and HA is.....
- Conjugate base of a weak acid has $K_b = 10^{-9}$. The equilibrium constant for the reaction of acid with strong base is.....
- K_{sp} of SrF_2 is 1×10^{-10} . The solubility of SrF_2 in 0.1 M NaF is $1 \times 10^{-a} M$. The value of a is.....
- pH of 0.1 M NaCl solution is.....
- The solubility of $\text{RNH}_2(\text{g})$ in water at 1 atm and 27°C is 24.63 litre per litre water. If pK_b of $\text{RNH}_2 = 4$, the maximum pOH of the resulting solution will be
- If K_{sp} of HgSO_4 is 6.4×10^{-5} mole² litre², then solubility of HgSO_4 in mole/m³ is
- The dissociation constant of a monoprotic acid, morphine a pain killer is 10^{-a} . Its 0.01 M solution with its 0.01 M sodium salt shows a pH 6, the value of pK_a is.....
- The molar concentration of SO_4^{2-} required to precipitate RaSO_4 for 2×10^{-4} mole of Ra^{2+} in 500 mL is $10^{-a} M$. If K_{sp} of RaSO_4 is 4×10^{-11} then value of a is.....
- The A^{2+} left in solution when 50.0 mL of 0.20 M $\text{A}(\text{NO}_3)_2$ is added to 50 mL of 1.5 M NaCl is $9.95 \times 10^{-a} M$. If K_{sp} of $\text{A}(\text{Cl})_2$ is 5.6×10^{-4} then the value of a is.....
- The solubility of CoS in 0.10 M H_2S and 0.15 M H^+ solution is $a \times 10^{-6} M$. The value of a is..... Given K for $\text{H}_2\text{S} \rightleftharpoons 2\text{H}^+ + \text{S}^{2-}$ is 1×10^{-21} and K_{sp} CoS = 3.11×10^{-26}
- If K_{a1} and K_{a2} for H_2S are 1×10^{-7} and 1×10^{-14} respectively. Then hydrolysis constant for $\text{S}^{2-} + \text{H}_2\text{O} \rightleftharpoons \text{HS}^- + \text{OH}^-$ is.....
- The pK_{eq} for $A^- + \text{H}_3\text{O}^+ \rightleftharpoons \text{HA} + \text{H}_2\text{O}$ is -6, then pK_{eq} for HA is.....
- pH of 0.1 M NaCl solution is.....
- A buffer mixture containing equal mole of HB and its conjugate base B^- ($K_b = 10^{-10}$) has pH equal to.....
- The pH at which an indicator $pK_b = 5.00$ changes its colour is.....
- Ethylenediamine $\text{NH}_2\text{C}_2\text{H}_4\text{NH}_2$ a base (B) can add 1 or 2 protons. The successive pK_b values are 4 and 7 respectively. The $p^{[\text{BH}_2^{2+}]}$ in 0.001 M solution of ethylenediamine is.....
- The per cent error in $[\text{H}_3\text{O}^+]$ made by neglecting the ionisation of water in a $1.0 \times 10^{-6} M$ NaOH solution.
- The pH of solution resulting when 50 mL of 0.20 M HCl is mixed with 50 mL of 0.20 M CH_3COOH ($pK_a = 1.8 \times 10^{-5}$)
- The pK_{eq} for the reaction of Na_2A and HCl at exactly half neutralisation of Na_2A is 6, then pK_{eq} for $A^{2-} + \text{H}^+ \rightleftharpoons \text{HA}^-$ will be.....
- If K_{eq} for $A^- + \text{H}_3\text{O}^+ \rightleftharpoons \text{HA} + \text{H}_2\text{O}$ is 10^6 , then pK_a for acid HA is.....
- The solubility of $M(\text{OH})_x$ is $10^{-4} M$. If its pK_{sp} is 4×10^{-12} , then the value of x is.....
- An acid indicator ($K_a = 10^{-5}$) will have 50% its basic form at pOH.
- The solubility product of AgCl is $10^{-10} M^2$ at 25°C . The minimum volume of water (in m³) required to dissolve 1.435 g AgCl at 25°C .

36. The molar solubility of Ag_2CO_3 ($K_{sp} = 4 \times 10^{-13}$) in $0.1 \text{ M Na}_2\text{CO}_3$ is 10^{-a} M . The value of a is
37. The pH of an aqueous solution of $0.1 \text{ M NH}_4\text{Cl}$ and $0.01 \text{ M NH}_4\text{OH}$ ($pK_b = 5$) is
38. The pOH of resulting 500 mL solution by mixing 25 mL of 0.1 M NaOH and 25 mL 0.08 M HCl , finally diluted with water is
39. If $a \text{ M}$ solution of weak acid HA has same pH as $b \text{ M}$ solution of weak acid HB . Also if $\frac{a}{b} = 4$, then the ratio of dissociation constant of two acids HB and HA will be
40. The minimum pH required to prevent the precipitation of ZnS in a solution having 0.1 M ZnCl_2 and $0.1 \text{ M H}_2\text{S}$ if $K_{a1} \times K_{a2}$ for H_2S is 10^{-20} and K_{sp} of $\text{ZnS} = 10^{-21}$.
41. The per cent error in $[\text{H}^+]$ made by neglecting ionisation of water in $1.0 \times 10^{-6} \text{ M NaOH}$ is
42. 0.01 mole of NaOH in 10 litre water are added. How much the pH water changes?
43. The approximate pH of a solution of benzoic acid ($K_a = 6.4 \times 10^{-5}$) having density 2.06 g/dm^3 is
44. A mixture of KOH and Ca(OH)_2 weighing 6.13 g is completely neutralized by an acid. If mass percent of KOH in mixture is 45.68 and normality of acid is 20 N, then find the approximate volume (in mL) of acid used in neutralisation.
45. 50 mL of 0.2 N NaCN is mixed with 50 mL of 0.2 M HCl . Calculate approximate concentration of $[\text{H}^+]$ in molarity $\times 10^{-6}$ if K_b for CN^- is 2×10^{-5} .
46. The dissociation constant of a substituted benzoic acid at 25°C is 1×10^{-4} . The pH of a 0.01 M solution of its sodium salt is : (IIT 2009)
47. The total number of diprotic acids among the following is..... (IIT 2010)
 H_3PO_4 , H_2SO_4 , H_3PO_3 , H_2CO_3 , $\text{H}_2\text{S}_2\text{O}_7$
 H_3BO_3 , H_3PO_2 , H_2CrO_4 , H_2SO_3
48. Amongst the following, the total number of compounds whose aqueous solution turns red litmus paper to blue is..... (IIT 2010)
 KCN , K_2SO_4 , $(\text{NH}_4)_2\text{C}_2\text{O}_4$, NaCl , $\text{Zn(NO}_3)_2$
 FeCl_3 , K_2CO_3 , NH_4NO_3 , LiCN
49. In one litre saturated solution of AgCl [K_{sp} of $\text{AgCl} = 1.6 \times 10^{-10}$], 0.1 mole of CuCl [K_{sp} of $\text{CuCl} = 1.0 \times 10^{-6}$] is added. The resultant concentration of Ag^+ in the solution is 1.6×10^{-X} . The value of X is..... (IIT 2011)

ANSWERS

1. Seven 2. Two 3. Nine 4. Five 5. Nine 6. Six 7. Five 8. Two 9. Six 10. Six 11. Two 12. Four
 13. Three 14. Nine 15. Eight 16. Seven 17. Two 18. Eight 19. Six 20. Seven 21. Four 22. Seven 23. One 24. Six
 25. Seven 26. Four 27. Nine 28. Seven 29. One 30. One 31. Six 32. Six 33. Two 34. Nine 35. One 36. Six
 37. Eight 38. Three 39. Four 40. One 41. One 42. Four 43. Three 44. Seven 45. Seven 46. Eight 47. Six 48. Three
 49. Seven

OBJECTIVE PROBLEMS (One Answer Correct)

- The equilibrium constant for :
 $\text{CN}^- + \text{CH}_3\text{COOH} \rightleftharpoons \text{HCN} + \text{CH}_3\text{COO}^-$ is :
 (Given $\text{p}K_b$ for $\text{CN}^- = 4.69$ and pH for $\text{CH}_3\text{COO}^- = 9.25$)
 (a) 3.7×10^4 (b) 2.8×10^{-5}
 (c) 1.97 (d) 0.5
- The ionisation of codein in 0.01 M solution shows a $\text{pH} = 10.1$. The ionisation constant of codein is :
 (a) 1.6×10^{-4} (b) 1.6×10^{-6}
 (c) 1.6×10^{-8} (d) 1.6×10^{-3}
- The solubility of a sparingly soluble salt $A(\text{OH})_2$ (molar mass 192.3 g mol^{-1}) is 19.23 g/litre at 300 K. The pH of its saturated solution assuming 80% ionisation at 300 K is :
 (a) 1.0970 (b) 12.9030
 (c) 13.2041 (d) 12.0000
- K_b for $\text{CH}_2\text{ClCOO}^-$ is 7.41×10^{-12} . The pH of 0.1 M $\text{CH}_2\text{ClCOONa}$ in water is :
 (a) 7.93 (b) 6.66
 (c) 1.94 (d) 12.06
- Equimolar solution of CaCl_2 and K_2SO_4 are mixed in equal volumes. If K_{sp} of CaSO_4 is 9.1×10^{-6} , the maximum molar concentrations of CaCl_2 and K_2SO_4 which will lead no precipitation of CaSO_4 is :
 (a) 0.006 (b) 0.004
 (c) 0.002 (d) 0.008
- If K_{sp} of ZnS is 1.0×10^{-21} , the minimum volume of water is required for the dissolution of 0.097 g ZnS is :
 (a) 3.16×10^5 litre (b) 3.16×10^7 litre
 (c) 3.16×10^9 litre (d) 3.16×10^3 litre
- K_{sp} of $M(\text{OH})_x$ is 27×10^{-12} and its solubility in water is $10^{-3} \text{ mole litre}^{-1}$. The value of x is :
 (a) 1 (b) 2
 (c) 3 (d) 4
- The dissociation constants for acetic acid are 1.8×10^{-5} and 1.805×10^{-5} at 300 and 310 K respectively. The enthalpy of deprotonation for acetic acid is :
 (a) 50 cal (b) 52.8 cal
 (c) 51.6 cal (d) 48.48 cal
- Solubility product of $A_2X_3 \cdot 4\text{H}_2\text{O}$ is given by :
 (a) $[A^{3+}]^2[X^{2-}]^3[\text{H}_2\text{O}]^4$
 (b) $[2A^{3+}]^2[3X^{2-}]^3[\text{H}_2\text{O}]^4$
 (c) $[2A^{2+}]^2[3X^{3-}]^3[\text{H}_2\text{O}]^4$
 (d) $[A^{3+}]^2[X^{2-}]^3$
- pH of some solutions is given by : $[\text{H}^+] = \frac{\text{p}K_{a1} + \text{p}K_{a2}}{2}$.
 For which type of compounds this formula is not valid?
 (a) NaHS (b) NaH_2PO_4
 (c) NaH_2PO_3 (d) NaH_2BO_3
- Which one is not correct statement about buffer mixture ?
 (a) NH_4OH + its conjugate acid
 (b) CH_3COONa + its conjugate base
 (c) 1 millimole of NaCN + 0.5 millimole of HCl in 1 litre
 (d) 0.5 millimole of NaCN + 1 millimole of HCl in 1 litre
- A buffer solution is prepared by dissolving 1 mole of CH_3COOH and 'a' mole of Zn acetate. The pH of solution is 4.7447 and $\text{p}K_a$ for $\text{CH}_3\text{COOH} = 4.7447$. The number of mole of zinc acetate are :
 (a) 1 (b) 2
 (c) 0.5 (d) 0.25
- The percentage error in H^+ concentration provided by 10^{-8} M HCl if ionisation of water is not neglected is :
 (a) 3% (b) 2%
 (c) 5% (d) 4%
- The degree of dissociation of water is 1.8×10^{-9} at 300 K. The self ionisation constant for water is :
 (a) 1.0×10^{-14} (b) 1.8×10^{-16}
 (c) 3.23×10^{-18} (d) none of these
- The change in pH of water when 0.01 mole of NaOH are added in 10 litre water is :
 (a) 11 units (b) 3 units
 (c) 4 units (d) 2 units
- Which of the following has concentration dependent pH ?
 (a) CH_3COONa (b) $\text{CH}_2\text{NH}_2\text{COOH}$
 (c) $\text{CH}_3\text{COONH}_4$ (d) NaHS
- An acid-base indicator ($\text{p}K_a = 4.5271$) has the acid form red and basic form blue. If we need 75% red to be converted into 75% blue form in solution, the change in pH of solution should be :
 (a) 4.05 (b) 5.0
 (c) 0.95 (d) 0.80
- 0.1 mole of CH_3NH_2 ($\text{p}K_b = 3.3010$) is mixed with 0.08 mol of HCl and volume is made to 1 litre. The pH of solution is :
 (a) 7.32 (b) 11.4642
 (c) 10.097 (d) 11.5020
- 0.08 mole of CH_3NH_2 ($\text{p}K_b = 3.3010$) is mixed with 1.0 mole of HCl and volume is made to 1 litre. The pH of solution is :
 (a) 1.699 (b) 2.699
 (c) 10.097 (d) 7.32
- 0.04 M solution of CH_3NH_2 ($\text{p}K_b = 3.3010$) is diluted to reduce its molar concentration to half. The change in pH during dilution is :

- (a) +0.1615 (b) -0.1615
(c) +0.2615 (d) -0.2615
21. The enthalpy change for first proton neutralisation of H_2S is $-33.7 \text{ kJ mol}^{-1}$. What is enthalpy change for first ionisation of H_2S ?
(a) 23.6 (b) 33.7
(c) 20.8 (d) 10.4
22. Which will have pH closer to 1.0?
(a) 100 mL of 0.1 M HCl + 100 mL 0.1 M NaOH
(b) 55 mL of 0.1 M HCl + 45 mL of 0.1 M NaOH
(c) 10 mL of 0.1 M HCl + 90 mL 0.1 M NaOH
(d) 75 mL of 0.2 M HCl + 25 mL 0.2 M NaOH
23. The pH of 0.1 M solution of the following salts increases in the order:
(a) $\text{NaCl} < \text{NH}_4\text{Cl} < \text{NaCN} < \text{HCl}$
(b) $\text{HCl} < \text{NH}_4\text{Cl} < \text{NaCl} < \text{NaCN}$
(c) $\text{NaCN} < \text{NH}_4\text{Cl} < \text{NaCl} < \text{HCl}$
(d) $\text{HCl} < \text{NaCl} < \text{NaCN} < \text{NH}_4\text{Cl}$
24. The conjugate acid of NH_2^- is:
(a) NH_3 (b) NH_2OH
(c) NH_4^+ (d) N_2H_4
25. The compound that is not Lewis acid is:
(a) BF_3 (b) AlCl_3
(c) SnCl_4 (d) CCl_4
26. The pK_a of acetyl salicylic acid (aspirin) is 3.5. The pH of gastric juice in human stomach is about 2–3 and the pH in the small intestine is about 8. Aspirin will be:
(a) unionised in small intestine and in the stomach
(b) completely ionised in the small intestine and in the stomach
(c) ionised in the stomach and almost unionised in the small intestine
(d) ionised in the small intestine and almost unionised in the stomach
27. Which of the following is the strongest acid?
(a) $\text{ClO}_3(\text{OH})$ (b) $\text{ClO}_2(\text{OH})$
(c) $\text{SO}(\text{OH})_2$ (d) $\text{SO}_2(\text{OH})_2$
28. pK_b of NH_4OH is 5. The pH of a mixture containing 10 mL of 0.3 M NH_4OH and 200 mL of 0.1 M $(\text{NH}_4)_3\text{PO}_4$ is:
(a) 5 (b) 5.3010
(c) 6.3010 (d) 7.6987
29. K_w for water is $9.5 \times 10^{-14} \text{ m}^2$ at 60°C . An aqueous solution of a salt at 60°C has pH = 7, it is:
(a) acidic (b) basic
(c) neutral (d) none of these
30. 100 mL of 0.005 M H_2SO_4 is diluted to one litre. The pH of resulting solution is:
(a) 3 (b) 4
(c) 5 (d) 2
31. How much water should be added to 300 mL of 0.2 M solution of acetic acid ($K_a = 1.8 \times 10^{-5}$) to increase its degree of dissociation two times?
(a) 1.2 litre (b) 1.5 litre
(c) 0.9 litre (d) 0.7 litre
32. If pK_b for $\text{C}_6\text{H}_5\text{O}^-$ is 4, what is the pH of 0.1 M $\text{C}_6\text{H}_5\text{O}^-$ in solution at 25°C ?
(a) 13.0 (b) 12.5
(c) 11.5 (d) 10.5
33. 2, 4-dinitrophenol ($\text{pK}_a = 4$) acts as buffer and if pH of this buffer is adjusted to 5, the ratio of concentration of dissociated and undissociated molecules of 2, 4-di-nitrophenol is:
(a) 1 (b) 10
(c) 0.1 (d) 0.01
34. In a saturated solution of AgCl , addition of NaCl is made drop by drop in excess. Which graph correctly represents the change?
- (a) $[\text{Ag}^+]$ vs $[\text{Cl}^-]$

(b) $[\text{Ag}^+]$ vs $[\text{Cl}^-]$

(c) $[\text{Ag}^+]$ vs $[\text{Cl}^-]$

(d) $[\text{Ag}^+]$ vs $[\text{Cl}^-]$
35. Solubility of Ag_2CrO_4 in 0.2 M K_2CrO_4 is 'a' and solubility of Ag_2CrO_4 in 0.4 M AgNO_3 is 'b', then:
(a) $a = b$
(b) $a > b$
(c) $a < b$
(d) cannot be predicted in absence of K_{sp} of Ag_2CrO_4
36. 0.1 M solution of H_3A being a weak tribasic acid having K_{a1} , K_{a2} and K_{a3} as 10^{-5} , 10^{-9} and 10^{-13} respectively. If pX represents $-\log \frac{[\text{A}^{3-}]}{[\text{HA}^{2-}]}$, then the value of pX is:
(a) 7 (b) 8
(c) 10 (d) 9
37. pH of mixture having 0.1 M NH_4OH and 0.1 M NH_4Cl is equal to $\text{pK}_w - \text{pK}_b$. If 0.1 M NH_4OH and 0.1 M $(\text{NH}_4)_2\text{SO}_4$ are taken, the pH will be:
(a) $\text{pK}_w - \text{pK}_b$ (b) $\text{pK}_w + \text{pK}_b$
(c) $\text{pK}_w - \text{pK}_b + \log 2$ (d) $\text{pK}_w - \text{pK}_b - \log 2$
38. In the equilibrium: $\text{BaF}_2(\text{s}) \rightleftharpoons \text{Ba}^{2+}(\text{aq}) + 2\text{F}^{-}(\text{aq})$ if $[\text{Ba}^{2+}]$ is increased two times, then $[\text{F}^-]$ in solution will:
(a) increase two times (b) increase four times
(c) decrease two times (d) decrease to $1/\sqrt{2}$ times

39. The pH of 100 mL solution containing 0.3 M NH_4^+ , 0.2 M NH_4OH ($K_b = 4.74$) and 0.01 M HCl in it is :
 (a) 9.05 (b) 8.66
 (c) 7.46 (d) 8.05
40. The pH of a 100 mL solution containing 0.3 mole of NH_4^+ , 0.2 M NH_4OH and 0.01 M NaOH in it is :
 (a) 10.12 (b) 9.12
 (c) 8.12 (d) 8.5
41. A weak acid has pH = 4, then :
 (a) $C = 10^{-3}$, $\alpha = 10\%$ (b) $C = 10^{-3}$, $\alpha = 10^{-6}$
 (c) $[A^-] = 10^{-5}$ (d) $K_a = 10^{-2}$, $\alpha = 10$
42. An acidic buffer solution can be prepared by mixing the solution of:
 (a) ammonium acetate and acetic acid
 (b) ammonium chloride and ammonium hydroxide
 (c) sulphuric acid and sodium sulphate
 (d) sodium chloride and sodium hydroxide
43. The pH of a 10^{-8} molar solution of HCl in water is :
 (a) 8 (b) -8
 (c) between 7 and 8 (d) between 6 and 7
44. Of the given anions, the strongest Bronsted base is :
 (a) ClO^- (b) ClO_2^-
 (c) ClO_3^- (d) ClO_4^-
45. At 90°C, pure water has $[\text{H}_3\text{O}^+] = 10^{-6}$ mole litre $^{-1}$. What is the value of K_w at 90°C?
 (a) 10^{-6} (b) 10^{-12}
 (c) 10^{-14} (d) 10^{-8}
46. The precipitate of CaF_2 ($K_{sp} = 1.7 \times 10^{-10}$) is obtained when equal volumes of the following are mixed :
 (a) 10^{-4} M Ca^{2+} + 10^{-4} M F^-
 (b) 10^{-2} M Ca^{2+} + 10^{-3} M F^-
 (c) 10^{-5} M Ca^{2+} + 10^{-3} M F^-
 (d) 10^{-3} M Ca^{2+} + 10^{-5} M F^-
47. A certain buffer solution contains equal concentration X^- and HX . Then K_b for X^- is 10^{-10} . The pH of the buffer is :
 (a) 4 (b) 7
 (c) 10 (d) 14
48. A certain weak acid has a dissociation constant of 1.0×10^{-4} . The equilibrium constant for its reaction with a strong base is :
 (a) 1.0×10^{-4} (b) 1.0×10^{-10}
 (c) 1.0×10^{10} (d) 1.0×10^{14}
49. The best indicator for detection of end point in titration of a weak acid and a strong base is :
 (a) methyl orange (3 to 4)
 (b) methyl red (5 to 6)
 (c) bromothymol blue (6 to 7.5)
 (d) phenolphthalein (8 to 9.6)
50. The following equilibrium is established when hydrogen chloride is dissolved in acetic acid.

$$\text{HCl} + \text{CH}_3\text{COOH} \rightleftharpoons \text{Cl}^- + \text{CH}_3\text{COOH}_2^+$$
 The set that characterises the conjugate acid-base pairs is :
 (a) (HCl , CH_3COOH) and ($\text{CH}_3\text{COOH}_2^+$, Cl^-)
 (b) (HCl , $\text{CH}_3\text{COOH}_2^+$) and (CH_3COOH , Cl^-)
 (c) ($\text{CH}_3\text{COOH}_2^+$, HCl) and (Cl^- , CH_3COOH)
 (d) (HCl , Cl^-) and ($\text{CH}_3\text{COOH}_2^+$, CH_3COOH)
51. The compound insoluble in acetic acid is :
 (a) calcium oxide (b) calcium carbonate
 (c) calcium oxalate (d) calcium hydroxide
52. The compound whose 0.1 M solution is basic is :
 (a) ammonium acetate (b) ammonium chloride
 (c) ammonium sulphate (d) sodium acetate
53. When equal volumes of the following solutions are mixed, precipitation of AgCl ($K_{sp} = 1.8 \times 10^{-10}$) will occur only with :
 (a) 10^{-4} M (Ag^+) and 10^{-4} M (Cl^-)
 (b) 10^{-5} M (Ag^+) and 10^{-5} M (Cl^-)
 (c) 10^{-6} M (Ag^+) and 10^{-6} M (Cl^-)
 (d) 10^{-10} M (Ag^+) and 10^{-10} M (Cl^-)
54. The degree of dissociation of water at 25°C is $1.9 \times 10^{-7}\%$ and density is 1.0 g cm^{-3} . The equilibrium constant for water is :
 (a) 1.0×10^{-14} (b) 2.0×10^{-14}
 (c) 2.0×10^{-16} (d) 1.0×10^{-8}
55. Which one is more acidic in aqueous solution ?
 (a) NiCl_2 (b) FeCl_3
 (c) AlCl_3 (d) BeCl_2
56. The following acids have been arranged in the order of decreasing acid strength. Identify the correct order
 ClOH (I), BrOH (II), IOH (III)
 (a) I > II > III (b) II > I > III
 (c) III > II > I (d) I > III > II
57. If $\text{p}K_b$ for fluoride ion at 25°C is 10.83, the ionisation constant of hydrofluoric acid in water at this temperature is :
 (a) 1.74×10^{-5} (b) 3.52×10^{-3}
 (c) 6.75×10^{-4} (d) 5.38×10^{-2}
58. The solubility of A_2X_3 is $y \text{ mol dm}^{-3}$. Its solubility product is :
 (a) $6y^4$ (b) $64y^4$
 (c) $36y^5$ (d) $108y^5$

59. Among the following the one having lowest K_{sp} is :
 (a) $Mg(OH)_2$ (b) $Ca(OH)_2$
 (c) $Ba(OH)_2$ (d) $Be(OH)_2$
60. Which of the following is correct for 1.0 M solution of strong acid HA and 1.0 M solution of weak acid HB ?
 (a) HA : pH = 0; HB : pH < 1
 (b) HA : $[H^+] = [A^-]$; HB : $[H^+] = [A^-]$
 (c) HA : $[HA] = 0$; HB : $[HB] = (1 - \alpha)$
 (d) All of these
61. Which order is not correct for basic nature ?
 (a) $NH_3 > PH_3$ (b) $H_2O > H_2S$
 (c) $NH_3 > NCl_3$ (d) $NF_3 > NH_3$
62. Which of the following acid-base neutralisation does not form water with salt ?
 (a) $Al_2O_3(s) + NaOH(aq.)$ (b) $Al_2O_3 + HCl(aq.)$
 (c) $SiO_2 + NaOH(aq.)$ (d) $ZnO + NaOH(aq.)$
63. How many mole of HCl must be added to 1 litre aqueous solution of HCl of pH = 3, in order to decrease its pH to 2?
 (a) 0.009 (b) 0.001
 (c) 1 (d) 0.1
64. Which one is correct decreasing order of solubility of AgCl ?
 (i) in water (ii) in 0.1 M NaCl
 (iii) in 0.1 M $BaCl_2$ (iv) in 0.1 M NH_3
 (a) (iv) > iii > ii > i (b) (iv) > i > ii > iii
 (c) (iv) > ii > iii > i (d) (i) > ii > iii > iv
65. 0.1 M aqueous solution of BOH (a weak base) has pH equal to 12 at 27°C. Assuming K_w at 27°C 1.0×10^{-14} , calculate the osmotic pressure of 0.1 M BOH(aq.)
 (a) 2.71 atm (b) 2.46 atm
 (c) 0.246 atm (d) 24.6 atm
66. 300 mL of saturated, clear solution of $CaC_2O_4(aq.)$ is completely oxidised by 6 mL of 0.001 M $KMnO_4$ in acid medium. The K_{sp} of CaC_2O_4 is :
 (a) 25×10^{-8} (b) 25×10^{-9}
 (c) 25×10^{-10} (d) 25×10^{-11}
67. Solubility of AgCl (in mole/litre) in a mixture of 1 litre solution containing 0.05M NaCl and 0.05 M $BaCl_2$ is : (Given K_{sp} of AgCl = 1.0×10^{-10})
 (a) 6.6×10^{-8} (b) 6.6×10^{-10}
 (c) 1.0×10^{-9} (d) 1.0×10^{-8}
68. Select the correct choice when 100 mL of 0.5 M hydrazoic acid (N_3H having $K_a = 3.6 \times 10^{-4}$) and 400 mL of 0.1 M cyanic acid (HOCN having $K_a = 8 \times 10^{-4}$) are mixed :
 (a) pH = 2 (b) $[N_3^-] = 3.6 \times 10^{-3}$
 (c) $[OCN^-] = 6.4 \times 10^{-3}$ (d) all of these
69. Autoprotonation constant for liquid NH_3 is 3.46×10^{-27} at $-50^\circ C$. The number of anions present in 1 mL of pure NH_3 at $-50^\circ C$ is :
 (a) 2.08×10^9 (b) 1×10^{-15}
 (c) 6.023×10^{-5} (d) 1×10^{15}
70. Solubility product of AgCl is 1.0×10^{-10} . What minimum volume of water in litre is required to dissolve 4.305 mg of AgCl?
 (a) 1 (b) 2
 (c) 3 (d) 4
71. How many mixed buffers can be obtained using KOH and H_3PO_4 ?
 (a) 1 (b) 2
 (c) 3 (d) 4
72. Which graph correctly represents addition of NaCl to a saturated solution of AgCl?
- (a)

(b)
- (c)

(d)
73. Which reagent can be used to separate one of ions as precipitate from a solution containing Cu^{2+} and Pb^{2+} ions ?
 (a) $H_2S(g)$ (b) NH_4NO_3
 (c) HNO_3 (d) H_2SO_4
74. Which can be used to separate F^- and Cl^- in solution ?
 (a) $AgNO_3$ (b) $Pb(NO_3)_3$
 (c) HNO_3 (d) H_2SO_4
75. Except one in rest all cases, K_2CO_3 solution on addition gives a precipitate :
 (a) $BaCl_2(aq.)$ (b) $CaBr_2(aq.)$
 (c) $(NH_4)_2SO_4$ (d) $Pb(NO_3)_2$
76. Arrange the following in increasing order of pH :
 1. 0.1 M CH_3COONa + 0.1 M CH_3COOH
 2. 0.1 M CH_3COOH + 0.1 M HCl
 3. 0.1 M CH_3COOH
 4. 0.1 M HCl
 (a) $2 < 4 < 3 < 1$ (b) $1 < 2 < 3 < 4$
 (c) $2 < 3 < 4 < 1$ (d) $4 < 3 < 2 < 1$
77. A solution has pH = 4. It is 1000 times less acidic than another solution of same species of pH :

- (a) 1 (b) 2
(c) 3 (d) 4
78. pOH of 0.003 M HCl is :
(a) $11 + \log 3$ (b) $11 - \log 3$
(c) $7 + \log 3$ (d) $7 - \log 3$
79. 10 mL of $10^{-4} \text{ N H}_2\text{SO}_4$ is diluted with water to have 1 dm^3 solution : The pH of this solution is :
(a) 6 (b) slightly more than 6
(c) slightly less than 6 (d) 7
80. The pH of a solution obtained by mixing 10 mL of 0.1 M HCl and 10 mL of 0.01 M Sr(OH)_2 .
(a) 1.40 (b) 1.50
(c) 11.70 (d) 7.00
81. If heat of ionisation $\Delta_i H^\circ$ for HCN and $\Delta_i H^\circ$ for CH_3COOH are 45.2 kJ mol^{-1} and 2.1 kJ mol^{-1} , then which one is not correct ?
(a) $pK_{a\text{HCN}} > pK_{a\text{CH}_3\text{COOH}}$
(b) K_a of $\text{HCN} < K_a$ of CH_3COOH
(c) acetic acid is stronger than HCN
(d) $pK_{a\text{HCN}} < pK_{a\text{CH}_3\text{COOH}}$
82. The ratio of degree of dissociation of HCN ($K_a = 10^{-9}$) in its 0.1 M and 0.001 M solution :
(a) 0.1 (b) 0.2
(c) 0.3 (d) 0.4
83. Which of the following ions can exist in solution ?
(a) H_3O^+ (b) H_3O_2^-
(c) H_5O_2^+ (d) all of these
84. What will be degree of hydrolysis and pH respectively of an aqueous solution of ammonium formate : $pK_a = 4$, $pK_b = 5$; $pK_w = 14$:
(a) 3.16×10^{-3} , 7 (b) 4.16×10^{-3} , 6.5
(c) 3.16×10^{-3} , 6.5 (d) 4.16×10^{-3} , 7.0
85. The solubility product of AgCl is 10^{-10} . Applying Debye Huckel limiting law, the correct solubility order of AgCl in different solutions of 0.1 M concentration is :
(a) $\text{NaCl} < \text{H}_2\text{O} < \text{NaNO}_3 < \text{Ca(NO}_3)_2$
(b) $\text{H}_2\text{O} < \text{NaCl} < \text{NaNO}_3 < \text{Ca(NO}_3)_2$
(c) $\text{NaNO}_3 < \text{Ca(NO}_3)_2 < \text{H}_2\text{O} < \text{NaCl}$
(d) $\text{Ca(NO}_3)_2 < \text{NaNO}_3 < \text{H}_2\text{O} < \text{NaCl}$
86. For, $\text{Ag(CN)}_2^- \rightleftharpoons \text{Ag}^+ + 2\text{CN}^-$
 $K_C = 4 \times 10^{-19}$ at 25°C .
Calculate Ag^+ concentration in a solution which is originally 0.1 M in KCN and 0.03 M in AgNO_3 .
(a) $7.5 \times 10^{-18} \text{ M}$ (b) $7.5 \times 10^{-15} \text{ M}$
(c) $6.2 \times 10^{-15} \text{ M}$ (d) $3.4 \times 10^{-15} \text{ M}$
87. The solubility product of PbI_2 is 7.47×10^{-9} and 1.39×10^{-8} at 15°C and 25°C respectively. The molar heat of solution of PbI_2 is :
(a) 9.969 k cal (b) 10.66 k cal
(c) 8.88 k cal (d) 11.11 k cal
88. A solution of a weak monobasic acid has $\wedge_M$ for 0.1 N solution and $\wedge_\infty$ equal to $60 \text{ S cm}^2 \text{ mol}^{-1}$ and $400 \text{ S cm}^2 \text{ mol}^{-1}$ respectively. The pH of this solution and K_a of acid is :
(a) 1.82, 2.65×10^{-3} (b) 2.82, 2.65×10^{-3}
(c) 1.82, 2.25×10^{-3} (d) 2.82, 2.25×10^{-3}
89. The dissociation constant of HA and A^- are equal. The pH of 0.001 HA solution is :
(a) 3 (b) 4
(c) 5 (d) 6
90. pK_a of acetic acid is 4.74. What mass of KOH is required to be added in 500 mL of 1 M acid in order to prepare a buffer solution of maximum capacity.
(a) 28 g (b) 14 g
(c) 7 g (d) 10 g
91. K_a of HCOOH is 10^{-5} and solubility of HCOOAg is 10^{-2} M . The solubility of HCOOAg in a buffer solution of $\text{pH} = 3$ is
(a) 0.01 M (b) 0.1 M
(c) 0.2 M (d) 0.02 M
92. From the following informations, select the strongest base :
Acid H_2PO_4^- HPO_4^{2-} HCO_3^- HSO_3^-
 K_a 6.0×10^{-8} 4.8×10^{-13} 4.7×10^{-11} 6.3×10^{-8}
(a) PO_4^{3-} (b) HPO_4^{2-}
(c) CO_3^{2-} (d) SO_3^{2-}
93. 0.125 mole of Ca A_2 are present in 0.5 litre solution. If K_a of HA is 8.0×10^{-4} and salt CaA_2 is completely ionised, the pH of solution is :
(a) 5.60 (b) 5.75
(c) 8.40 (d) 8.25
94. At what $\frac{[\text{Br}^-]}{\sqrt{[\text{CO}_3^{2-}]}}$, the following cell shows equilibrium :
 $\text{Ag(s)} | \text{Ag}_2\text{CO}_3(\text{s}), \text{Na}_2\text{CO}_3(\text{aq.}) || \text{AgBr(aq.)} | \text{Ag(s)}$
 K_{SP} of $\text{Ag}_2\text{CO}_3 = 8 \times 10^{-12}$ and
 K_{SP} of $\text{AgBr} = 4 \times 10^{-13}$
(a) 2×10^{-7} (b) $\sqrt{2} \times 10^{-7}$
(c) $\sqrt{3} \times 10^{-7}$ (d) 2
95. Concentration of the Ag^+ ions in a saturated solution of $\text{Ag}_2\text{C}_2\text{O}_4$ is $2.2 \times 10^{-4} \text{ mol L}^{-1}$. Solubility product of $\text{Ag}_2\text{C}_2\text{O}_4$ is :
(a) 2.42×10^{-8} (b) 2.66×10^{-12}
(c) 4.5×10^{-11} (d) 5.3×10^{-12}

SOLUTIONS (One Answer Correct)

1. (a) $K_{eq} = \frac{[HCN][CH_3COO^-]}{[CN^-][CH_3COOH]}$
 $K_{CH_3COOH} = \frac{[CH_3COO^-][H^+]}{[CH_3COOH]}$
 $K_{HCN} = \frac{[H^+][CN^-]}{[HCN]}$
 $pK_{HCN} = 4.69 \therefore pK_{HCN} = 9.31$
 $pK_{CH_3COOH} = 4.75$
 $\therefore K_{eq} = \frac{K_{CH_3COOH}}{K_{HCN}} = \frac{1.77 \times 10^{-5}}{4.9 \times 10^{-10}} = 3.7 \times 10^4$
2. (b) pH of Codein = 10.1 $\therefore$ it is a base.
 Now $Codein + H_2O \rightleftharpoons Codein^+ + OH^-$
 $K_b = \frac{[Codein^+][OH^-]}{[Codein]}$
 Also, $[OH^-] = c\alpha = 0.01 \times \alpha$
 $-\log 3.9 = 0.01\alpha$
 $\therefore \alpha = \frac{1.259 \times 10^{-4}}{0.01} = 0.0125$
 $K_a = \frac{c\alpha^2}{(1-\alpha)}$
 $K_a = \frac{0.01 \times (0.0125)^2}{(1-0.0125)} = 1.6 \times 10^{-6}$
3. (c) $S = \frac{19.23}{192.3} \text{ mol/litre} = 0.1 M$
 $A(OH)_2 \rightleftharpoons A + 2OH^-$
 $\therefore [OH^-] = \frac{0.1 \times 2 \times 80}{100} = 0.16$
 $\therefore pOH = 0.7958$
 $\therefore pH = 13.2041$
4. (a) $CH_2ClCOO^- + H_2O \rightleftharpoons CH_2ClCOOH + OH^-$
 $[OH^-] = ch = \sqrt{\frac{K_H}{c}} = \sqrt{K_H \cdot c} = \sqrt{\frac{K_w \times c}{K_a}}$
 $= \sqrt{K_b \times c} = \sqrt{7.41 \times 10^{-12} \times 0.1}$
 $= 8.606 \times 10^{-7}$
 $\therefore pOH = 6.065$
 $pH = 7.93$
5. (a) Let molar concentration of salts be $a M$
 $[Ca^{2+}][SO_4^{2-}] = K_{sp}$
 $\left[\frac{a \times V}{2V}\right]\left[\frac{a \times V}{2V}\right] = 9.1 \times 10^{-6}$
 $\therefore a^2 = 4 \times 9.1 \times 10^{-6} \therefore a = 6.033 \times 10^{-3}$
6. (b) Solubility of $ZnS = S = \sqrt{K_{sp}} = \sqrt{1 \times 10^{-21}}$
 $= 3.16 \times 10^{-11} \text{ mol/litre}$
 $= 3.16 \times 10^{-11} \times 97 \text{ g/L}$
 $= 3.065 \times 10^{-9} \text{ g/L}$
 $\therefore \text{Volume of water in litre required} = \frac{0.097}{3.065 \times 10^{-9}}$
 $= 3.16 \times 10^7 \text{ litre}$
7. (c) $K_{sp} \text{ of } M(OH)_X = X^X \cdot (S)^{X+1} = 27 \times 10^{-12}$
 $\therefore X^X \cdot (10^{-3})^{X+1} = 27 \times 10^{-12}$
 By putting $X = 1, 2, 3, \dots$
 $X = 3$
8. (c) $2.303 \log \frac{K_2}{K_1} = \frac{\Delta H}{R} \left[\frac{T_2 - T_1}{T_1 T_2} \right]$
 $2.303 \log \frac{1.805 \times 10^{-5}}{1.8 \times 10^{-5}} = \frac{\Delta H}{R} \times \frac{10}{300 \times 310}$
 $\therefore \Delta H = 51.6 \text{ cal}$
9. (d) Hydrated water molecules do not interfere in K_{sp} .
10. (d) The formula is valid for acidic salts as well as for zwitter ionic molecules. NaH_2BO_3 is neither an acid salt nor it exists.
11. (d) $NaCN + HCl \rightarrow NaCl + HCN$

1	0.5	0	0
0.5	0	0.5	0.5

i.e., 0.5 NaCN + 0.5 HCN, buffer
 $NaCN + HCl \rightarrow NaCl + HCN$


0.5	1	0	0
0	0.5	0.5	0.5

 Not a buffer
12. (c) $pH = pK_a + \log \frac{2a}{1}$
 $(CH_3COO)_2Zn$ gives 2 mol CH_3COO^-
 $4.7447 = 4.7447 + \log \frac{2a}{1} \therefore a = 1/2$
13. (c) $10^{-8} M$ $HCl \therefore [H^+] = 10^{-8} = 10 \times 10^{-9}$ if dissociation of water is neglected
 if not neglected then: $H_2O \rightleftharpoons H^+ + OH^-$
 $a + 10^{-8} \quad a$
 $\therefore a \times (a + 10^{-8}) = 10^{-14}$
 $a = 0.95 \times 10^{-7}$
 $\therefore \% \text{ error} = \frac{(10 \times 10^{-9} - 9.5 \times 10^{-9})}{10 \times 10^{-9}} \times 100 = 5\%$
14. (c) $H_2O + H_2O \rightleftharpoons H_3O^+ + OH^-$
 $[H_3O^+] = C\alpha = 10^{-7} = [OH^-]$
 $\therefore C = \frac{10^{-7}}{1.8 \times 10^{-9}} = 55.6$
 $K_{s.t.} = \frac{[H_3O^+][OH^-]}{[H_2O]^2} = \frac{10^{-14}}{(55.6)^2} = 3.23 \times 10^{-18}$

15. (c) $[\text{OH}^-] = \frac{0.01}{10} = 10^{-3} \therefore \text{pOH} = 3 \text{ or } \text{pH} = 11$

Thus, pH changes from 7 to 11 by 4 units.

16. (a) pH of acidic salts, salts of weak-acid + weak-base and zwitter ionic species is independent of concentration.



$$K_a = \frac{[\text{H}^+][\text{I}_n^-]}{[\text{HI}_n]}$$

$$\text{pH} = \text{p}K_a + \log \frac{25}{75} = 4.5271 - 0.4771 = 4.05$$

$$\begin{aligned} \text{For 75% blue form } \text{pH} &= 4.5271 + \log \frac{75}{25} \\ &= 4.5271 + 0.4771 = 5.0 \\ \therefore \Delta \text{pH} &= 5.0 - 4.05 = 0.95 \end{aligned}$$



$$\begin{array}{cccc} 0.1 & 0.08 & 0 & 0 \\ 0.02 & 0 & 0.08 & 0.08 \end{array}$$


$$[\text{CH}_3\text{NH}_2] = \frac{0.02}{1}, [\text{CH}_3\text{NH}_3^+] = 0.08$$

This acts as basic buffer.

$$\text{pOH} = \text{p}K_b + \log \frac{0.08}{0.02} = 3.3010 + 0.6020$$


$$\text{pOH} = 3.903$$

$$\therefore \text{pH} = 10.097$$



$$\begin{array}{cccc} 0.08 & 1 & 0 & 0 \\ 0 & 0.02 & 0.08 & 0.08 \end{array}$$

$$\therefore [\text{H}^+] = 0.02 = 2 \times 10^{-2}; \therefore \text{pH} = 1.699$$




$$K_b = \frac{c\alpha^2}{(1-\alpha)} = 5 \times 10^{-4} = \frac{0.04 \times \alpha^2}{(1-\alpha)} \quad (1-\alpha \neq 1)$$

$$\therefore \alpha = 0.1056$$

$$[\text{OH}^-] = c\alpha = 0.04 \times 0.1056 = 4.224 \times 10^{-3}$$

$$\text{pOH} = 2.3743$$




$$K_b = \frac{c\alpha^2}{(1-\alpha)} = 5 \times 10^{-4} = \frac{0.02 \times \alpha^2}{(1-\alpha)} \quad (1-\alpha \neq 1)$$

$$\therefore \alpha = 0.1456$$


$$\therefore [\text{OH}^-] = c\alpha = 0.02 \times 0.1456 = 2.912 \times 10^{-3}$$

$$\text{pOH} = 2.5358$$

$\therefore -\Delta \text{pOH} = \Delta \text{pH} = +0.1615$, i.e., pOH increases on dilution or pH decreases.









$$\therefore \text{H}_2\text{S} \rightleftharpoons \text{H}^+ + \text{HS}^-; \Delta H = +57.3 - 33.7 = +23.6 \text{ kJ}$$



$$\begin{array}{cccc} 75 \times 0.2 & 25 \times 0.2 & 0 & 0 \\ = 15 & 5 & 0 & 0 \\ 10 & 0 & 5 & 5 \end{array}$$


$$[\text{H}^+] = \frac{10}{100} = 0.1$$

$$\therefore \text{pH} = 1$$



$$\therefore \text{pOH} = 5 + \log \frac{60}{3} = 5 + \log 20 = 6.3010$$

$$\therefore \text{pH} = 14 - 6.3010 = 7.6987$$




$$\therefore [\text{H}^+] = 3.1 \times 10^{-7}$$

$$\text{If } [\text{H}^+] > 3.1 \times 10^{-7} \text{ acidic}$$

$$< 3.1 \times 10^{-7} \text{ basic}$$


Thus, if $[\text{H}^+] = 10^{-7}$ i.e., $< 3.1 \times 10^{-7}$, solution is basic.



$$0.005 \times 2 \times 100 = 1000 \times N$$

$$\therefore [\text{H}^+] = 1 \times 10^{-3}$$

$$\therefore \text{pH} = 3$$



If α is doubled, then new $\alpha_2 = 1.898 \times 10^{-2}$

Since α remains negligible in both cases, therefore

$$c_1\alpha_1^2 = c_2\alpha_2^2$$


$$\therefore c_2 = 0.2 \times \left(\frac{1}{2}\right)^2 = 0.05$$

$$\text{On dilution } c_1V_1 = c_2V_2 \quad (\text{or } M_1V_1 = M_2V_2)$$

$$300 \times 0.2 = 0.05 \times V_2$$

$$\therefore V_2 = 1200 \text{ mL}$$

$$\therefore \text{Volume of water added} = 900 \text{ mL}$$




$$[\text{OH}^-] = c \cdot h = c \cdot \sqrt{\frac{K_H}{c}} = \sqrt{\frac{K_w}{K_a}} \cdot c$$

$$= \sqrt{K_b \cdot c} = \sqrt{10^{-4} \times 0.1} = 3.16 \times 10^{-3}$$

$$\therefore \text{pOH} = -2.5$$

$$\therefore \text{pH} = 11.5$$



$$5 = 4 + \log \frac{[\text{conjugate base}]}{[\text{acid}]}$$

$$\log \frac{[\text{conjugate base}]}{[\text{acid}]} = 1$$

$$\therefore \frac{[\text{conjugate base}]}{[\text{acid}]} = \frac{[\text{dissociated ions}]}{[\text{undissociated acid}]} = 10$$

$$34. (b) [\text{Ag}^+][\text{Cl}^-] = K_{sp} \quad \text{or} \quad [\text{Ag}^+] = \frac{K_{sp}}{[\text{Cl}^-]}$$

if $[\text{Cl}^-]$ increases, then to have K_{sp} constant and thus $[\text{Ag}^+]$ decreases. Also after sufficient addition of $[\text{Cl}^-]$, $[\text{Ag}^+]$ will become almost constant.

$$35. (d) K_{sp} \text{Ag}_2\text{CrO}_4 = [\text{Ag}^+]^2[\text{CrO}_4^{2-}]$$

$$= (2s)^2 \times s = 4s^3$$

Now in 0.2 M K_2CrO_4

$$K_{sp} = [\text{Ag}^+]^2 \times 0.2$$

$$\therefore [\text{Ag}^+] = \sqrt{\frac{K_{sp}}{0.2}}$$

$$\text{or} \quad a = \sqrt{\frac{K_{sp}}{0.2}}$$

$$= 2.23 \times \sqrt{K_{sp}}$$

In 0.4 M AgNO_3

$$K_{sp} = [0.4 + s]^2 \times s$$

$$\therefore s = \frac{K_{sp}}{0.16}$$

$$b = \frac{K_{sp}}{0.16}$$


$$= 6.25 \times K_{sp}$$

36. (c) $[\text{H}^+]$ are obtained in dissociation of H_3A in all the three steps, but the $[\text{H}^+]$ obtained in II & III steps are too low. Thus $[\text{H}^+]$ can be obtained from I step

$$[\text{H}^+] = \sqrt{K_{a1} \times c} \quad (1 - \alpha \approx 1)$$

$$= \sqrt{10^{-5} \times 0.1} = 10^{-3}$$

$$\text{Now, } K_3 = \frac{[\text{H}^+][\text{A}^{3-}]}{[\text{HA}^{2-}]}$$



$$10^{-13} = \frac{10^{-3} \times [\text{A}^{3-}]}{[\text{HA}^{2-}]}$$

$$\therefore \frac{[\text{A}^{3-}]}{[\text{HA}^{2-}]} = 10^{-10} \quad \therefore pX = 10$$

37. (d) In I Case :

$$pOH = pK_b + \log \frac{[\text{NH}_4^+]}{[\text{NH}_4\text{OH}]} = pK_b$$

$$\therefore pH = pK_w - pK_b \quad [\because [\text{NH}_4^+] = [\text{NH}_4\text{OH}] = 0.1]$$

In II Case :

$$pOH = pK_b + \log \frac{[\text{NH}_4^+]}{[\text{NH}_4\text{OH}]} = pK_b + \log 2$$

$$\therefore pH = pK_w - pK_b - \log 2 \quad [\because [\text{NH}_4^+] = 0.1 \times 2]$$

$$38. (d) \text{BaF}_2 \rightleftharpoons \text{Ba}^{2+} + 2\text{F}^-$$

$$K_{sp} = [\text{Ba}^{2+}][\text{F}^-]^2$$

$$\therefore [\text{F}^-] = \sqrt{\frac{K_{sp}}{[\text{Ba}^{2+}]}}$$

$$\text{Again } K_{sp} = 2 \times [\text{Ba}^{2+}][\text{F}^-]^2$$

$[\text{Ba}^{2+}]$ is increased two times

$$[\text{F}^-]^2 = \frac{K_{sp}}{2 \times [\text{Ba}^{2+}]}$$

$$\text{or } [\text{F}^-] = \sqrt{\frac{K_{sp}}{2[\text{Ba}^{2+}]}}$$

$$39. (a) \text{NH}_4\text{OH} + \text{HCl} \rightleftharpoons \text{NH}_4\text{Cl} + \text{H}_2\text{O}$$

$$\begin{array}{ccc} 0.2 & 0.01 & 0 \\ 0.19 & 0 & 0.01 \end{array}$$

$$\therefore [\text{NH}_4^+] = 0.30 + 0.01 = 0.31$$

$$[\text{NH}_4\text{OH}] = 0.19$$

$$\therefore pOH = 4.74 + \log \frac{0.31}{0.19}$$

$$= 4.95$$

$$\therefore pH = 14 - 4.95 = 9.05$$

$$40. (b) \text{NH}_4^+ + \text{NaOH} \longrightarrow \text{NH}_4\text{OH} + \text{Na}^+$$

$$\begin{array}{ccc} 0.3 & 0.01 & 0 \\ 0.29 & 0 & 0.01 \end{array}$$

$$\therefore [\text{NH}_4^+] = 0.29$$

$$[\text{NH}_4\text{OH}] = 0.2 + 0.01 = 0.21$$

$$\therefore pOH = 4.74 + \log \frac{0.29}{0.21}$$

$$= 4.880$$

$$\therefore pH = 14 - 4.880 = 9.12$$

$$41. (a) \text{HA} \rightleftharpoons \text{H}^+ + \text{A}^-$$

$$[\text{H}^+] = [\text{A}^-] = \alpha \cdot c = \frac{10}{100} \times 10^{-3} = 10^{-4}$$

$$\therefore pH = 4$$

42. (a) Weak acid (CH_3COOH) and its conjugate base CH_3COO^- and this buffer is an acidic buffer.

43. (d) HCl is an acid.

44. (a) Stronger is the acid, weaker is its conjugate base.

45. (b) $K_w = [\text{H}_3\text{O}^+][\text{OH}^-]$. Also $[\text{H}_3\text{O}^+] = [\text{OH}^-]$

$$10^{-6} \times 10^{-6} = 10^{-12}$$

$$46. (b) Q \text{ for } \text{CaF}_2 = [\text{Ca}^{2+}][\text{F}^-]^2$$

$$= \left[\frac{10^{-2} \times V}{2V} \right] \left[\frac{10^{-3} \times V}{2V} \right]^2 = \frac{1}{8} \times 10^{-8} = 1.2 \times 10^{-9}$$

$$Q > K_{sp}$$

$$47. (a) pH = -\log K_a + \log \frac{[\text{conjugate base}]}{[\text{acid}]} = 10^{-4} + \log \frac{1}{1} = 4$$

$$[\because pK_a + pK_a = 14]$$

$$48. (c) \text{HX} \rightleftharpoons \text{H}^+ + \text{X}^-$$

$$K_a = 10^{-4} = \frac{[\text{H}^+][\text{X}^-]}{[\text{HX}]}$$

...(i)



$$K_{eq} = \frac{[Na^+][X^-][H_2O]}{[HX][Na^+][OH^-]} = \frac{[X^-][H_2O]}{[HX][OH^-]} \dots(ii)$$


By Eqns. (i) and (ii)

$$\frac{K_{eq}}{K_a} = \frac{1}{K_w} \therefore K_{eq} = \frac{K_a}{K_w} = \frac{10^{-4}}{10^{-14}} = 10^{10}$$

49. (d) Phenolphthalein is good indicator for strong alkali titrations.




51. (c) Calcium oxalate is weaker base.



53. (a) Q for $AgCl = [Ag^+][Cl^-] = \left[\frac{10^{-4} \times V}{2V} \right] \left[\frac{10^{-4} \times V}{2V} \right]$
 $= 2.5 \times 10^{-9}$

$\therefore Q > K_{sp}$, precipitation will occur.

54. (c) $K_{eq} = \frac{[H^+][OH^-]}{[H_2O]}$




$\therefore [H^+] = c\alpha$ and $c_{H_2O} = \frac{1000}{18} = 55.6$

$K_{eq} = \frac{c\alpha \cdot c\alpha}{c(1-\alpha)} \quad (1-\alpha=1)$

$\therefore K_{eq} = \frac{(55.6 \times 19 \times 10^{-9})^2}{55.6} = 2 \times 10^{-16}$

55. (b) Both $FeCl_3$ and $AlCl_3$ will furnish $3OH^-$ ions and $Fe(OH)_3$ is relatively stronger base than others.

56. (a) $HOCl > HOBr > HOI$ due to more electronegative nature of halogen.



$pK_a + pK_b = 14$



$pK_a + 10.83 = 14$

$\therefore pK_a = 3.17$


$K_a = 6.76 \times 10^{-4}$

58. (d) K_{sp} of $A_2X_3 = 2^2 \cdot (S)^2 \times 3^3 \times S^3$
 $= 108(S)^5 = 108y^5$

59. (d) The solubility of alkaline earth metal hydroxides increases down the gp. more is solubility, more is K_{sp} .



61. (d) F is more electronegative than H and thus it reduces electron pair donation nature in NF_3 .




63. (a) $[H^+] = [HCl]$ or mole of HCl of $pH = 3$ in 1 litre is $10^{-3} = 0.001$

$[H^+] = [HCl]$ or mole of HCl of $pH = 2$ in 1 litre is $10^{-2} = 0.01$

$\therefore$ Mole of HCl to be added in HCl of $pH = 3$ are $0.01 - 0.001 = 0.009$

64. (b) $AgCl$ forms complex with NH_3 and thus more soluble. Also solubility of a salt decreases in presence of common ion.




$[OH^-] = C\alpha = 10^{-2}$

[For BOH , $pH = 12 \therefore [OH^-] = 10^{-2}$]

$\therefore \alpha = \frac{10^{-2}}{0.1} = 10^{-1} = 0.1$

$\therefore$ Total number of particles present are $(1-\alpha+\alpha+\alpha) = 1+\alpha = 1.1$


Now $\pi = CST(1+\alpha)$
 $= 0.1 \times 0.0821 \times 300 \times 1.1$
 $= 2.71 \text{ atm}$



meq. of $C_2O_4^{2-} = \text{meq. of } KMnO_4$

$M \times 2 \times 300 = 0.001 \times 5 \times 6$

$\therefore M_{C_2O_4^{2-}} = 5 \times 10^{-5}$



$K_{sp} = [Ca^{2+}][C_2O_4^{2-}] = 5 \times 10^{-5} \times 5 \times 10^{-5}$
 $= 25 \times 10^{-10}$

67. (b) $[Cl^-]$ in mixture = $0.05 + 2 \times 0.05 = 0.15 M$

from $NaCl$ from $BaCl_2$




$K_{sp} = [Ag^+][Cl^-] = S \times 0.15 = 1.0 \times 10^{-10}$

$S = 6.6 \times 10^{-10}$


68. (d) After mixing the two $[N_3H] = \frac{100 \times 0.5}{500} = 10^{-1}$

$[HOCN] = \frac{400 \times 0.1}{500} = 8 \times 10^{-2}$



at eq. $(0.1-x) \quad x \quad x+y$

$K_{N_3H} = \frac{x(x+y)}{(0.1-x)}$



$(0.08-y) \quad y \quad x+y$

$K_{HOCN} = \frac{y(x+y)}{0.08-y}$

[Both are weak, thus $0.1-x \approx 0.1$, $0.08-y \approx 0.08$]

$\therefore \frac{K_{N_3H}}{K_{HOCN}} = \frac{x(x+y)/0.1}{y(x+y)/0.08}$

or $\frac{3.6 \times 10^{-4}}{8 \times 10^{-4}} = \frac{0.08x}{0.1y}$

- or $\frac{x}{y} = \frac{3.6 \times 10^{-5}}{6.4 \times 10^{-5}} = 0.5625$
- Also, $8 \times 10^{-4} = \frac{y(x+y)}{0.08} = \frac{y(0.5625y+y)}{0.08}$
- or $1.5625 y^2 = 8 \times 10^{-4} \times 0.08$
- or $y = \sqrt{\frac{8 \times 10^{-4} \times 0.08}{1.5625}} = 6.4 \times 10^{-3} = [\text{OCN}^-]$
- $\therefore x = 0.5625 \times 6.4 \times 10^{-3} = 3.6 \times 10^{-3} = [\text{N}_3^-]$
- $\text{H}^+ = 6.4 \times 10^{-3} + 3.6 \times 10^{-3} = 10 \times 10^{-3} = 10^{-2}$
- $\therefore \text{pH} = 2$
69. (a) Given $2\text{NH}_3 \rightleftharpoons \text{NH}_4^+ + \text{NH}_2^-$
- $$K = \frac{[\text{NH}_4^+][\text{NH}_2^-]}{[\text{NH}_3]^2 (l)}$$
- or $3.46 \times 10^{-27} = \frac{[\text{NH}_4^+][\text{NH}_2^-]}{(58.82)^2}$
- $$[\text{NH}_3] = \frac{1000}{17} = 58.82 \text{ M}$$
- $\therefore [\text{NH}_4^+] = [\text{NH}_2^-] = 3.46 \times 10^{-12} \text{ M}$
- $$= 6.023 \times 10^{23} \times 3.46 \times 10^{-12}$$
- $$= 2.08 \times 10^{12} \text{ anions / litre}$$
- $$= 2.08 \times 10^9 \text{ anions / mL}$$
70. (c) K_{sp} of $\text{AgCl} = 1 \times 10^{-10} = S^2$
- $\therefore S = 10^{-5} \text{ mole/l} = 10^{-5} \times 143.5 \text{ g/l}$
- $$= 1.435 \times 10^{-3} \text{ g/l} = 1.435 \text{ mg/l}$$
- $\therefore 1.435 \text{ g of AgCl requires 1 litre}$
- $\therefore 4.305 \text{ g of AgCl requires } \frac{1 \times 4.305}{1.435} = 3 \text{ litre}$
71. (c) $\text{KH}_2\text{PO}_4 + \text{H}_3\text{PO}_4, \text{KH}_2\text{PO}_4 + \text{K}_2\text{HPO}_4$
and $\text{K}_2\text{HPO}_4 + \text{K}_3\text{PO}_4$
72. (d) $[\text{Ag}^+] = \frac{K_{sp}}{[\text{Cl}^-]}$
- Addition of Cl^- shows a decrease in $[\text{Ag}^+]$.
- or $[\text{Ag}^+] \propto \frac{1}{[\text{Cl}^-]}$ (K_{sp} is constant)
73. (d) PbSO_4 is insoluble. H_2S precipitates both CuS and PbS .
74. (a) AgF is soluble in water.
75. (c) K_2SO_4 and $(\text{NH}_4)_2\text{CO}_3$ both are water soluble
- $$\text{K}_2\text{CO}_3 + (\text{NH}_4)_2\text{SO}_4 \longrightarrow \text{K}_2\text{SO}_4 + (\text{NH}_4)_2\text{CO}_3$$
76. (a) pH of $0.1 \text{ M HCl} + 0.1 \text{ M CH}_3\text{COOH}$ is lowest as it has strong acid and CH_3COOH .
77. (a) $[\text{H}^+] = 10^{-4}$
 $[\text{H}^+] = 10^{-1}$
- 10^3 times more acidic than $[\text{H}^+] = 10^{-4}$
78. (a) $[\text{H}^+] = 3 \times 10^{-3}$
- $\therefore \text{pH} = 3 - \log 3$
- $\text{pOH} = 14 - 3 + \log 3 = 11 + \log 3$
79. (b) meq. of dil solution = meq. of conc. solution
- $$N \times 1000 = 10 \times 10^{-4}$$
- $\therefore N = \frac{10^{-3}}{1000} = 10^{-6}$
- $$\text{H}^+ \approx 10^{-6} + 10^{-7} = 1.01 \times 10^{-7}$$
- $\therefore \text{pH} > 6$
80. (a) meq. of $\text{HCl} = 10 \times 0.1 = 1$
meq. of $\text{Sr}(\text{OH})_2 = 10 \times 0.01 \times 2 = 0.2$
meq. of HCl left = $1 - 0.2 = 0.8$
- $\therefore N_{\text{HCl}} = \frac{0.8}{10+10} = \frac{0.8}{20} = 0.04$
- $\therefore \text{pH} = -\log \frac{4}{100} = -\log 4 + \log 100$
- $$= 2 - 0.6020 = 1.3979$$
81. (d) Lesser amount of energy is required for dissociation of acetic acid or acetic acid is stronger than HCN .
82. (a) $\alpha = \sqrt{\frac{K_a}{C}}$
- $\therefore \alpha_1 = \sqrt{\frac{K_a}{0.1}}$
- $$\alpha_2 = \sqrt{\frac{K_a}{0.001}}$$
- or $\frac{\alpha_1}{\alpha_2} = \sqrt{\frac{0.001}{0.1}} = 10^{-1} = 0.1$
83. (d) $\text{H}_2\text{O} + \text{H}^+ \longrightarrow \text{H}_3\text{O}^+$
 $\text{H}_3\text{O}^+ + \text{H}_2\text{O} \longrightarrow \text{H}_5\text{O}_2^+$
 $\text{H}_2\text{O} + \text{OH}^- \longrightarrow \text{H}_3\text{O}_2^-$
84. (c) $h = \sqrt{\frac{k_w}{k_a \cdot k_b}} = \sqrt{\frac{10^{-14}}{10^{-5} \times 10^{-4}}} = \sqrt{10^{-5}}$
- $$= 3.16 \times 10^{-3}$$
- $$\text{pH} = \frac{1}{2} \text{p}K_w + \frac{1}{2} \text{p}K_a - \frac{1}{2} \text{p}K_b$$
- $$= \frac{14}{2} + \frac{4}{2} - \frac{5}{2}$$
- $$= 9 - 2.5 = 6.5$$
85. (a) Solubility of AgCl in NaCl is lower than H_2O . Also other orders are derived by Debye-Huckel limiting law.
86. (a) $\text{AgNO}_3 + \text{KCN} \longrightarrow \text{AgCN} \downarrow + \text{KNO}_3$
 $\text{AgCN} + \text{KCN} \longrightarrow \text{KAg}(\text{CN})_2$
- or $\text{AgNO}_3 + 2\text{KCN} \longrightarrow \text{K}[\text{Ag}(\text{CN})_2] + \text{KNO}_3$
- For $[\text{Ag}(\text{CN})_2]^- \rightleftharpoons \text{Ag}^+ + 2\text{CN}^-$
- $\therefore K_C = \frac{[\text{Ag}^+][\text{CN}^-]^2}{[\text{Ag}(\text{CN})_2]^-} = \frac{[\text{Ag}^+][0.04]^2}{[0.03]}$
- $$= 4 \times 10^{-19}$$
- $[\text{Ag}^+] = 7.5 \times 10^{-18} \text{ M}$

$$87. (b) \quad 2.303 \log \frac{K_{sp2}}{K_{sp1}} = \frac{\Delta H}{R} \left[\frac{T_2 - T_1}{T_1 T_2} \right]$$

$$2.303 \log \frac{1.39 \times 10^{-8}}{7.47 \times 10^{-9}} = \frac{\Delta H}{2} \times \left[\frac{298 - 288}{288 \times 298} \right]$$

$$\Delta H = 10.66 \text{ k cal mol}^{-1}$$

$$88. (a) \quad \alpha = \frac{\wedge_m}{\wedge_\infty} = \frac{60}{400} = \frac{3}{20} = 0.15$$

$$\text{Now } [H^+] = C\alpha = 0.1 \times \frac{3}{20}$$

$$\therefore \text{pH} = -\log H^+$$

$$= -\log \frac{3}{200} = -0.4771 + 2.3010$$

$$= 1.8239$$

$$\text{Also } K_a = \frac{C\alpha^2}{(1-\alpha)} = \frac{0.1 \times (0.15)^2}{(1-0.15)}$$

$$= 2.65 \times 10^{-3}$$

$$89. (c) \quad HA \rightleftharpoons H^+ + A^-$$

Acid Conjugate base

$$pK_{aHA} + pK_{bA^-} = 14$$

$$\therefore pK_{aHA} = \frac{14}{2} = 7 \quad (\because pK_{aHA} + pK_{bA^-})$$

$$\therefore K_a = 10^{-7}$$

$$[H^+] = \sqrt{K_a \times C} = \sqrt{10^{-7} \times 0.001} = 10^{-5}$$

$$\therefore \text{pH} = 5$$

$$90. (b) \text{ Maximum buffer capacity} = pK_a \pm 1$$

This is possible when $\frac{[\text{conjugate base}]}{[\text{acid}]} = 1$

Thus milli mole of acetic acid taken = $500 \times 1 = 500$
 milli mole of acetic acid to be neutralised = 250
 $\therefore$ milli mole of KOH required = 250

$$\frac{w}{56} \times 1000 = 250$$

$$w_{\text{KOH}} = 14 \text{ g}$$

$$91. (b) \text{ Let solubility of HCOOAg be } a \text{ mol/litre}$$

$$\begin{array}{ccc} H^+ + HCOO^- & \rightleftharpoons & HCOOH \\ 10^{-3} - a & & a \\ 10^{-3} - a & & a \end{array}$$

(buffer)

Since HCOOH is a weak acid and has low degree of dissociation. Also its dissociation is suppressed in presence of H^+ . Thus,

$$HCOOH \rightleftharpoons H^+ + HCOO^-$$

$$\text{or } K_a = \frac{[H^+][HCOO^-]}{[HCOOH]}$$

$$[HCOO^-] = \frac{K_a \times [HCOOH]}{[H^+]} = \frac{10^{-5} \times a}{10^{-3}}$$

$$= a \times 10^{-2}$$

Also for HCOOAg:

$$HCOOAg(s) \rightleftharpoons HCOO^-(aq.) + Ag^+(aq.)$$

$$K_{sp} = S^2 = (10^{-2})^2 = 10^{-4}$$

Also $K_{sp} = [HCOO^-][Ag^+]$


$$10^{-4} = a \times 10^{-2} \times a$$

$$\therefore a = 10^{-1} = 0.1 \text{ M}$$

92. (a) Lesser is K_a , weaker is acid (HPO_4^{2-}), stronger is its conjugate base (PO_4^{3-})

$$93. (c) \quad [CaA_2] = \frac{0.125}{0.5} = 0.25 \text{ M}$$

$$[A^-] = 2 \times 0.25 = 0.50 \text{ M}$$



$$[OH^-] = C \cdot h = C \sqrt{\frac{K_h}{C}}$$

$$= \sqrt{\frac{K_w \cdot C}{K_a}} = \sqrt{\frac{10^{-14} \times 0.5}{8 \times 10^{-4}}}$$

$$= \sqrt{6.25 \times 10^{-12}} = 2.5 \times 10^{-6}$$

$$\text{pOH} = 5.60$$

$$\text{pH} = 14 - 5.60 = 8.40$$

or

$$94. (b) E_{\text{Cell}} = E_{OP_{Ag^+/Ag}}^\circ - \frac{0.059}{2} \log [Ag^+]_{LHS} + E_{RP_{Ag^+/Ag}}^\circ$$

$$+ \frac{0.059}{1} \log [Ag^+]_{RHS}$$

$\therefore$ at equilibrium $E_{\text{Cell}} = 0$ and $E_{OP_{Ag^+/Ag}}^\circ = -E_{RP_{Ag^+/Ag}}^\circ$

$$0 = \frac{0.059}{2} \log \frac{[Ag^+]_{RHS}}{[Ag^+]_{LHS}}$$

or $\frac{[Ag^+]_{RHS}}{[Ag^+]_{LHS}} = 1$ or $[Ag^+]_{RHS} = [Ag^+]_{LHS}$

At LHS:

$$[Ag^+]^2 [CO_3^{2-}] = K_{sp} Ag_2CO_3 = 8 \times 10^{-12}$$

$$\therefore [Ag^+] = \sqrt{\frac{8 \times 10^{-12}}{[CO_3^{2-}]}} = \frac{2\sqrt{2} \times 10^{-6}}{\sqrt{[CO_3^{2-}]}}$$

At RHS: $[Ag^+][Br^-] = K_{sp} AgBr = 4 \times 10^{-13}$

$$\therefore [Ag^+] = \frac{4 \times 10^{-13}}{[Br^-]}$$

$$\therefore \frac{4 \times 10^{-13}}{[Br^-]} = \frac{2\sqrt{2} \times 10^{-6}}{\sqrt{[CO_3^{2-}]}}$$

or $\frac{[Br^-]}{\sqrt{[CO_3^{2-}]}} = \sqrt{2} \times 10^{-7}$

$$95. (d) \quad Ag_2C_2O_4(s) \rightleftharpoons 2Ag^+ + C_2O_4^{2-}$$

Given $[Ag^+] = 2s = 2.2 \times 10^{-4}$

$$s = 1.1 \times 10^{-4}$$

$$K_{sp} = (2s)^2 \times s = 4s^3$$

$$= 4(1.1 \times 10^{-4})^3 = 5.3 \times 10^{-12}$$

● PREVIOUS YEARS PROBLEMS ●

- The correct acidic strength order is : (IIT 2001)
 - $\text{HClO} < \text{HClO}_2 < \text{HClO}_3 < \text{HClO}_4$
 - $\text{HClO}_4 < \text{HClO}_3 < \text{HClO}_2 < \text{HClO}$
 - $\text{HClO} < \text{HClO}_4 < \text{HClO}_3 < \text{HClO}_2$
 - $\text{HClO}_4 < \text{HClO}_2 < \text{HClO}_3 < \text{HClO}$
- For a sparingly soluble salt A_pB_q , the relationship of its solubility product L_s with its solubility (S) is : (IIT 2001)
 - $L_s = S^{p+q} \cdot p^p \cdot q^q$
 - $L_s = S^{p+q} \cdot p^q \cdot q^p$
 - $L_s = S^{pq} \cdot p^p \cdot q^q$
 - $L_s = S^{pq} \cdot (pq)^{p+q}$
- The correct order of acidic strength is : (IIT 2002)
 - $\text{CaO} < \text{CuO} < \text{H}_2\text{O} < \text{CO}_2$
 - $\text{H}_2\text{O} < \text{CuO} < \text{CaO} < \text{CO}_2$
 - $\text{CaO} < \text{H}_2\text{O} < \text{CuO} < \text{CO}_2$
 - $\text{H}_2\text{O} < \text{CO}_2 < \text{CaO} < \text{CuO}$
- H_3BO_3 is : (IIT 2003)
 - monobasic and weak Lewis acid
 - monobasic and weak Bronsted acid
 - monobasic and strong Lewis acid
 - tribasic and weak Bronsted acid
- A solution which is $10^{-3} M$ each in Mn^{2+} , Fe^{2+} , Zn^{2+} and Hg^{2+} is treated with $10^{-16} M$ sulphide ion. If K_{sp} of MnS , FeS , ZnS and HgS are 10^{-15} , 10^{-23} , 10^{-20} and 10^{-45} respectively which one will precipitate first ? (IIT 2003)
 - FeS
 - MgS
 - HgS
 - ZnS
- A weak-acid HX has the dissociation constant $1 \times 10^{-5} M$. It forms a salt NaX on reaction with alkali. The degree of hydrolysis of $0.1 M$ solution of NaX is : (IIT 2004)
 - 0.0001%
 - 0.01%
 - 0.1%
 - 0.15%
- 0.1 mole of CH_3NH_2 ($K_b = 5 \times 10^{-4}$) is mixed with 0.08 mole of HCl and diluted to one litre. The $[\text{H}^+]$ in solution is : (IIT 2005)
 - $8 \times 10^{-2} M$
 - $8 \times 10^{-11} M$
 - $1.6 \times 10^{-11} M$
 - $8 \times 10^{-5} M$
- The species present in solution when CO_2 is dissolved in water : (IIT 2006)
 - $\text{CO}_2, \text{H}_2\text{CO}_3, \text{HCO}_3^-, \text{CO}_3^{2-}$
 - $\text{H}_2\text{CO}_3, \text{CO}_3^{2-}$
 - $\text{CO}_3^{2-}, \text{HCO}_3^-$
 - $\text{CO}_2, \text{H}_2\text{CO}_3$
- 2.5 mL of $\frac{2M}{5}$ weak monoacidic base ($K_b = 1 \times 10^{-12}$ at 25°C) is titrated with $\frac{2M}{15}$ HCl in water at 25°C . The concentration of H^+ at equivalence point is : (IIT 2008)
 - $3.7 \times 10^{-13} M$
 - $3.2 \times 10^{-7} M$
 - $3.2 \times 10^{-2} M$
 - $2.7 \times 10^{-2} M$
- Solubility product K_{sp} of salt of type MX , MX_2 and M_3X at temperature T are 4.0×10^{-8} , 3.2×10^{-14} and 2.7×10^{-15} respectively. Solubilities of these salts in mol dm^{-3} at temperature T are in the order : (IIT 2008)
 - $\text{MX} > \text{MX}_2 > \text{M}_3\text{X}$
 - $\text{M}_3\text{X} < \text{MX}_2 > \text{MX}$
 - $\text{MX}_2 > \text{M}_3\text{X} > \text{MX}$
 - $\text{MX} > \text{M}_3\text{X} > \text{MX}_2$
- The dissociation constant of a substituted benzoic acid at 25°C is 1.0×10^{-4} . The pH of a $0.01 M$ solution of its sodium salt is : (IIT 2009)
 - 10
 - 8
 - 12
 - 6
- How many litres of water must be added to 1 litre of an aqueous solution of HCl with a pH of 1 to create an aqueous solution with pH of 2 ? [JEE (Main) 2013]
 - 2.0 L
 - 9.0 L
 - 0.1 L
 - 0.9 L
- The K_{sp} of Ag_2CrO_4 is 1.1×10^{-12} at $298 K$. The solubility (in mol/L) of Ag_2CrO_4 in a $0.1 M$ AgNO_3 solution is : [JEE (Advanced) I 2013]
 - 1.1×10^{-11}
 - 1.1×10^{-10}
 - 1.1×10^{-12}
 - 1.1×10^{-9}
- The initial rate of hydrolysis of methyl acetate ($1M$) by a weak acid (HA , $1M$) is $1/100^{\text{th}}$ of that of a strong acid (HX , $1M$), at 25°C . The K_a of HA is : [JEE (Advanced) II 2013]
 - 1×10^{-4}
 - 1×10^{-5}
 - 1×10^{-6}
 - 1×10^{-3}
- $\text{p}K_a$ of a weak acid (HA) and $\text{p}K_b$ of a weak base (BOH) are 3.2 and 3.4, respectively. The pH of their salt (AB) solution is : [JEE (Main) 2017]
 - 7.2
 - 6.9
 - 7.0
 - 1.0
- In the following reactions, ZnO is respectively acting as a/an : [JEE (Main) 2017]
 - $\text{ZnO} + \text{Na}_2\text{O} \longrightarrow \text{Na}_2\text{ZnO}_2$
 - $\text{ZnO} + \text{CO}_2 \longrightarrow \text{ZnCO}_3$
 - base and acid
 - base and base
 - acid and acid
 - acid and base

SOLUTIONS (Previous Years Problems)

1. (a) Higher is the oxidation number of central non-metal, more is acidic nature.
2. (a) $A_p B_q \rightleftharpoons pA^{q+} + qB^{p-}$

$$K_{sp} = \frac{pS}{(pS)^p} \frac{qS}{(qS)^q}$$

$$= p^p \cdot q^q \cdot (S)^{p+q}$$
3. (a) CaO is strongest base, CO₂ is acidic. Also, CuO is strong base than H₂O.
4. (a) $H_3BO_3 + OH^- \longrightarrow B(OH)_4^- + H^+$
 Accepts lone pair of electron thus Lewis acid but weak.
5. (c) K_{sp} of HgS is minimum and thus possesses less solubility. Note all salts are AB type.
6. (b)
$$h = \sqrt{\frac{K_H}{C}} = \sqrt{\frac{K_w}{K_a \cdot C}}$$

$$= \sqrt{\frac{10^{-14}}{10^{-5} \times 0.1}} = \sqrt{10^{-8}}$$

$$= 10^{-4}$$
 or $h = 10^{-4} \times 100 = 10^{-2} = 0.01\%$
7. (b) $CH_3NH_2 + HCl \longrightarrow CH_3NH_3^+ Cl^-$

0.1	0.08	0
0.02	0	0.08

This is basic buffer solution.

$$[OH^-] = K_b \times \frac{[base]}{[conjugate\ acid]}$$

$$= 5 \times 10^{-4} \times \frac{0.02}{0.08} = 1.25 \times 10^{-4}$$

$$\therefore [H^+] = \frac{10^{-14}}{[OH^-]} = \frac{10^{-14}}{1.25 \times 10^{-4}} = 8 \times 10^{-11} M$$
8. (a) All the following equilibrium exist together

$$CO_2 + H_2O \rightleftharpoons H_2CO_3$$

$$H_2CO_3 \rightleftharpoons H^+ + HCO_3^-$$

$$HCO_3^- \rightleftharpoons H^+ + CO_3^{2-}$$
9. (d) $BOH + HCl \longrightarrow BCl + H_2O$
 Meq. of BOH = Meq. of HCl = Meq. of BCl
 $2.5 \times \frac{2}{5} \times 1 = V \times \frac{2}{15} \times 1 = 1$
 $\therefore V = 7.5 \text{ mL}$
 $\therefore \text{Total volume} = 2.5 + 7.5 = 10 \text{ mL}$
 Thus, $[BCl] = \frac{1}{10} = 0.1$
 Now for hydrolysis of BCl; $K_H = \frac{Ch^2}{1-h} = \frac{K_w}{K_b}$
- $\therefore h = 0.27$
 or $[H^+] = c \cdot h = 0.27 \times 0.1 = 2.7 \times 10^{-2} M$
10. (d) Solubility of MX , MX_2 and M_3X are respectively

$$S = \sqrt{K_{sp}} = \sqrt{4 \times 10^{-8}} = 2 \times 10^{-4} M$$

$$S = \sqrt[3]{\frac{K_{sp}}{4}} = \sqrt[3]{\frac{3.2 \times 10^{-14}}{4}} = 2 \times 10^{-5} M$$

$$S = \sqrt[4]{\frac{K_{sp}}{27}} = \sqrt[4]{\frac{2.7 \times 10^{-15}}{27}} = 1 \times 10^{-4} M$$
11. (b) $C_6H_5COO^- + H_2O \rightleftharpoons C_6H_5COOH + OH^-$
 $\therefore [OH^-] = ch = c \sqrt{\frac{K_H}{c}} = \sqrt{\frac{K_w \cdot c}{K_a}}$

$$= \sqrt{\frac{10^{-14} \times 0.01}{1 \times 10^{-4}}} = 10^{-6}$$
 $\therefore [H^+] = 10^{-8} \text{ or } pH = 8$
12. (b) meq. of I HCl = meq. of II HCl
 $1 \times 10^{-1} = V \times 10^{-2}$
 $\therefore V = 10 \text{ litre}$
 $\therefore \text{volume of water added} = (10 - 1) = 9L$
13. (b) Ionic product of $Ag_2CrO_4 = [Ag^+]^2 [CrO_4^{2-}]$
 $= (2S + 0.1)^2 \times S = 1.1 \times 10^{-12}$
 neglecting S as compared to 0.1
 $S = 1.1 \times 10^{-10}$
14. (a) Rate = $K[\text{Ester}][H^+]$
 For strong acid $[H^+] = 1 M$
 For weak acid $[H^+] = \sqrt{K_a \cdot C}$

$$\frac{\text{Rate}_1}{\text{Rate}_2} = \frac{[H^+]_1 \text{ for strong acid}}{[H^+]_2 \text{ for weak acid}}$$

$$\frac{1}{1/100} = \frac{1}{\sqrt{K_a \times 1}} \quad (C = 1M \text{ for weak acid})$$

$$K_a = 10^{-4} M$$
15. (b) $pH = 7 + \frac{1}{2}(pK_a - pK_b)$
 $= 7 + \frac{1}{2}(3.2 - 3.4)$
 $= 6.9$
16. (d) In (a), Na₂O is basic oxide and thus ZnO acts as acidic oxide.
 In (b), CO₂ is acidic oxide and thus ZnO acts as basic oxide.

OBJECTIVE PROBLEMS (More Than One Answer Correct)

- Which one is not acidic salts ?
 (a) Na_2HPO_4 (b) NaH_2PO_2
 (c) Na_2HPO_3 (d) NaH_2PO_4
- The resulting mixtures which act buffer are :
 (a) 10 mL 0.1 M HCl + 20 mL 0.1 M NaCN
 (b) 10 mL 0.1 M NaOH + 20 mL 0.1 M NH_4CN
 (c) 10 mL 0.1 M NH_4OH + 20 mL 0.1 M $\text{CH}_3\text{COONH}_4$
 (d) 10 mL 0.1 M CH_3COOH + 20 mL 0.1 M $\text{CH}_3\text{COONH}_4$
- Which one is correct for the saturated solution of $\text{Ca}_3(\text{PO}_4)_2$ salt if its K_{sp} is 2.05×10^{-33} ?
 (a) Solubility of Ca_3PO_4 is $1.63 \times 10^{-6} \text{ M}$
 (b) $[\text{Ca}^{2+}]_{eq.} = 4.9 \times 10^{-6} \text{ M}$
 (c) $[\text{PO}_4^{3-}]_{eq.} = 3.26 \times 10^{-6} \text{ M}$
 (d) $[\text{Ca}_3\text{PO}_4]_{eq.} = \text{Zero}$
- Which of the following are correct ?
 (a) Aniline is a weak-acid in acetic acid
 (b) Ammonium salts act as acid in liquid NH_3
 (c) The reaction $\text{CsF} + \text{LiI} \longrightarrow \text{CsI} + \text{LiF}$ is acid-base reaction
 (d) HCl acts as base in HF.
- Which of the following acid-base reactions are possible?
 (a) $\text{PH}_3 + \text{NH}_4^+ \longrightarrow \text{PH}_4^+ + \text{NH}_3$
 (b) $\text{NH}_3 + \text{PH}_4^+ \longrightarrow \text{NH}_4^+ + \text{PH}_3$
 (c) $(\text{CH}_3)_3\text{P} + \text{NH}_4^+ \longrightarrow (\text{CH}_3)_3\text{P}^+\text{H} + \text{NH}_3$
 (d) $(\text{CH}_3)_3\text{N} + \text{PH}_4^+ \longrightarrow (\text{CH}_3)_3\text{NH}^+ + \text{PH}_3$
- Select the correct statements :
 (a) All Bronsted bases are Lewis bases
 (b) All Bronsted acids are Lewis acids
 (c) All Arrhenius acids are Bronsted acids
 (d) All Arrhenius bases are Bronsted bases
- The correct acidic orders are :
 (a) $\text{Li}^+ > \text{Na}^+ > \text{K}^+$ (b) $\text{Li}^+ < \text{Be}^{2+} < \text{B}^{3+}$
 (c) $\text{Fe}^{3+} > \text{Fe}^{2+} > \text{Fe}^+$ (d) $\text{NF}_3 < \text{NCl}_3 < \text{NBr}_3$
- Select the correct statements :
 (a) HCN is weak acid
 (b) Reaction of $\text{HCl}_{(g)}$ and $\text{NH}_{3(l)}$ is Arrhenius acid-base reaction
 (c) Pure H_2SO_4 and HClO_4 do not conduct current but in presence of each other they are good conductor
 (d) Mn_2O_7 is acidic oxide.
- The oxo acids of P_2O_5 are :
 (a) H_3PO_4 (b) $\text{H}_4\text{P}_2\text{O}_7$
 (c) HPO_3 (d) H_3PO_3
- Which of the following statement(s) is (are) correct?
 (a) pH of $1.0 \times 10^{-18} \text{ M}$ solution of HCl is 8
 (b) The conjugate base of H_2PO_4^- is HPO_4^{2-}
 (c) Autoprotolysis constant of water increases with temperature
 (d) When a solution of a weak monoprotic acid is titrated against a strong base, at half-neutralisation point $\text{pH} = (1/2) \text{p}K_a$
- A buffer solution can be prepared from a mixture of :
 (a) sodium acetate and acetic acid in water
 (b) sodium acetate and hydrochloric acid in water
 (c) ammonia and ammonium chloride in water
 (d) ammonia and sodium hydroxide in water
- In a buffer solution of NaH_2PO_4 and Na_2HPO_4 :
 (a) NaH_2PO_4 is acid and Na_2HPO_4 is salt
 (b) $\text{pH} = \text{p}K_2$ of $\text{H}_3\text{PO}_4 + \log \frac{[\text{HPO}_4^{2-}]}{[\text{H}_2\text{PO}_4^-]}$
 (c) Na_2HPO_4 is acid and NaH_2PO_4 is salt
 (d) $\text{pH} = \text{p}K_3$ of $\text{H}_3\text{PO}_4 + \log \frac{[\text{H}_2\text{PO}_4^-]}{[\text{HPO}_4^{2-}]}$
- Select the correct statements :
 (a) The K_{a1} values for H_2SO_3 is 1.3×10^{-2}
 (b) H_2SO_3 exist in only minute concentration in aqueous solution of SO_2
 (c) The K_{a1} , values of H_2SO_3 refers for the process $\text{H}_2\text{SO}_3 \rightleftharpoons \text{H}^+ + \text{HSO}_3^-$
 (d) The K_a , value of H_2SO_3 refers for the process $\text{SO}_2 + \text{H}_2\text{O} \rightleftharpoons \text{H}^+ + \text{HSO}_3^-$
 (g) (l) (aq.) (aq.)
- Select the correct statements :
 (a) HF is a weak acid
 (b) The strength of weak acids increases with dilution ($\text{HA} \rightleftharpoons \text{H}^+ + \text{A}^-$)
 (c) The strength of HF increases with concentration ($\text{HF} \rightleftharpoons \text{H}^+ + \text{F}^-$)
 (d) The F^- furnished by HF reacts with HF to give HF_2^- and thereby shifting the reaction to right
- An aqueous solution of HNO_3 , KOH , CH_3COOH and CH_3COONa of identical concentrations are provided. The pairs of solution which form a buffer upon mixing is/are :
 (a) HNO_3 and CH_3COOH
 (b) KOH and CH_3COONa
 (c) HNO_3 and CH_3COONa
 (d) CH_3COOH and CH_3COONa

(IIT 2010)

16. The correct statement(s) for orthoboric acid is/are :

[JEE (Advanced) 2014]

(a) It behaves as a weak acid in water due to self ionization.

(b) Acidity of its aqueous solution increases upon addition of ethylene glycol.

(c) It has a three dimensional structure due to hydrogen bonding.

(d) It is a weak electrolyte in water.

SOLUTIONS (More Than One Answer Correct)

1. (b,c) H_3PO_2 is monobasic and H_3PO_3 is dibasic.

2. (a,b,c,d) $\text{HCl} + \text{NaCN} \longrightarrow \text{NaCl} + \text{HCN}$

$$\begin{array}{ccccccc} 10 \times 0.1 & 20 \times 0.1 & & & & & \\ = 1 & = 2 & & 0 & & 0 & \\ 0 & 1 & & 1 & & 1 & \end{array}$$

$$[\text{HCN}] = \frac{1}{30} \text{ and } [\text{NaCN}] = \frac{1}{30}$$

3. (a,b,c,d) $\text{Ca}_3(\text{PO}_4)_2(s) \rightleftharpoons 3\text{Ca}^{2+} + 2\text{PO}_4^{3-}$

$$\therefore K_{sp} = 2.05 \times 10^{-33} = 108s^5$$


$$\therefore s = 1.63 \times 10^{-6} M$$

$$\therefore [\text{Ca}^{2+}] = 3 \times 1.63 \times 10^{-6} = 4.9 \times 10^{-6} M$$


$$[\text{PO}_4^{3-}] = 2 \times 1.63 \times 10^{-6} = 3.26 \times 10^{-6} M$$

$[\text{Ca}_3\text{PO}_4]_{eq} = 0$, because all the salt is in ionic state.

4. (b,c,d) Aniline is strong base in acetic acid. $\text{NH}_3(l)$ ionises as :



This reaction is an interesting example of soft acid-hard base and hard acid-soft base combination





5. (b,c) Proton affinity of NH_3 is greater than PH_3 . Proton affinity of $(\text{CH}_3)_3\text{P}$ is greater than NH_3 .

6. (a,c) NaOH is Arrhenius base but not Bronsted base. HCl is Bronsted acid but not Lewis acid.

7. (a,b,c,d) Higher is effective nuclear charge more is acidic nature of Lewis acid cations. The increasing electronegativity from I to F in NX_3 give rise to lesser tendency for N to donate electron pair as it acquires more +ve charge on N-atom.

8. (a,b,c,d) Arrhenius concept is in aqueous medium



9. (a,b,c) Each has P^{5+} state.

10. (b,c) HCl is acid; for (d) $\text{pH} = \text{p}K_a$

11. (a,b,c) For (b) $\text{CH}_3\text{COONa} + \text{HCl} \rightarrow \text{CH}_3\text{COOH} + \text{NaCl}$

more less


12. (a,b) HPO_4^{2-} is conjugate base of H_2PO_4^- and thus.

13. (a,b,d) Choice (b) is correct explanation for H_2SO_3 .


14. (a,b,c,d) All are facts. Choice (d) is correct explanation for HF in aqueous medium.

15. (c,d) CH_3COONa (in more amount) than HNO_3 forms a buffer mixture of $(\text{CH}_3\text{COONa} + \text{CH}_3\text{COOH})$ in solution.

16. (b,d) Orthoboric acid H_3BO_3 is a weak monobasic Lewis acid as it accepts OH^- ions



The equilibrium is shifted in forward direction by the addition of ethylene glycol (a syndiol) which forms a stable complex with $\text{B}(\text{OH})_4^-$.



It has a planar sheet like structure due to sp^2 hybridization of B atom as well as hydrogen bonding.

COMPREHENSION BASED PROBLEMS

Comprehension 1 : Solubility of a substance in its saturated solution can be derived from its K_{sp} values. Higher is the K_{sp} for same type of compound more is the solubility. If S is the solubility in mol/litre then K_{sp} of a compound A_xB_y is expressed as

$$K_{sp} = x^x \cdot y^y [S]^{x+y}$$

Consider a compound $M(OH)_x$ having $K_{sp} = 27 \times 10^{-12}$ and solubility in pure water is 10^{-3} mol litre⁻¹.

- [1] The value of x is :
 (a) 1 (b) 2
 (c) 3 (d) 4
- [2] The solubility (in M) of $M(OH)_x$ in 0.1 M NaOH solution is :
 (a) 2.7×10^{-10} (b) 2.7×10^{-3}
 (c) 2.7×10^{-8} (d) 2.7×10^{-4}
- [3] The solubility (in M) of $M(OH)_x$ in 0.1 Ca(OH)₂ solution is :
 (a) 6.46×10^{-4} (b) 2.7×10^{-8}
 (c) 2.7×10^{-4} (d) 3.375×10^{-9}
- [4] The solubility (in mole/litre) of $M(OH)_x$ in 0.1 M (NO₃)_x solution is :
 (a) 2.15×10^{-4} (b) 2.15×10^{-5}
 (c) 2.15×10^{-6} (d) 2.15×10^{-7}

Comprehension 2: K_b of CH_3COO^- is 5.26×10^{-10} . Calculate for 0.01 N solution of sodium acetate.

- [1] Hydrolysis constant of CH_3COO^- is :
 (a) 5.26×10^{-10} (b) 5.26×10^{-11}
 (c) 5.26×10^{-12} (d) 5.26×10^{-9}
- [2] Degree of hydrolysis of CH_3COO^- is :
 (a) 2.29×10^{-6} (b) 2.29×10^{-4}
 (c) 2.29×10^{-3} (d) 2.29×10^{-5}
- [3] pH of solution is :
 (a) 8.56 (b) 5.44
 (c) 8.36 (d) 9.56


Comprehension 3: Oxides of metals are usually basic and oxides of non metals are usually acidic. However some acidic oxides of metals and neutral oxides of non metals such as N₂O, NO, CO, H₂O are known.

- [1] Which of the following oxide is acidic ?
 (a) Cr₂O₃ (b) CrO₃
 (c) MnO₂ (d) MnO
- [2] In the reaction of CO with NaOH at high P and T to give sodium formate, CO acts as :
 (a) acid (b) base
 (c) neutral (d) amphoteric

[3] Number of OH gp present in H₃PO₂ are :

- (a) 1 (b) 2
(c) 3 (d) none of these

Comprehension 4 : The dissolution of ammonia gas in water does not obey Henry's law. On dissolving, a major portion of ammonia, molecules unite with H₂O to form NH₄OH molecules. NH₄OH again dissociates into NH₄⁺ and OH⁻ ions. In solution therefore, we have NH₃ molecules, NH₄OH molecules and NH₄⁺ ions and the following equilibrium exist :



Let C_1 mol/L of NH₃ pass in solution state a part of which on dissolution in water forms C_2 mol/L of NH₄OH. The solution contains C_3 mol/L of NH₄⁺ ions.

- [1] Total concentration of ammonia, which can be determined by volumetric analysis is equal to :
 (a) $C_1 + C_2$ (b) $C_1 + C_2 + C_3$
 (c) $C_1 + C_3$ (d) $C_2 + C_3$
- [2] Concentration of undissociated ammonium hydroxide is :
 (a) $C_1 + C_2$ (b) $C_2 - C_3$
 (c) $C_1 + C_3$ (d) $C_1 - C_2$
- [3] Degree of dissociation of ammonium hydroxide is :
 (a) C_1 (b) C_3 / C_1
 (c) C_3 / C (d) C_3 / C_2
- [4] If p is the partial pressure of ammonia at equilibrium, then which of the following is constant? α is degree of dissociation of NH₃.
 (a) $\frac{P}{C_1}$ (b) $\frac{P}{C}$
 (c) $\frac{P}{C_2}$ (d) $\frac{P}{C_3}$
- [5] The dissociation constant of NH₄OH can be given as :
 (a) $K_b = \frac{(C_3)^2}{(C_2 - C_3)}$ (b) $K_b = \frac{(C_3)^2}{C_2}$
 (c) $K_b = \frac{C_3}{(C_2 - C_1)}$ (d) $K_b = \frac{C_3}{C_1 - C_2}$
- [6] The pH of solution can be given by :
 (a) $pK_w + \log C_3$
 (b) $pK_w - \frac{1}{2} pK_b + \frac{1}{2} \log C_2$
 (c) $pK_a + \frac{1}{2} \log C_2 + \frac{1}{2} pK_b$
 (d) either of these

Comprehension 5 : The pH of pure water at 25°C and 60°C are 7 and 6.5 respectively. HCl gas is passed through water at 25°C till the resulting 1 litre solution acquires a pH of 3. Now 4×10^{-3} mole of NaCN are added into this solution. Also a fresh 0.1 MHCN solution has pH 5.1936. Now in the one part of solution obtained after addition of NaCN, one millimole of NaOH are added and in the second part of this solution 0.5 millimole of HCl are added.

- [1] The heat of formation of water from H^+ and OH^- is :
(a) 13.06 kcal (b) - 13.06 kcal
(c) 16.32 kcal (d) - 16.32 kcal
- [2] The volume of HCl passed through the solution at 25°C and 1 atm is :
(a) 24.46 mL (b) 2.446 mL
(c) 244.6 mL (d) 0.2446 mL
- [3] The dissociation constant of HCN is :
(a) 4.1×10^{-10} (b) 4.1×10^{-6}
(c) 4.1×10^{-3} (d) 4.1×10^{-8}
- [4] The degree of dissociation of 0.1 M HCN solution is :
(a) 6.4×10^{-5} (b) 6.4×10^{-3}
(c) 6.4×10^{-2} (d) 6.4×10^{-6}
- [5] The pH of resulting solution after addition of NaCN is :
(a) 9.86 (b) 8.86
(c) 6.86 (d) 5.86
- [6] The pH of resulting solution after addition of 0.1 millimole of HCl is :
(a) 9.68 (b) 9.39
(c) 9.21 (d) 9.98
- [7] The pH of resulting solution after addition of 0.05 millimole of NaOH is :
(a) 10.23 (b) 11.23
(c) 9.23 (d) 8.23

SOLUTIONS

Comprehension 1

- [1] (c) $M(OH)_x \rightleftharpoons M^{+x} + xOH^-$
 $K_{sp} = x^x \cdot (S)^{1+x} = 27 \times 10^{-12}$
 $\therefore x = 3$
- [2] (c) $K_{sp} = [M^{+3}][OH^-]^3 = [M^{+3}][OH^-]^3$
 $27 \times 10^{-12} = S \times [3S + 0.1]^3 \quad [3S \ll 0.1]$
 $\therefore S = \frac{27 \times 10^{-12}}{(0.1)^3} = 2.7 \times 10^{-8}$
- [3] (d) $K_{sp} = [M^{+3}][OH^-]^3$
 $27 \times 10^{-12} = S \times [3S + 0.2]^3 \quad [3S \ll 0.2]$
 $S = \frac{27 \times 10^{-12}}{(0.2)^3} = 3.375 \times 10^{-9}$
- [4] (a) $K_{sp} = [M^{+3}][OH^-]^3$
 $= [S + 0.1][3S]^3 \quad [S \ll 0.1]$
 $27S^3 = \frac{27 \times 10^{-12}}{0.1}$
 $S = 2.15 \times 10^{-4}$

Comprehension 2

- For $CH_3COONa + H_2O \rightleftharpoons CH_3COOH + NaOH$
 Before hydrolysis 1 0 0
 After hydrolysis (1-h) h h
- K_b for $CH_3COO^- = 5.26 \times 10^{-10}$
 $\therefore K_a$ for $CH_3COOH = 1.9 \times 10^{-5}$
- [1] (a) $K_H = \frac{K_w}{K_a} = \frac{10^{-14}}{1.9 \times 10^{-5}} = 5.26 \times 10^{-10}$
- [2] (b) $h = \sqrt{\left(\frac{K_H}{C}\right)} = \sqrt{\left(\frac{5.26 \times 10^{-10}}{0.01}\right)} = 2.29 \times 10^{-4}$
- [3] (c) $[OH^-]$ from NaOH, a strong alkali = Ch
 $= 0.01 \times 2.29 \times 10^{-4}$
 $= 2.29 \times 10^{-6} M$
 $\therefore pOH = 5.64 \therefore pH = 8.36$

Comprehension 3


- [1] (b) Some metal oxides in their highest oxidation state are acidic.
- [2] (a) $CO + NaOH \xrightarrow{P.T.} HCOONa$
 acid base (salt)
- [3] (a) $HO-P(=O)(H)-H$

Comprehension 4

- [1] (b) The intermediate solution of acid will react with all the NH_3 present in solution.
- [2] (b) NH_4OH left undissociated = $C_2 - C_3$.
- [3] (d) $\alpha = \frac{\text{Mole of } NH_4OH \text{ dissociated}}{\text{Total mole}} = \frac{C_3}{C_2}$
- [4] (a) Acc. to Henry's law $a \propto P_{NH_3}$
 Where a is the amount of gas dissolved per unit volume of solvent.
 or $C_{NH_3} \propto P_{NH_3}$
 $\therefore K = \frac{P_{NH_3}}{C_{NH_3}} = \frac{P}{C_1}$
- [5] (a) $NH_4OH \rightleftharpoons NH_4^+ + OH^-$
 $\begin{matrix} C_2 & 0 & 0 \\ (C_2 - C_3) & C_3 & C_3 \end{matrix}$
 $\therefore K_b = \frac{C_3^2}{C_2 - C_3}$
- [6] (d) $NH_4OH \rightleftharpoons NH_4^+ + OH^-$
 $\begin{matrix} t=0 & C_2 & 0 & 0 \\ \text{At eq.} & C_2(1-\alpha) & C_2\alpha & C_2\alpha \end{matrix}$
 Also $C_2\alpha = C_3$
 $\therefore [OH^-] = C_3$
 or $pOH = -\log C_3$
 $\therefore pH = pK_w - pOH = pK_w + \log C_3$
 Also $[OH^-] = C_2\alpha = C_2\sqrt{\frac{K_b}{C_2}} = \sqrt{K_b \cdot C_2}$
 $\therefore pOH = -\frac{1}{2} \log K_b - \log C_2$
 $= \frac{1}{2} pK_b - \frac{1}{2} \log C_2$
 $\therefore pH = pK_w - pOH = pK_w - \frac{1}{2} pK_b + \frac{1}{2} \log C_2$
 $= pK_a + \frac{1}{2} pK_b + \frac{1}{2} \log C_2$

Comprehension 5

- [1] (b) At $25^\circ C$ $pH = 7 \therefore K_w = 10^{-14}$;
 At $60^\circ C$, $pH = 6.5 \therefore K_w = 10^{-13}$
 $H_2O \rightleftharpoons H^+ + OH^-; \Delta H = ?$
 $2.303 \log \frac{10^{-13}}{10^{-14}} = \frac{\Delta H}{R} \left[\frac{35}{333 \times 298} \right]$
 $\therefore \Delta H = 13.06 \text{ kcal}$
 $\therefore H^+ + OH^- \rightarrow H_2O; \Delta H = -13.06 \text{ kcal}$
- [2] (a) pH of solution after passage of $HCl = 3$
 $\therefore [H^+] = 10^{-3} M$ or $[HCl] = 10^{-3} M$
 From $PV = nRT$
 $1 \times V = 10^{-3} \times 0.0821 \times 298$
 $V = 24.46 \text{ mL}$




$$\text{Also } [\text{H}^+] = c \cdot \alpha = c \sqrt{\frac{K_a}{c}} = \sqrt{K_a c}$$

$$\text{or } -\log \text{H}^+ = -\frac{1}{2} [\log K_a + \log c]$$


$$5.1936 = -\frac{1}{2} [\log K_a + \log 0.1]$$

$$\therefore K_a = 4.1 \times 10^{-10}$$




$$4.1 \times 10^{-10} = 0.1 \times \alpha^2$$

$$\therefore \alpha = 6.4 \times 10^{-5}$$




$$\therefore \text{pH} = \text{p}K_a + \log \frac{[\text{conjugate base}]}{[\text{acid}]}$$

$$= 9.3872 + \log \frac{0.003}{0.001} = 9.8643$$



$$\therefore \text{pH} = 9.3872 + \log \frac{0.002}{0.002} = 9.3872$$



$$\therefore \text{pH} = 9.3872 + \log \frac{0.0035}{0.0005} = 10.2323$$

In each sub question given below a statement S and explanation E is given. Choose the correct answers from the codes a, b, c and d given for each question :

- (a) S is correct but E is wrong
 (b) S is wrong but E is correct
 (c) Both S and E are correct and E is correct explanation for S
 (d) Both S and E are correct but E is not correct explanation for S
- S : The pH of a basic buffer mixture is given by :

$$\text{pH} = \text{p}K_a + \log \frac{[\text{base}]}{[\text{conjugate acid}]}$$

 E : The pH of an acidic buffer mixture is given by :

$$\text{pH} = \text{p}K_a + \log \frac{[\text{conjugate base}]}{[\text{acid}]}$$
 - S : On passing HCl(g) through a saturated solution of BaCl_2 , a white turbidity appears.
 E : The common ion effect is responsible for white turbidity.
 - S : Degree of hydrolysis and pH of a salt say NH_4CN is independent of concentration of NH_4CN .
 E : The solution of NH_4CN in water has pH slightly greater than 7.
 - S : In a pair of two electrolytes one having higher value of K_{sp} is more soluble in water than the other having lower value of K_{sp} .
 E : Solubility of electrolyte depends upon K_{sp} as well as on the nature of electrolyte.
 - S : HgCl_2 and SnCl_2 cannot coexist in a solution.
 E : Increase in concentration of Cl^- in solution brings in precipitation of either of them.
 - S : The solubility of HgI_2 in water decreases in presence of KI.
 E : HgI_2 is insoluble in water but it becomes soluble in KI(aq) .
 - S : The pH of $\text{NH}_4\text{OH(aq)}$ decreases on addition of little NH_4Cl in it.
 E : The pH of NH_4Cl brings in decrease in degree of dissociation of NH_4OH due to common ion effect.
 - S : The $[\text{H}^+]$ in 10^{-8} N HCl is $1.05 \times 10^{-7} \text{ M}$.
 E : The H^+ is obtained from HCl and H_2O ; the later furnishes less H^+ than 10^{-7} due to common ion effect.
 - S : HCl acts as weak base in liquid HF but weak acid in acetic acid.
 E : The tendency to act as acid or base also depends upon the nature of other substance to accept or donate a proton.
 - S : Cu^+ , Ag^+ are soft acids whereas CN^- , H^- are soft bases.
 E : Soft acids do not possess noble gas configuration whereas soft bases have donor atom of easily polarised nature.
 - S : Metal oxides are usually either acidic or amphoteric but Mn_2O_7 is acidic.
 E : On dissolution in water, Mn_2O_7 forms per manganic acid.
 - S : The dissociation constants of polyprotic acid are in the order $K_1 > K_2 > K_3$.
 E : The $[\text{H}^+]$ furnished in first step of dissociation exerts common ion effect to reduce the second dissociation so on.
 - S : All strong acid in water show almost same acidic nature.
 E : This is due to levelling effect of water on account of its high dielectric constant and strong proton accepting tendency.
 - S : CCl_4 , C_6H_6 and liquid SO_2 are aprotic solvents.
 E : Aprotic solvents do not influence the acidic or basic nature of solute.
 - S : The acidic nature of some cations is :

$$\text{Al}^{3+} > \text{Be}^{2+} > \text{Na}^+ > \text{K}^+$$

 E : More is the effective nuclear charge on cation more is its acidic nature.
 - S : Acidic nature of boron trihalides is in the order :

$$\text{BF}_3 < \text{BCl}_3 < \text{BBr}_3 < \text{BI}_3$$

 E : Basic nature of nitrogen trihalides is in the order :

$$\text{NF}_3 > \text{NCl}_3 > \text{NBr}_3 > \text{NI}_3$$
 - S : $\text{CO} + \text{NaOH} \xrightarrow[\text{High } T]{\text{High } P} \text{HCOONa}$.
 E : CO although being neutral can acts as acid in the given reaction.
 - S : The dissociation constant of water at 60°C is 10^{-13} .
 E : The pH of water is 6.5 and that it behaves as acid at 60°C .
 - S : Salting out action of sodium soap in presence of NaCl is based on common ion effect.
 E : Salting out action of soap is based on the fact that as the concentration of Na^+ increases, the RCOONa shows precipitation because $[\text{RCOO}^-][\text{Na}^+] > K_{sp}$.
 - S : Hydrolysis of salt is an exothermic phenomenon.
 E : It involves breaking up of water molecule of produce acid and base respectively.

21. S : 0.1 M NaCN + 0.05 M HCl solution on mixing in equal volume form a buffer solution.
E : The solution after mixing contains a weak acid and its conjugate base and thus act as buffer.
22. S : The pH of NH_4OH remains unchanged on addition of NH_4Cl .
E : Addition of NH_4Cl suppresses the dissociation of NH_4OH due to common ion effect.
23. S : Heat given out during neutralisation of NaOH and HF is -13.7 kcal/eq .
E : F^- ion is more easily hydrated and thus heat of neutralisation of HF and NaOH is more.
24. S : The pH of pure water is less than 7 at 60°C .
E : As the temperature increases, pure water becomes slightly acidic.
25. S : The pH of human blood at body temperature is found to be 6.9.
E : Blood is alkaline in nature.
26. S : Solubility of AgCl is more in conc. HCl than in water.
E : AgCl form a complex with conc. HCl and thus solubility of AgCl increases in conc. HCl.
27. S : All Arrhenius acids are also Bronsted acids.
E : All Bronsted bases are also Lewis bases.
28. S : Cl^- is weak base than $\text{C}_2\text{H}_5\text{O}^-$.
E : Stronger is acid, weaker is its conjugate base.
29. S : H_3BO_3 in water behaves as monobasic acid.
E : The ionisation reaction is :
$$\text{H}_3\text{BO}_3 + \text{H}_2\text{O} \rightleftharpoons \text{B}(\text{OH})_4^- + \text{H}^+$$
30. S : Solubility of AgCl is less in 0.1 M NaCl than in water.
E : In presence of NaCl, the solubility of AgCl is lowered on account of common ion effect.
31. S : In water orthoboric acid behaves as a weak monobasic acid.
E : In water orthoboric acid behaves as a proton donor.
32. S : HNO_3 is a stronger acid than HNO_2 .
E : In HNO_3 there are two nitrogen-to-oxygen bonds whereas in HNO_2 there is only one.

Ionic Equilibrium

591

ANSWERS (Statement Explanation Problems)

1. (d) For basic buffer $\text{pOH} = \text{p}K_b + \log \frac{[\text{conjugate acid}]}{[\text{base}]}$
 $\therefore \text{pH} = \text{p}K_w - \text{pOH}$
2. (a) This is not common ion effect (At least one should be weak electrolyte). Here product of ionic concentration exceeds K_{sp} of BaCl_2 .
3. (d) NH_4OH is relatively stronger than HCN.
4. (b) $K_{sp} = p^p \cdot q^q \cdot s^{(p+q)}$ for $A_p B_q$.
5. (a) $\text{SnCl}_2 + 2\text{HgCl}_2 \longrightarrow \text{SnCl}_4 + \text{Hg}_2\text{Cl}_2 \downarrow$
 $\text{Hg}_2\text{Cl}_2 + \text{SnCl}_2 \longrightarrow \text{SnCl}_4 + 2\text{Hg} \downarrow$
This is redox change.
6. (b) HgI_2 forms soluble complex with KI
 $2\text{KI} + \text{HgI}_2 \longrightarrow \text{K}_2\text{HgI}_4$
7. (c) Explanation is correct reason for statement.
8. (c) —do—
9. (c) —do—
10. (c) —do—
11. (c) —do—
12. (c) Explanation is correct reason for statement.
13. (c) Explanation is correct reason for statement.
14. (c) Explanation is correct reason for statement.
15. (c) Explanation is correct reason for statement.
16. (a) Statement is correct due to back bonding in boron. In nitrogen halides the order is $\text{NF}_3 < \text{NCl}_3 < \text{NBr}_3 < \text{NI}_3$. On account of decreasing electronegativity of halogens, in NF_3 , the lone pair is not released easily, due to more +ve charge on N.
17. (c) CO is acid, NaOH is base and salt formed is HCOONa .
18. (a) $K_w = 10^{-13}$ at 60°C $\therefore \text{pH} = 6.5$; but water is neutral because pH scale contracts to 0 to 13.
19. (b) $\text{RCOONa} \rightarrow \text{RCOO}^- + \text{Na}^+$; In presence of NaCl, $[\text{Na}^+]$ increases and $[\text{RCOO}^-][\text{Na}^+]$ exceeds than K_{sp} of RCOONa .
20. (b) Breaking up of bonds is endothermic.
21. (c)
$$\begin{array}{ccccccc} \text{NaCN} + \text{HCl} & \longrightarrow & \text{NaCl} + \text{HCN} \\ 0.1 & 0.05 & 0 & 0 \\ 0.05 & 0 & 0.05 & 0.05 \end{array}$$

The solution contains a weak acid HCN and its conjugate base CN^- and thus acts as buffer.
22. (b) The dissociation of NH_4OH is suppressed in presence of NH_4Cl and thus pH of NH_4OH decreases.
23. (b) Heat given out during complete neutralisation of HF and NaOH is -16.4 kcal/eq due to extensive hydration of F^- on NaF account of its smaller size.
24. (a) The scale of pH (0 to 14 at 25°C) changes to (0 to less than 14) as the temperature rises because K_w of water increases with temperature. Note that $[\text{H}^+] = [\text{OH}^-]$ and thus water remains neutral.
25. (d) Blood is alkaline and at body temperature (98°F) scale of pH lies between 0 to 13.6.
26. (c) Explanation is correct reason for statement.
27. (a) All Bronsted bases are not Lewis bases.
28. (c) Explanation is correct reason for statement.
29. (c) Explanation is correct reason for statement.
30. (a) The equilibrium $\text{AgCl}(s) \rightleftharpoons \text{Ag}^+ + \text{Cl}^-$
 $K_{sp} = [\text{Ag}^+][\text{Cl}^-]$
if $[\text{Cl}^-]$ increases, the equilibrium is shifted in the backward direction, i.e., solubility of AgCl decreases in presence of NaCl. Note this is Le Chatelier's principle application to solubility product. In common ion effect, there must be a weak electrolyte.
31. (a) $\text{B}(\text{OH})_3 + \text{H}_2\text{O} \rightleftharpoons \text{B}(\text{OH})_4^- + \text{H}^+$
32. (a)
$$\begin{array}{ccc} & +5 & \\ & \text{O} & \\ \text{HO}-\text{N} & \searrow & \\ & \text{O} & \end{array} \quad \text{HO}-\text{N} \begin{array}{l} +3 \\ =\text{O} \end{array}$$

MATCHING TYPE PROBLEMS

Type I : Only One Match Is Possible

- | 1. | List A | List B |
|-----|----------------------|-------------------|
| (a) | Adsorption indicator | (i) $K_3Fe(CN)_6$ |
| (b) | External indicator | (ii) Starch |
| (c) | Self indicator | (iii) Methyl red |
| (d) | Acid-base indicator | (iv) $KMnO_4$ |

- | 2. | List A | List B |
|-----|--|--|
| (a) | pH of amphiprotic salt | 1. $-\frac{1}{2}[\log(K_{a1}c_1 + K_{a2}c_2)]$ |
| (b) | pH of mixtures of two weak acids | 2. $\frac{1}{2}[pK_{a1} + pK_{a2}]$ |
| (c) | pH of a basic buffer mixture | 3. $pK_a + \log \frac{[\text{conjugate base}]}{[\text{acid}]}$ |
| (d) | pH of an acidic buffer mixture | 4. $pK_a + \log \frac{[\text{base}]}{[\text{conjugate acid}]}$ |
| (e) | pH of a salt solution of weak acid + strong base | 5. $\frac{1}{2}[pK_w + pK_a + \log c]$ |
| (f) | pH of a salt solution of strong acid + weak base | 6. $\frac{1}{2}[pK_w - pK_b - \log c]$ |

- | 3. | List A | List B |
|-----|----------------------------------|--------------------------------|
| (a) | K_{sp} of $BaCl_2 \cdot 2H_2O$ | 1. $[Ba^{2+}][Cl^-]^2[H_2O]^2$ |
| (b) | K_{a2} of dibasic acid | 2. $[Ba^{2+}][Cl^-]^2$ |
| (c) | Conjugate acid of RNH_2 | 3. $[A^{-2}]$ |
| (d) | Conjugate base of RNH_2 | 4. RNH^- |
| (e) | Solubility of $AgCl$ in $NaCl$ | 5. RNH_3^+ |
| | | 6. $S = \frac{K_{sp}}{[Cl^-]}$ |

Type II : More Than One Match Possible

- | 4. | List A | List B |
|-----|------------------------|------------------------------------|
| (a) | Solubility of PbS | 1. Increases in dil. HCl_{aq} . |
| (b) | Solubility of Ag_2S | 2. Increases in dil. $NaCN_{aq}$. |
| (c) | Solubility of CuS | 3. Increases in dil. NH_3_{aq} . |
| (d) | Solubility of $BaSO_4$ | 4. Increases in conc. HCl |

Type III : Only One Match From Each List

- | 5. | List A | List B | List C |
|----|-------------------------|----------------------------------|---|
| A. | Melting of ice | 1. Favoured with decrease in P | a. $pH = \frac{1}{2}[pK_{a1} + pK_{a2}]$ |
| B. | Freezing of water | 2. Favoured with increase in P | b. $pH = \frac{1}{2}[pK_w + pK_a - pK_b]$ |
| C. | 0.1 M $NaHS(aq)$ | 3. Hydrolysis and protonation | c. Favoured by addition of ice |
| D. | 0.1 M $CH_3COONH_4(aq)$ | 4. Hydrolysis | d. Not favoured by addition of ice |
| E. | 0.1 M $NH_4Cl(aq)$ | 5. Acidic | e. $pH = \frac{1}{2}[pK_a - \log C]$ |

ANSWERS

1. a-ii; b-i; c-iv; d-iii
 2. a-2; b-1; c-4; d-3; e-5; f-6
 3. a-2; b-3; c-5; d-4; e-6;

4. a-1,3,4; b-1,2,3,4; c-1,2,3,4; d-4
 5. A-2-d; B-1-c; C-3-a; D-4-b; E-5-e