

Ratio and Proportion (Including Properties & Uses)

Question 1.

If $a : b = 5 : 3$, find: $\frac{5a - 3b}{5a + 3b}$.

Solution:

$$a : b = 5 : 3$$

$$\Rightarrow \frac{a}{b} = \frac{5}{3}$$

$$\frac{5a - 3b}{5a + 3b} = \frac{5\left(\frac{a}{b}\right) - 3}{5\left(\frac{a}{b}\right) + 3} \quad (\text{Dividing each term by } b)$$

$$= \frac{5\left(\frac{5}{3}\right) - 3}{5\left(\frac{5}{3}\right) + 3}$$

$$= \frac{\frac{25}{3} - 3}{\frac{25}{3} + 3}$$

$$= \frac{25 - 9}{25 + 9}$$

$$= \frac{16}{34} = \frac{8}{17}$$

Question 2.

If $x : y = 4 : 7$, find the value of $(3x + 2y) : (5x + y)$.

Solution:

$$x : y = 4 : 7$$

$$\Rightarrow \frac{x}{y} = \frac{4}{7}$$

$$\frac{3x+2y}{5x+y} = \frac{3\left(\frac{x}{y}\right)+2}{5\left(\frac{x}{y}\right)+1}$$

(Dividing each term by y)

$$= \frac{3\left(\frac{4}{7}\right)+2}{5\left(\frac{4}{7}\right)+1}$$

$$= \frac{\frac{12}{7}+2}{\frac{20}{7}+1}$$

$$= \frac{12+14}{20+7}$$

$$= \frac{26}{27}$$

Question 3.

If $a : b = 3 : 8$, find the value of $\frac{4a+3b}{6a-b}$.

Solution:

$$a : b = 3 : 8$$

$$\Rightarrow \frac{a}{b} = \frac{3}{8}$$

$$\frac{4a+3b}{6a-b} = \frac{4\left(\frac{a}{b}\right)+3}{6\left(\frac{a}{b}\right)-1}$$

(Dividing each term by b)

$$= \frac{4\left(\frac{3}{8}\right)+3}{6\left(\frac{3}{8}\right)-1}$$

$$\begin{aligned}
 &= \frac{\frac{3}{2} + 3}{\frac{9}{4} - 1} \\
 &= \frac{\frac{9}{2}}{\frac{5}{4}} \\
 &= \frac{18}{5}
 \end{aligned}$$

Question 4.

If $(a - b) : (a + b) = 1 : 11$, find the ratio $(5a + 4b + 15) : (5a - 4b + 3)$.

Solution:

$$\frac{a - b}{a + b} = \frac{1}{11}$$

$$11a - 11b = a + b$$

$$10a = 12b$$

$$\frac{a}{b} = \frac{12}{10} = \frac{6}{5}$$

So, let $a = 6k$ and $b = 5k$

$$\begin{aligned}
 \frac{5a + 4b + 15}{5a - 4b + 3} &= \frac{5(6k) + 4(5k) + 15}{5(6k) - 4(5k) + 3} \\
 &= \frac{30k + 20k + 15}{30k - 20k + 3} \\
 &= \frac{50k + 15}{10k + 3} \\
 &= \frac{5(10k + 3)}{10k + 3} \\
 &= 5
 \end{aligned}$$

Hence, $(5a + 4b + 15) : (5a - 4b + 3) = 5 : 1$

Question 5.

Find the number which bears the same ratio to

$$\frac{7}{33} \text{ that } \frac{8}{21} \text{ does to } \frac{4}{9}.$$

Solution:

Let the required number be $\frac{x}{y}$.

$$\text{Now, Ratio of } \frac{8}{21} \text{ to } \frac{4}{9} = \frac{\frac{8}{21}}{\frac{4}{9}} = \frac{8}{21} \times \frac{9}{4} = \frac{6}{7}$$

Thus, we have

$$\frac{\frac{x}{y}}{\frac{7}{33}} = \frac{6}{7}$$

$$\Rightarrow \frac{x}{y} = \frac{6}{7} \times \frac{7}{33}$$

$$\Rightarrow \frac{x}{y} = \frac{6}{7} \times \frac{7}{33}$$

$$\Rightarrow \frac{x}{y} = \frac{2}{11}$$

Hence, the required number is $\frac{2}{11}$.

Question 6.

$$\text{If } \frac{m+n}{m+3n} = \frac{2}{3}, \text{ find: } \frac{2n^2}{3m^2 + mn}.$$

Solution:

$$\frac{m+n}{m+3n} = \frac{2}{3}$$

$$\Rightarrow 3m + 3n = 2m + 6n$$

$$\Rightarrow m = 3n$$

$$\Rightarrow \frac{m}{n} = \frac{3}{1}$$

$$\frac{2n^2}{3m^2 + mn} = \frac{2}{3\left(\frac{m}{n}\right)^2 + \left(\frac{m}{n}\right)} \quad (\text{Dividing each term by } n^2)$$

$$= \frac{2}{3\left(\frac{3}{1}\right)^2 + \left(\frac{3}{1}\right)}$$

$$= \frac{2}{27+3} = \frac{1}{15}$$

Question 7.

Find $\frac{x}{y}$, when $x^2 + 6y^2 = 5xy$.

Solution:

$$x^2 + 6y^2 = 5xy$$

Dividing both sides by y^2 , we get,

$$\frac{x^2}{y^2} + \frac{6y^2}{y^2} = \frac{5xy}{y^2}$$

$$\left(\frac{x}{y}\right)^2 + 6 = 5\left(\frac{x}{y}\right)$$

$$\left(\frac{x}{y}\right)^2 - 5\left(\frac{x}{y}\right) + 6 = 0$$

$$\text{Let } \frac{x}{y} = a$$

$$\therefore a^2 - 5a + 6 = 0$$

$$\Rightarrow (a-2)(a-3) = 0$$

$$\Rightarrow a = 2, 3$$

$$\text{Hence, } \frac{x}{y} = 2, 3$$

Question 8.

If the ratio between 8 and 11 is the same as the ratio of $2x - y$ to $x + 2y$, find the value of $\frac{7x}{9y}$.

Solution:

$$\frac{2x - y}{x + 2y} = \frac{8}{11}$$

$$22x - 11y = 8x + 16y$$

$$14x = 27y$$

Given,

$$\frac{x}{y} = \frac{27}{14}$$

$$\therefore \frac{7x}{9y} = \frac{7 \times 27}{9 \times 14} = \frac{3}{2}$$

Question 9.

Divide Rs 1,290 into A, B and C such that A is $\frac{2}{5}$ of B and $B : C = 4 : 3$.

Solution:

$$\text{Given, } B : C = 4 : 3 \Rightarrow \frac{B}{C} = \frac{4}{3}$$

$$\text{And, } A = \frac{2}{5}B \Rightarrow \frac{A}{B} = \frac{2}{5}$$

$$\text{Now, } \frac{A}{B} = \frac{2}{5} = \frac{2 \times 4}{5 \times 4} = \frac{8}{20} \text{ and } \frac{B}{C} = \frac{4 \times 3}{3 \times 5} = \frac{20}{15}$$

$$\Rightarrow A : B : C = 8 : 20 : 15$$

$$\Rightarrow A = 8x, B = 20x \text{ and } C = 15x$$

$$\therefore 8x + 20x + 15x = 1290$$

$$\Rightarrow 43x = 1290$$

$$\Rightarrow x = 30$$

$$A's \text{ share} = 8x = 8 \times 30 = \text{Rs. } 240$$

$$B's \text{ share} = 20x = 20 \times 30 = \text{Rs. } 600$$

$$C's \text{ share} = 15x = 15 \times 30 = \text{Rs. } 450$$

Question 10.

A school has 630 students. The ratio of the number of boys to the number of girls is 3 : 2. This ratio changes to 7 : 5 after the admission of 90 new students. Find the number of newly admitted boys.

Solution:

Let the number of boys be $3x$.

Then, number of girls = $2x$

$$\therefore 3x + 2x = 630$$

$$\Rightarrow 5x = 630$$

$$\Rightarrow x = 126$$

$$\Rightarrow \text{Number of boys} = 3x = 3 \times 126 = 378$$

$$\text{And, Number of girls} = 2x = 2 \times 126 = 252$$

After admission of 90 new students, we have

$$\text{total number of students} = 630 + 90 = 720$$

Now, let the number of boys be $7x$.

Then, number of girls = $5x$

$$\therefore 7x + 5x = 720$$

$$\Rightarrow 12x = 720$$

$$\Rightarrow x = 60$$

$$\Rightarrow \text{Number of boys} = 7x = 7 \times 60 = 420$$

$$\text{And, Number of girls} = 5x = 5 \times 60 = 300$$

$$\therefore \text{Number of newly admitted boys} = 420 - 378 = 42$$

Question 11.

What quantity must be subtracted from each term of the ratio 9: 17 to make it equal to 1: 3?

Solution:

Let x be subtracted from each term of the ratio 9: 17.

$$\frac{9-x}{17-x} = \frac{1}{3}$$

$$27 - 3x = 17 - x$$

$$10 = 2x$$

$$x = 5$$

Thus, the required number which should be subtracted is 5.

Question 12.

The monthly pocket money of Ravi and Sanjeev are in the ratio 5 : 7. Their expenditures are in the ratio 3 : 5. If each saves Rs. 80 every month, find their monthly pocket money.

Solution:

Given that the pocket money of Ravi and Sanjeev are in the ratio 5 : 7

Thus, the pocket money of Ravi is 5k and that of Sanjeev is 7k.

Also given that the expenditure of Ravi and Sanjeev are in the ratio 3 : 5

Thus, the expenditure of Ravi is 3m and that of Sanjeev is 5m.

And each of them saves Rs. 80

$$\Rightarrow 5k - 3m = 80 \dots (1)$$

$$7k - 5m = 80 \dots (2)$$

Solving equations (1) and (2), we have,

$$k = 40, m = 40$$

Hence the monthly pocket money of Ravi is Rs. 200 and that of Sanjeev is Rs. 280.

Question 13.

The work done by $(x - 2)$ men in $(4x + 1)$ days and the work done by $(4x + 1)$ men in $(2x - 3)$ days are in the ratio 3: 8. Find the value of x.

Solution:

Assuming that all the men do the same amount of work in one day and one day work of each man = 1 units, we have,

$$\begin{aligned} &\text{Amount of work done by } (x - 2) \text{ men in } (4x + 1) \text{ days} \\ &= \text{Amount of work done by } (x - 2)(4x + 1) \text{ men in one day} \\ &= (x - 2)(4x + 1) \text{ units of work} \end{aligned}$$

Similarly,

Amount of work done by $(4x + 1)$ men in $(2x - 3)$ days
= $(4x + 1)(2x - 3)$ units of work

According to the given information,

$$\frac{(x - 2)(4x + 1)}{(4x + 1)(2x - 3)} = \frac{3}{8}$$

$$\frac{x - 2}{2x - 3} = \frac{3}{8}$$

$$8x - 16 = 6x - 9$$

$$2x = 7$$

$$x = \frac{7}{2} = 3.5$$

Question 14.

The bus fare between two cities is increased in the ratio 7: 9. Find the increase in the fare, if:

- (i) the original fare is Rs 245;
- (ii) the increased fare is Rs 207.

Solution:

According to the given information,

$$\text{Increased (new) bus fare} = \frac{9}{7} \times \text{original bus fare}$$

(i) We have:

$$\text{Increased (new) bus fare} = \frac{9}{7} \times \text{Rs } 245 = \text{Rs } 315$$

$$\therefore \text{Increase in fare} = \text{Rs } 315 - \text{Rs } 245 = \text{Rs } 70$$

(ii) We have:

$$\text{Rs } 207 = \frac{9}{7} \times \text{original bus fare}$$

$$\text{Original bus fare} = \text{Rs } 207 \times \frac{7}{9} = \text{Rs } 161$$

$$\therefore \text{Increase in fare} = \text{Rs } 207 - \text{Rs } 161 = \text{Rs } 46$$

Question 15.

By increasing the cost of entry ticket to a fair in the ratio 10: 13, the number of visitors to the fair has decreased in the ratio 6: 5. In what ratio has the total collection increased

or decreased?

Solution:

Let the cost of the entry ticket initially and at present be $10x$ and $13x$ respectively.

Let the number of visitors initially and at present be $6y$ and $5y$ respectively.

Initially, total collection = $10x \times 6y = 60xy$

At present, total collection = $13x \times 5y = 65xy$

Ratio of total collection = $60xy : 65xy = 12 : 13$

Thus, the total collection has increased in the ratio $12 : 13$.

Question 16.

In a basket, the ratio between the number of oranges and the number of apples is $7 : 13$.

If 8 oranges and 11 apples are eaten, the ratio between the number of oranges and the number of apples becomes $1 : 2$. Find the original number of oranges and the original number of apples in the basket.

Solution:

Let the original number of oranges and apples be $7x$ and $13x$.

According to the given information,

$$\frac{7x - 8}{13x - 11} = \frac{1}{2}$$
$$14x - 16 = 13x - 11$$
$$x = 5$$

Thus, the original number of oranges and apples are $7 \times 5 = 35$ and $13 \times 5 = 65$ respectively.

Question 17.

In a mixture of 126 kg of milk and water, milk and water are in ratio $5 : 2$. How much water must be added to the mixture to make this ratio $3 : 2$?

Solution:

Quantity of milk : Quantity of water = $5 : 2$

$$\therefore \text{Quantity of milk} = 126 \times \frac{5}{7} = 90 \text{ kg}$$

$$\Rightarrow \text{Quantity of water} = 126 - 90 = 36 \text{ kg}$$

New ratio = $3 : 2$

Let the quantity of water to be added be x kg.

$$\text{Then, milk : water} = \frac{90}{36+x}$$

$$\therefore \frac{90}{36+x} = \frac{3}{2}$$

$$\Rightarrow 180 = 108 + 3x$$

$$\Rightarrow 3x = 72$$

$$\Rightarrow x = 24$$

Thus, quantity of water to be added is 24 kg.

Question 18.

(A) If $A : B = 3 : 4$ and $B : C = 6 : 7$, find:

(i) $A : B : C$

(ii) $A : C$

(B) If $A : B = 2 : 5$ and $A : C = 3 : 4$, find

(i) $A : B : C$

Solution:

(A)

(i)

$$\frac{A}{B} = \frac{3}{4} = \frac{3}{4} \times \frac{3}{3} = \frac{9}{12}$$

$$\frac{B}{C} = \frac{6}{7} = \frac{6}{7} \times \frac{2}{2} = \frac{12}{14}$$

$$A : B : C = 9 : 12 : 14$$

(ii)

$$\frac{A}{B} = \frac{3}{4}$$

$$\frac{B}{C} = \frac{6}{7}$$

$$\therefore \frac{A}{C} = \frac{\frac{A}{B}}{\frac{B}{C}} = \frac{\frac{3}{4}}{\frac{6}{7}} = \frac{3}{4} \times \frac{7}{6} = \frac{7}{8}$$

$$\therefore A : C = 7 : 8$$

(B)

(i)

To compare 3 ratios, the consequent of the first ratio and the antecedent of the 2nd ratio must be made equal.

Given that $A:B = 2:5$ and $A:C = 3:4$

Interchanging the first ratio, we have,

$B:A = 5:2$ and $A:C = 3:4$

L.C.M. of 2 and 3 is 6.

$\Rightarrow B:A = 5 \times 3 : 2 \times 3$ and $A:C = 3 \times 2 : 4 \times 2$

$\Rightarrow B:A = 15:6$ and $A:C = 6:8$

$\Rightarrow B:A:C = 15:6:8$

$\Rightarrow A:B:C = 6:15:8$

Question 19(i).

If $3A = 4B = 6C$; find $A:B:C$.

Solution:

$$3A = 4B = 6C$$

$$3A = 4B \Rightarrow \frac{A}{B} = \frac{4}{3}$$

$$4B = 6C \Rightarrow \frac{B}{C} = \frac{6}{4} = \frac{3}{2}$$

Hence, $A:B:C = 4:3:2$

Question 19(ii).

If $2a = 3b$ and $4b = 5c$, find: $a:c$.

Solution:

We have,

$$2a = 3b \Rightarrow \frac{a}{b} = \frac{3}{2}$$

$$\text{And } 4b = 5c \Rightarrow \frac{b}{c} = \frac{5}{4}$$

$$\text{Now, } \frac{a}{b} = \frac{3}{2} = \frac{3 \times 5}{2 \times 5} = \frac{15}{10} \text{ and } \frac{b}{c} = \frac{5}{4} = \frac{5 \times 2}{4 \times 2} = \frac{10}{8}$$

$$\Rightarrow a:b:c = 15:10:8$$

$$\Rightarrow a:c = 15:8$$

Question 20.

Find the compound ratio of:

(i) $2:3, 9:14$ and $14:27$

(ii) $2a:3b, mn:x^2$ and $x:n$.

(iii) $\sqrt{2}:1, 3:\sqrt{5}$ and $\sqrt{20}:9$.

Solution:

(i) Required compound ratio = $3 \times 8:5 \times 15$

$$= \frac{3 \times 8}{5 \times 15}$$

$$= \frac{8}{25} = 8:25$$

(ii) Required compound ratio = $2 \times 9 \times 14:3 \times 14 \times 27$

$$= \frac{2 \times 9 \times 14}{3 \times 14 \times 27}$$

$$= \frac{2}{9} = 2:9$$

(iii) Required compound ratio = $2a \times mn \times x:3b \times x^2 \times n$

$$= \frac{2a \times mn \times x}{3b \times x^2 \times n}$$

$$= \frac{2am}{3bx} = 2am:3bx$$

(iv) Required compound ratio = $\sqrt{2} \times 3 \times \sqrt{20}:1 \times \sqrt{5} \times 9$

$$= \frac{\sqrt{2} \times 3 \times \sqrt{20}}{1 \times \sqrt{5} \times 9}$$

$$= \frac{\sqrt{2} \times \sqrt{4}}{3}$$

$$= \frac{2\sqrt{2}}{3} = 2\sqrt{2}:3$$

Question 21.

Find duplicate ratio of:

(i) $3:4$ (ii) $3\sqrt{3}:2\sqrt{5}$

Solution:

(i) Duplicate ratio of $3:4 = 3^2:4^2 = 9:16$

(ii) Duplicate ratio of $3\sqrt{3}:2\sqrt{5} = (3\sqrt{3})^2:(2\sqrt{5})^2 = 27:20$

Question 22.

Find the triplicate ratio of:

(i) 1: 3 (ii) $\frac{m}{2} : \frac{n}{3}$

Solution:

(i) Triplicate ratio of 1: 3 = $1^3 : 3^3 = 1 : 27$

(ii) Triplicate ratio of $\frac{m}{2} : \frac{n}{3}$

$$= \left(\frac{m}{2}\right)^3 : \left(\frac{n}{3}\right)^3 = \frac{m^3}{8} : \frac{n^3}{27} = \frac{\frac{m^3}{8}}{\frac{n^3}{27}} = 27m^3 : 8n^3$$

Question 23.

Find sub-duplicate ratio of:

(i) 9: 16 (ii) $(x - y)^4 : (x + y)^6$

Solution:

(i) Sub-duplicate ratio of 9: 16 = $\sqrt{9} : \sqrt{16} = 3 : 4$

(ii) Sub-duplicate ratio of $(x - y)^4 : (x + y)^6$

$$= \sqrt{(x - y)^4} : \sqrt{(x + y)^6} = (x - y)^2 : (x + y)^3$$

Question 24.

Find the sub-triplicate ratio of:

(i) 64: 27 (ii) $x^3 : 125y^3$

Solution:

(i) Sub-triplicate ratio of 64 : 27 = $\sqrt[3]{64} : \sqrt[3]{27} = 4 : 3$

(ii) Sub-triplicate ratio of $x^3 : 125y^3 = \sqrt[3]{x^3} : \sqrt[3]{125y^3} = x : 5y$

Question 25.

Find the reciprocal ratio of:

(i) 5: 8 (ii) $\frac{x}{3} : \frac{y}{7}$

Solution:

(i) Reciprocal ratio of $5:8 = \frac{1}{5} : \frac{1}{8} = 8:5$

(ii) Reciprocal ratio of $\frac{x}{3} : \frac{y}{7} = \frac{1}{\frac{x}{3}} : \frac{1}{\frac{y}{7}} = \frac{3}{x} : \frac{7}{y} = \frac{\frac{3}{x}}{\frac{7}{y}} = \frac{3y}{7x} = 3y:7x$

Question 26.

If $(x + 3) : (4x + 1)$ is the duplicate ratio of $3 : 5$, find the value of x .

Solution:

If $(x + 3) : (4x + 1)$ is the duplicate ratio of $3 : 5$,
find the value of x .

We have,

$$\frac{x+3}{4x+1} = \frac{3^2}{5^2}$$

$$\Rightarrow \frac{x+3}{4x+1} = \frac{9}{25}$$

$$\Rightarrow 25x + 75 = 36x + 9$$

$$\Rightarrow 11x = 66$$

$$\Rightarrow x = 6$$

Question 27.

If $m:n$ is the duplicate ratio of $m+x:n+x$; show that $x^2 = mn$.

Solution:

$$\frac{m}{n} = \frac{(m+x)^2}{(n+x)^2}$$

$$\frac{m}{n} = \frac{m^2 + x^2 + 2mx}{n^2 + x^2 + 2nx}$$

$$mn^2 + mx^2 + 2mnx = m^2n + nx^2 + 2mnx$$

$$x^2(m-n) = mn(m-n)$$

$$x^2 = mn$$

Question 28.

If $(3x - 9) : (5x + 4)$ is the triplicate ratio of $3 : 4$, find the value of x .

Solution:

We have,

$$\frac{3x - 9}{5x + 4} = \frac{3^3}{4^3}$$

$$\Rightarrow \frac{3x - 9}{5x + 4} = \frac{27}{64}$$

$$\Rightarrow \frac{3(x - 3)}{5x + 4} = \frac{27}{64}$$

$$\Rightarrow \frac{x - 3}{5x + 4} = \frac{9}{64}$$

$$\Rightarrow 64x - 192 = 45x + 36$$

$$\Rightarrow 19x = 228$$

$$\Rightarrow x = 12$$

Question 29.

Find the ratio compounded of the reciprocal ratio of $15 : 28$, the sub-duplicate ratio of $36 : 49$ and the triplicate ratio of $5 : 4$.

Solution:

Reciprocal ratio of $15 : 28 = 28 : 15$

Sub-duplicate ratio of $36 : 49 = \sqrt{36} : \sqrt{49} = 6 : 7$

Triplicate ratio of $5 : 4 = 5^3 : 4^3 = 125 : 64$

Required compounded ratio

$$= \frac{28 \times 6 \times 125}{15 \times 7 \times 64} = \frac{25}{8} = 25 : 8$$

Question 30(a).

If $r^2 = pq$, show that $p : q$ is the duplicate ratio of $(p + r) : (q + r)$.

Solution:

Given, $r^2 = pq$

$$\begin{aligned}
 \text{Duplicate ratio of } (p + r) : (q + r) &= (p + r)^2 : (q + r)^2 \\
 &= (p^2 + r^2 + 2pr) : (q^2 + r^2 + 2qr) \\
 &= (p^2 + pq + 2pr) : (q^2 + pq + 2qr) \\
 &= p(p + q + 2r) : q(q + p + 2r) \\
 &= p : q
 \end{aligned}$$

Thus, $p : q$ is the duplicate ratio of $(p + r) : (q + r)$.

Question 30(b).

If $(p - x) : (q - x)$ be the duplicate ratio of $p : q$

then show that: $\frac{1}{p} + \frac{1}{q} = \frac{1}{x}$

Solution:

We have,

$$\frac{(p - x)}{(q - x)} = \frac{p^2}{q^2}$$

$$\Rightarrow q^2(p - x) = p^2(q - x)$$

$$\Rightarrow pq^2 - q^2x = p^2q - p^2x$$

$$\Rightarrow p^2x - q^2x = p^2q - pq^2$$

$$\Rightarrow x(p^2 - q^2) = pq(p - q)$$

$$\Rightarrow x(p - q)(p + q) = pq(p - q)$$

$$\Rightarrow x = \frac{pq}{p + q}$$

$$\Rightarrow \frac{p + q}{pq} = \frac{1}{x}$$

$$\Rightarrow \frac{p}{pq} + \frac{q}{pq} = \frac{1}{x}$$

$$\Rightarrow \frac{1}{q} + \frac{1}{p} = \frac{1}{x}$$

$$\Rightarrow \frac{1}{p} + \frac{1}{q} = \frac{1}{x}$$

Exercise 7B

Question 1.

Find the fourth proportional to:

(i) 1.5, 4.5 and 3.5 (ii) $3a$, $6a^2$ and $2ab^2$

Solution:

(i) Let the fourth proportional to 1.5, 4.5 and 3.5 be x .

$$\Rightarrow 1.5 : 4.5 = 3.5 : x$$

$$\Rightarrow 1.5 \times x = 3.5 \times 4.5$$

$$\Rightarrow x = 10.5$$

(ii) Let the fourth proportional to $3a$, $6a^2$ and $2ab^2$ be x .

$$\Rightarrow 3a : 6a^2 = 2ab^2 : x$$

$$\Rightarrow 3a \times x = 2ab^2 \times 6a^2$$

$$\Rightarrow 3a \times x = 12a^3b^2$$

$$\Rightarrow x = 4a^2b^2$$

Question 2.

Find the third proportional to:

(i) $2\frac{2}{3}$ and 4 (ii) $a - b$ and $a^2 - b^2$

Solution:

(i) Let the third proportional to $2\frac{2}{3}$ and 4 be x .

$\Rightarrow 2\frac{2}{3}, 4, x$ are in continued proportion.

$$\Rightarrow 2\frac{2}{3} : 4 = 4 : x$$

$$\Rightarrow \frac{8}{3} = \frac{4}{x}$$

$$\Rightarrow x = 4 \times \frac{3}{8} = 1.5$$

(ii) Let the third proportional to $a - b$ and $a^2 - b^2$ be x .

$\Rightarrow a - b, a^2 - b^2, x$ are in continued proportion.

$$\Rightarrow a - b : a^2 - b^2 = a^2 - b^2 : x$$

$$\Rightarrow \frac{a - b}{a^2 - b^2} = \frac{a^2 - b^2}{x}$$

$$\Rightarrow x = \frac{(a^2 - b^2)^2}{a - b}$$

(i) Let the third proportional to $2\frac{2}{3}$ and 4 be x.

$\Rightarrow 2\frac{2}{3}, 4, x$ are in continued proportion.

$$\Rightarrow 2\frac{2}{3} : 4 = 4 : x$$

$$\Rightarrow \frac{8}{3} = \frac{4}{x}$$

$$\Rightarrow x = 16 \times \frac{3}{8} = 6$$

(ii) Let the third proportional to $a - b$ and $a^2 - b^2$ be x.

$\Rightarrow a - b, a^2 - b^2, x$ are in continued proportion.

$$\Rightarrow a - b : a^2 - b^2 = a^2 - b^2 : x$$

$$\Rightarrow \frac{a - b}{a^2 - b^2} = \frac{a^2 - b^2}{x}$$

$$\Rightarrow x = \frac{(a^2 - b^2)^2}{a - b}$$

$$\Rightarrow x = \frac{(a + b)(a - b)(a^2 - b^2)}{a - b}$$

$$\Rightarrow x = (a + b)(a^2 - b^2)$$

Question 3.

Find the mean proportional between:

(i) $6 + 3\sqrt{3}$ and $8 - 4\sqrt{3}$

(ii) $a - b$ and $a^3 - a^2b$

Solution:

(i) Let the mean proportional between $6 + 3\sqrt{3}$ and $8 - 4\sqrt{3}$ be x.

$\Rightarrow 6 + 3\sqrt{3}, x$ and $8 - 4\sqrt{3}$ are in continued proportion.

$$\Rightarrow 6 + 3\sqrt{3} : x = x : 8 - 4\sqrt{3}$$

$$\Rightarrow x \times x = (6 + 3\sqrt{3})(8 - 4\sqrt{3})$$

$$\Rightarrow x^2 = 48 + 24\sqrt{3} - 24\sqrt{3} - 36$$

$$\Rightarrow x^2 = 12$$

$$\Rightarrow x = 2\sqrt{3}$$

(ii) Let the mean proportional between $a - b$ and $a^3 - a^2b$ be x.

$\Rightarrow a - b, x, a^3 - a^2b$ are in continued proportion.

$$\Rightarrow a - b : x = x : a^3 - a^2b$$

$$\Rightarrow x \times x = (a - b)(a^3 - a^2b)$$

$$\Rightarrow x^2 = (a - b) a^2(a - b) = [a(a - b)]^2$$

$$\Rightarrow x = a(a - b)$$

Question 4.

If $x + 5$ is the mean proportional between $x + 2$ and $x + 9$; find the value of x .

Solution:

Given, $x + 5$ is the mean proportional between $x + 2$ and $x + 9$.

$\Rightarrow (x + 2), (x + 5)$ and $(x + 9)$ are in continued proportion.

$$\Rightarrow (x + 2) : (x + 5) = (x + 5) : (x + 9)$$

$$\Rightarrow (x + 5)^2 = (x + 2)(x + 9)$$

$$\Rightarrow x^2 + 25 + 10x = x^2 + 2x + 9x + 18$$

$$\Rightarrow 25 - 18 = 11x - 10x$$

$$\Rightarrow x = 7$$

Question 5.

If $x^2, 4$ and 9 are in continued proportion, find x .

Solution:

Given, $x^2, 4$ and 9 are in continued proportion.

$$\therefore \frac{x^2}{4} = \frac{4}{9}$$

$$\Rightarrow 9x^2 = 16$$

$$\Rightarrow x^2 = \frac{16}{9}$$

$$\Rightarrow x = \frac{4}{3}$$

Question 6.

What least number must be added to each of the numbers $6, 15, 20$ and 43 to make them proportional?

Solution:

Let the number added be x .

$$\therefore (6 + x) : (15 + x) :: (20 + x) : (43 + x)$$

$$\frac{6 + x}{15 + x} = \frac{20 + x}{43 + x}$$

$$(6 + x)(43 + x) = (20 + x)(15 + x)$$

$$258 + 6x + 43x + x^2 = 300 + 20x + 15x + x^2$$

$$49x - 35x = 300 - 258$$

$$14x = 42$$

$$x = 3$$

Thus, the required number which should be added is 3.

Question 7(i).

If a, b, c are in continued proportion,

show that: $\frac{a^2 + b^2}{b(a + c)} = \frac{b(a + c)}{b^2 + c^2}$.

Solution:

Since a, b, c are in continued proportion,

$$\frac{a}{b} = \frac{b}{c}$$

$$\Rightarrow b^2 = ac$$

$$\begin{aligned} \text{Now, } (a^2 + b^2)(b^2 + c^2) &= (a^2 + ac)(ac + c^2) \\ &= a(a + c)c(a + c) \\ &= ac(a + c)^2 \\ &= b^2(a + c)^2 \end{aligned}$$

$$\Rightarrow (a^2 + b^2)(b^2 + c^2) = [b(a + c)][b(a + c)]$$

$$\Rightarrow \frac{a^2 + b^2}{b(a + c)} = \frac{b(a + c)}{b^2 + c^2}$$

Question 7(ii).

If a, b, c are in continued proportion and $a(b - c) = 2b$,

prove that: $a - c = \frac{2(a + b)}{a}$.

Solution:

Since a, b, c are in continued proportion,

$$\frac{a}{b} = \frac{b}{c}$$

$$\Rightarrow b^2 = ac$$

$$a(b - c) = 2b$$

$$\Rightarrow ab - ac = 2b$$

$$\Rightarrow ab - b^2 = 2b$$

$$\Rightarrow b(a - b) = 2b$$

$$\Rightarrow a - b = 2$$

Now,

$$\text{L.H.S.} = a - c$$

$$= \frac{a(a - c)}{a}$$

$$= \frac{a^2 - ac}{a}$$

$$= \frac{a^2 - b^2}{a}$$

$$= \frac{(a - b)(a + b)}{a}$$

$$= \frac{2(a + b)}{2}$$

$$= \text{R.H.S.}$$

Question 7(iii).

If $\frac{a}{b} = \frac{c}{d}$, show that: $\frac{a^3c + ac^3}{b^3d + bd^3} = \frac{(a + c)^4}{(b + d)^4}$.

Solution:

$$\text{Let } \frac{a}{b} = \frac{c}{d} = k$$

$$\Rightarrow a = bk \text{ and } c = dk$$

$$\text{L.H.S.} = \frac{a^3c + ac^3}{b^3d + bd^3}$$

$$\begin{aligned}
&= \frac{ac(a^2 + c^2)}{bd(b^2 + d^2)} \\
&= \frac{(bk \times dk)(b^2k^2 + d^2k^2)}{bd(b^2 + d^2)} \\
&= \frac{k^2 \times k^2(b^2 + d^2)}{(b^2 + d^2)} \\
&= k^4
\end{aligned}$$

$$\text{R.H.S.} = \frac{(a+c)^4}{(b+d)^4} = \frac{(bk+dk)^4}{(b+d)^4} = \left[\frac{k(b+d)}{b+d} \right]^4 = k^4$$

$$\text{Hence, } \frac{a^3c + ac^3}{b^3d + bd^3} = \frac{(a+c)^4}{(b+d)^4}$$

Question 8.

What least number must be subtracted from each of the numbers 7, 17 and 47 so that the remainders are in continued proportion?

Solution:

Let the number subtracted be x .

$$\therefore (7-x) : (17-x) :: (17-x) : (47-x)$$

$$\frac{7-x}{17-x} = \frac{17-x}{47-x}$$

$$(7-x)(47-x) = (17-x)^2$$

$$329 - 47x - 7x + x^2 = 289 - 34x + x^2$$

$$329 - 289 = -34x + 54x$$

$$20x = 40$$

$$x = 2$$

Thus, the required number which should be subtracted is 2.

Question 9.

If y is the mean proportional between x and z ; show that $xy + yz$ is the mean proportional between $x^2 + y^2$ and $y^2 + z^2$.

Solution:

Since y is the mean proportion between x and z

Therefore, $y^2 = xz$

Now, we have to prove that $xy + yz$ is the mean proportional between $x^2 + y^2$ and $y^2 + z^2$, i.e.,

$$(xy + yz)^2 = (x^2 + y^2)(y^2 + z^2)$$

$$\text{LHS} = (xy + yz)^2$$

$$= [y(x + z)]^2$$

$$= y^2(x + z)^2$$

$$= xz(x + z)^2$$

$$\text{RHS} = (x^2 + y^2)(y^2 + z^2)$$

$$= (x^2 + xz)(xz + z^2)$$

$$= x(x + z)z(x + z)$$

$$= xz(x + z)^2$$

$$\text{LHS} = \text{RHS}$$

Hence, proved.

Question 10.

If q is the mean proportional between p and r , show that:
 $pqr(p + q + r)^3 = (pq + qr + rp)^3$.

Solution:

Given, q is the mean proportional between p and r .

$$\Rightarrow q^2 = pr$$

$$\text{L.H.S.} = pqr(p + q + r)^3$$

$$= qq^2(p + q + r)^3 \quad [\because q^2 = pr]$$

$$= q^3(p + q + r)^3$$

$$= [q(p + q + r)]^3$$

$$= (pq + q^2 + qr)^3$$

$$= (pq + pr + qr)^3 \quad [\because q^2 = pr]$$

$$= \text{R.H.S.}$$

Question 11.

If three quantities are in continued proportion; show that the ratio of the first to the third is the duplicate ratio of the first to the second.

Solution:

Let x , y and z be the three quantities which are in continued proportion.

Then, $x : y :: y : z \Rightarrow y^2 = xz \dots (1)$

Now, we have to prove that

$$x : z = x^2 : y^2$$

That is we need to prove that

$$xy^2 = x^2z$$

$$\text{LHS} = xy^2 = x(xz) = x^2z = \text{RHS [Using (1)]}$$

Hence, proved.

Question 12.

If y is the mean proportional between x and z , prove that:

$$\frac{x^2 - y^2 + z^2}{x^{-2} - y^{-2} + z^{-2}} = y^4.$$

Solution:

Given, y is the mean proportional between x and z .

$$\Rightarrow y^2 = xz$$

$$\begin{aligned} \text{LHS} &= \frac{x^2 - y^2 + z^2}{x^{-2} - y^{-2} + z^{-2}} \\ &= \frac{x^2 - y^2 + z^2}{\frac{1}{x^2} - \frac{1}{y^2} + \frac{1}{z^2}} \\ &= \frac{x^2 - xz + z^2}{\frac{1}{x^2} - \frac{1}{xz} + \frac{1}{z^2}} \quad (\because y^2 = xz) \\ &= \frac{x^2 - xz + z^2}{\frac{z^2 - xz + x^2}{x^2z^2}} \\ &= x^2z^2 \\ &= (xz)^2 \\ &= (y^2)^2 \quad (\because y^2 = xz) \\ &= y^4 \\ &= \text{RHS} \end{aligned}$$

Question 13.

Given four quantities a, b, c and d are in proportion. Show that:

$$(a - c)b^2 : (b - d)cd = (a^2 - b^2 - ab) : (c^2 - d^2 - cd)$$

Solution:

$$\text{Let } \frac{a}{b} = \frac{c}{d} = k$$

$$\Rightarrow a = bk \text{ and } c = dk$$

$$\begin{aligned} \text{LHS} &= \frac{(a - c)b^2}{(b - d)cd} \\ &= \frac{(bk - dk)b^2}{(b - d)dkd} \\ &= \frac{k(b - d)b^2}{(b - d)d^2k} \\ &= \frac{b^2}{d^2} \end{aligned}$$

$$\begin{aligned} \text{RHS} &= \frac{(a^2 - b^2 - ab)}{(c^2 - d^2 - cd)} \\ &= \frac{(b^2k^2 - b^2 - bkb)}{(d^2k^2 - d^2 - dkd)} \\ &= \frac{b^2(k^2 - 1 - k)}{d^2(k^2 - 1 - k)} \\ &= \frac{b^2}{d^2} \end{aligned}$$

$$\Rightarrow \text{LHS} = \text{RHS}$$

Hence proved.

Question 14.

Find two numbers such that the mean mean proportional between them is 12 and the third proportional to them is 96.

Solution:

Let a and b be the two numbers, whose mean proportional is 12.

$$\therefore ab = 12^2 \Rightarrow ab = 144 \Rightarrow b = \frac{144}{a} \dots\dots\dots (1)$$

Now, third proportional is 96

$$\therefore a : b :: b : 96$$

$$\Rightarrow b^2 = 96a$$

$$\Rightarrow \left(\frac{144}{a}\right)^2 = 96a$$

$$\Rightarrow \frac{(144)^2}{a^2} = 96a$$

$$\Rightarrow a^3 = \frac{144 \times 144}{96}$$

$$\Rightarrow a^3 = 216$$

$$\Rightarrow a = 6$$

$$b = \frac{144}{6} = 24$$

Therefore, the numbers are 6 and 24.

Question 15.

Find the third proportional to $\frac{x}{y} + \frac{y}{x}$ and $\sqrt{x^2 + y^2}$

Solution:

Let the required third proportional be p.

$$\Rightarrow \frac{x}{y} + \frac{y}{x}, \sqrt{x^2 + y^2}, p \text{ are in continued proportion.}$$

$$\Rightarrow \frac{x}{y} + \frac{y}{x} : \sqrt{x^2 + y^2} = \sqrt{x^2 + y^2} : p$$

$$\Rightarrow p \left(\frac{x}{y} + \frac{y}{x} \right) = \left(\sqrt{x^2 + y^2} \right)^2$$

$$\Rightarrow p \left(\frac{x^2 + y^2}{xy} \right) = x^2 + y^2$$

$$\Rightarrow p = xy$$

Question 16.

If $p : q = r : s$, then show that:
 $mp + nq : q = mr + ns : s$.

Solution:

$$\frac{p}{q} = \frac{r}{s}$$

$$\Rightarrow \frac{mp}{q} = \frac{mr}{s}$$

$$\Rightarrow \frac{mp}{q} + n = \frac{mr}{s} + n$$

$$\Rightarrow \frac{mp + nq}{q} = \frac{mr + ns}{s}$$

Hence, $mp + nq : q = mr + ns : s$.

Question 17.

If $p + r = mq$ and $\frac{1}{q} + \frac{1}{s} = \frac{m}{r}$; then prove that $p : q = r : s$.

Solution:

$$\frac{1}{q} + \frac{1}{s} = \frac{m}{r}$$

$$\frac{s + q}{qs} = \frac{m}{r}$$

$$\frac{s + q}{s} = \frac{mq}{r}$$

$$\frac{s + q}{s} = \frac{p + r}{r} \quad (\because p + r = mq)$$

$$1 + \frac{q}{s} = \frac{p}{r} + 1$$

$$\frac{q}{s} = \frac{p}{r}$$

$$\frac{p}{q} = \frac{r}{s}$$

Hence, proved.

Question 18.

If $\frac{a}{b} = \frac{c}{d}$, prove that each of the given ratio is equal to:

(i) $\frac{5a + 4c}{5b + 4d}$

(ii) $\frac{13a - 8c}{13b - 8d}$

(iii) $\sqrt{\frac{3a^2 - 10c^2}{3b^2 - 10d^2}}$

(iv) $\left(\frac{8a^3 + 15c^3}{8b^3 + 15d^3}\right)^{\frac{1}{3}}$

Solution:

Let $\frac{a}{b} = \frac{c}{d} = k$

Then, $a = bk$ and $c = dk$

(i) $\frac{5a + 4c}{5b + 4d} = \frac{5(bk) + 4(dk)}{5b + 4d} = \frac{k(5b + 4d)}{5b + 4d} = k = \text{each given ratio}$

(ii) $\frac{13a - 8c}{13b - 8d} = \frac{13(bk) - 8(dk)}{13b - 8d} = \frac{k(13b - 8d)}{13b - 8d} = k = \text{each given ratio}$

(iii) $\sqrt{\frac{3a^2 - 10c^2}{3b^2 - 10d^2}} = \sqrt{\frac{3(bk)^2 - 10(dk)^2}{3b^2 - 10d^2}} = \sqrt{\frac{k^2(3b^2 - 10d^2)}{3b^2 - 10d^2}} = k$
= each given ratio

(iv) $\left(\frac{8a^3 + 15c^3}{8b^3 + 15d^3}\right)^{\frac{1}{3}} = \left[\frac{8(bk)^3 + 15(dk)^3}{8b^3 + 15d^3}\right]^{\frac{1}{3}} = \left[\frac{k^3(8b^3 + 15d^3)}{8b^3 + 15d^3}\right]^{\frac{1}{3}} = k$
= each given ratio

Question 19.

If a, b, c and d are in proportion, prove that:

$$(i) \frac{13a + 17b}{13c + 17d} = \sqrt{\frac{2ma^2 - 3nb^2}{2mc^2 - 3nd^2}}$$

$$(ii) \sqrt{\frac{4a^2 + 9b^2}{4c^2 + 9d^2}} = \left(\frac{xa^3 - 5yb^3}{xc^3 - 5yd^3} \right)^{\frac{1}{3}}$$

Solution:

a, b, c and d are in proportion

$$\frac{a}{b} = \frac{c}{d} = k \text{ (say)}$$

Then, a = bk and c = dk

$$(i) \text{L.H.S.} = \frac{13a + 17b}{13c + 17d} = \frac{13(bk) + 17b}{13(dk) + 17d} = \frac{b(13k + 17)}{d(13k + 17)} = \frac{b}{d}$$

$$\text{R.H.S.} = \sqrt{\frac{2ma^2 - 3nb^2}{2mc^2 - 3nd^2}} = \sqrt{\frac{2m(bk)^2 - 3nb^2}{2m(dk)^2 - 3nd^2}} = \sqrt{\frac{b^2(2mk^2 - 3n)}{d^2(2mk^2 - 3n)}} = \frac{b}{d}$$

Hence, L.H.S. = R.H.S.

$$(ii) \text{L.H.S.} = \sqrt{\frac{4a^2 + 9b^2}{4c^2 + 9d^2}} = \sqrt{\frac{4(bk)^2 + 9b^2}{4(dk)^2 + 9d^2}} = \sqrt{\frac{b^2(4k^2 + 9)}{d^2(4k^2 + 9)}} = \frac{b}{d}$$

$$\begin{aligned} \text{R.H.S.} &= \left(\frac{xa^3 - 5yb^3}{xc^3 - 5yd^3} \right)^{\frac{1}{3}} = \left[\frac{x(bk)^3 - 5yb^3}{x(dk)^3 - 5yd^3} \right]^{\frac{1}{3}} \\ &= \left[\frac{b^3(xk^3 - 5y)}{d^3(xk^3 - 5y)} \right]^{\frac{1}{3}} \\ &= \left[\frac{b^3}{d^3} \right]^{\frac{1}{3}} = \frac{b}{d} \end{aligned}$$

Hence, L.H.S. = R.H.S.

Question 20.

If $\frac{x}{a} = \frac{y}{b} = \frac{z}{c}$, prove that:

$$\frac{2x^3 - 3y^3 + 4z^3}{2a^3 - 3b^3 + 4c^3} = \left(\frac{2x - 3y + 4z}{2a - 3b + 4c} \right)^3$$

Solution:

$$\text{Let } \frac{x}{a} = \frac{y}{b} = \frac{z}{c} = k$$

Then, $x = ak$, $y = bk$ and $z = ck$

$$\begin{aligned} \text{L.H.S.} &= \frac{2x^3 - 3y^3 + 4z^3}{2a^3 - 3b^3 + 4c^3} \\ &= \frac{2(ak)^3 - 3(bk)^3 + 4(ck)^3}{2a^3 - 3b^3 + 4c^3} \\ &= \frac{2a^3k^3 - 3b^3k^3 + 4c^3k^3}{2a^3 - 3b^3 + 4c^3} \\ &= \frac{k^3(2a^3 - 3b^3 + 4c^3)}{2a^3 - 3b^3 + 4c^3} \\ &= k^3 \end{aligned}$$

$$\begin{aligned} \text{R.H.S.} &= \left(\frac{2x - 3y + 4z}{2a - 3b + 4c} \right)^3 \\ &= \left(\frac{2ak - 3bk + 4ck}{2a - 3b + 4c} \right)^3 \\ &= \left[\frac{k(2a - 3b + 4c)}{2a - 3b + 4c} \right]^3 \\ &= k^3 \end{aligned}$$

Hence, L.H.S. = R.H.S.

Exercise 7C**Question 1.**

If $a : b = c : d$, prove that:

- (i) $5a + 7b : 5a - 7b = 5c + 7d : 5c - 7d$.
- (ii) $(9a + 13b)(9c - 13d) = (9c + 13d)(9a - 13b)$.

(iii) $xa + yb : xc + yd = b : d$.

Solution:

(i) Given, $\frac{a}{b} = \frac{c}{d}$

$$\Rightarrow \frac{5a}{7b} = \frac{5c}{7d} \quad \left(\text{Multiplying each side by } \frac{5}{7} \right)$$

$$\Rightarrow \frac{5a+7b}{5a-7b} = \frac{5c+7d}{5c-7d} \quad (\text{By componendo and dividendo})$$

(ii) Given, $\frac{a}{b} = \frac{c}{d}$

$$\Rightarrow \frac{9a}{13b} = \frac{9c}{13d} \quad \left(\text{Multiplying each side by } \frac{9}{13} \right)$$

$$\Rightarrow \frac{9a+13b}{13a-13b} = \frac{9c+13d}{9c-13d} \quad (\text{By componendo and dividendo})$$

$$\Rightarrow (9a+13b)(9c-13d) = (9c+13d)(9a-13b)$$

(iii) Given, $\frac{a}{b} = \frac{c}{d}$

$$\Rightarrow \frac{xa}{yb} = \frac{xc}{yd} \quad \left(\text{Multiplying each side by } \frac{x}{y} \right)$$

$$\Rightarrow \frac{xa+yb}{yb} = \frac{xc+yd}{yd} \quad (\text{By componendo})$$

$$\Rightarrow \frac{xa+yb}{xc+yd} = \frac{yb}{yd}$$

$$\Rightarrow \frac{xa+yb}{xc+yd} = \frac{b}{d}$$

Question 2.

If $a : b = c : d$, prove that:

$$(6a + 7b) (3c - 4d) = (6c + 7d) (3a - 4b).$$

Solution:

Given, $\frac{a}{b} = \frac{c}{d}$

$$\Rightarrow \frac{6a}{7b} = \frac{6c}{7d} \quad \left(\text{Multiplying each side by } \frac{6}{7} \right)$$

$$\Rightarrow \frac{6a+7b}{7b} = \frac{6c+7d}{7d} \quad (\text{By componendo})$$

$$\Rightarrow \frac{6a+7b}{6c+7d} = \frac{7b}{7d} = \frac{b}{d} \quad \dots(1)$$

Also, $\frac{a}{b} = \frac{c}{d}$

$$\Rightarrow \frac{3a}{4b} = \frac{3c}{4d} \quad \left(\text{Multiplying each side by } \frac{3}{4} \right)$$

$$\Rightarrow \frac{3a-4b}{4b} = \frac{3c-4d}{4d} \quad (\text{By dividendo})$$

$$\Rightarrow \frac{3a-4b}{3c-4d} = \frac{4b}{4d} = \frac{b}{d} \quad \dots(2)$$

From (1) and (2),

$$\frac{6a+7b}{6c+7d} = \frac{3a-4b}{3c-4d}$$

$$(6a+7b)(3c-4d) = (6c+7d)(3a-4b)$$

Question 3.

Given, $\frac{a}{b} = \frac{c}{d}$, prove that:

$$\frac{3a-5b}{3a+5b} = \frac{3c-5d}{3c+5d}$$

Solution:

$$\frac{a}{b} = \frac{c}{d}$$

$$\frac{3a}{5b} = \frac{3c}{5d} \quad \left(\text{Multiplying each side by } \frac{3}{5} \right)$$

$$\frac{3a+5b}{3a-5b} = \frac{3c+5d}{3c-5d} \quad (\text{By componendo and dividendo})$$

$$\frac{3a-5b}{3a+5b} = \frac{3c-5d}{3c+5d} \quad (\text{By alternendo})$$

Question 4.

If $\frac{5x + 6y}{5u + 6v} = \frac{5x - 6y}{5u - 6v}$; then prove that:

$x : y = u : v$.

Solution:

$$\frac{5x + 6y}{5u + 6v} = \frac{5x - 6y}{5u - 6v}$$

(By alternendo)

$$\frac{5x + 6y}{5x - 6y} = \frac{5u + 6v}{5u - 6v}$$

$$\frac{5x + 6y + 5x - 6y}{5x + 6y - 5x + 6y} = \frac{5u + 6v + 5u - 6v}{5u + 6v - 5u + 6v}$$

(By componendo and dividendo)

$$\frac{10x}{12y} = \frac{10u}{12v}$$

$$\frac{x}{y} = \frac{u}{v}$$

Question 5.

If $(7a + 8b)(7c - 8d) = (7a - 8b)(7c + 8d)$, prove that $a : b = c : d$.

Solution:

Given, $\frac{7a + 8b}{7a - 8b} = \frac{7c + 8d}{7c - 8d}$

Applying componendo and dividendo,

$$\frac{7a + 8b + 7a - 8b}{7a + 8b - 7a + 8b} = \frac{7c + 8d + 7c - 8d}{7c + 8d - 7c + 8d}$$

$$\Rightarrow \frac{14a}{16b} = \frac{14c}{16d}$$

$$\Rightarrow \frac{a}{b} = \frac{c}{d}$$

Hence, $a : b = c : d$.

Question 6.

(i) If $x = \frac{6ab}{a+b}$, find the value of:

$$\frac{x+3a}{x-3a} + \frac{x+3b}{x-3b}.$$

(ii) If $a = \frac{4\sqrt{6}}{\sqrt{2}+\sqrt{3}}$, find the value of:

$$\frac{a+2\sqrt{2}}{a-2\sqrt{2}} + \frac{a+2\sqrt{3}}{a-2\sqrt{3}}.$$

Solution:

$$(i) x = \frac{6ab}{a+b}$$

$$\Rightarrow \frac{x}{3a} = \frac{2b}{a+b}$$

Applying componendo and dividendo,

$$\frac{x+3a}{x-3a} = \frac{2b+a+b}{2b-a-b}$$

$$\frac{x+3a}{x-3a} = \frac{3b+a}{b-a} \quad \dots (1)$$

$$\text{Again, } x = \frac{6ab}{a+b}$$

$$\Rightarrow \frac{x}{3b} = \frac{2a}{a+b}$$

Applying componendo and dividendo,

$$\frac{x+3b}{x-3b} = \frac{2a+a+b}{2a-a-b}$$

$$\frac{x+3b}{x-3b} = \frac{3a+b}{a-b} \quad \dots (2)$$

From (1) and (2),

$$\frac{x+3a}{x-3a} + \frac{x+3b}{x-3b} = \frac{3b+a}{b-a} + \frac{3a+b}{a-b}$$

$$\frac{x+3a}{x-3a} + \frac{x+3b}{x-3b} = \frac{-3b-a+3a+b}{a-b}$$

$$\frac{x+3a}{x-3a} + \frac{x+3b}{x-3b} = \frac{2a-2b}{a-b} = 2$$

$$(ii) a = \frac{4\sqrt{6}}{\sqrt{2} + \sqrt{3}}$$

$$\frac{a}{2\sqrt{2}} = \frac{2\sqrt{3}}{\sqrt{2} + \sqrt{3}}$$

Applying componendo and dividendo,

$$\frac{a + 2\sqrt{2}}{a - 2\sqrt{2}} = \frac{2\sqrt{3} + \sqrt{2} + \sqrt{3}}{2\sqrt{3} - \sqrt{2} - \sqrt{3}}$$

$$\frac{a + 2\sqrt{2}}{a - 2\sqrt{2}} = \frac{3\sqrt{3} + \sqrt{2}}{\sqrt{3} - \sqrt{2}} \quad \dots (1)$$

$$\frac{a}{2\sqrt{3}} = \frac{2\sqrt{2}}{\sqrt{2} + \sqrt{3}}$$

Applying componendo and dividendo,

$$\frac{a + 2\sqrt{3}}{a - 2\sqrt{3}} = \frac{2\sqrt{2} + \sqrt{2} + \sqrt{3}}{2\sqrt{2} - \sqrt{2} - \sqrt{3}}$$

$$\frac{a + 2\sqrt{3}}{a - 2\sqrt{3}} = \frac{3\sqrt{2} + \sqrt{3}}{\sqrt{2} - \sqrt{3}} \quad \dots (2)$$

From (1) and (2),

$$\frac{a + 2\sqrt{2}}{a - 2\sqrt{2}} + \frac{a + 2\sqrt{3}}{a - 2\sqrt{3}} = \frac{3\sqrt{3} + \sqrt{2}}{\sqrt{3} - \sqrt{2}} + \frac{3\sqrt{2} + \sqrt{3}}{\sqrt{2} - \sqrt{3}}$$

$$\frac{a + 2\sqrt{2}}{a - 2\sqrt{2}} + \frac{a + 2\sqrt{3}}{a - 2\sqrt{3}} = \frac{3\sqrt{2} + \sqrt{3} - 3\sqrt{3} - \sqrt{2}}{\sqrt{2} - \sqrt{3}}$$

$$\frac{a + 2\sqrt{2}}{a - 2\sqrt{2}} + \frac{a + 2\sqrt{3}}{a - 2\sqrt{3}} = \frac{2\sqrt{2} - 2\sqrt{3}}{\sqrt{2} - \sqrt{3}} = 2$$

Question 7.

If $(a + b + c + d)(a - b - c + d) = (a + b - c - d)(a - b + c - d)$, prove that $a : b = c : d$.

Solution:

Given, $\frac{a+b+c+d}{a+b-c-d} = \frac{a-b+c-d}{a-b-c+d}$

Applying componendo and dividendo,

$$\frac{(a+b+c+d)+(a+b-c-d)}{(a+b+c+d)-(a+b-c-d)} = \frac{(a-b+c-d)+(a-b-c+d)}{(a-b+c-d)-(a-b-c+d)}$$

$$\frac{2(a+b)}{2(c+d)} = \frac{2(a-b)}{2(c-d)}$$

$$\frac{a+b}{c+d} = \frac{a-b}{c-d}$$

$$\frac{a+b}{a-b} = \frac{c+d}{c-d}$$

Applying componendo and dividendo,

$$\frac{a+b+a-b}{a+b-a+b} = \frac{c+d+c-d}{c+d-c+d}$$

$$\frac{2a}{2b} = \frac{2c}{2d}$$

$$\frac{a}{b} = \frac{c}{d}$$

Question 8.

If $\frac{a-2b-3c+4d}{a+2b-3c-4d} = \frac{a-2b+3c-4d}{a+2b+3c+4d}$, show that $2ad = 3bc$.

Solution:

$$\frac{a-2b-3c+4d}{a+2b-3c-4d} = \frac{a-2b+3c-4d}{a+2b+3c+4d}$$

Applying componendo and dividendo,

$$\begin{aligned} \frac{(a-2b-3c+4d)+(a+2b-3c-4d)}{(a-2b-3c+4d)-(a+2b-3c-4d)} &= \frac{(a-2b+3c-4d)+(a+2b+3c+4d)}{(a-2b+3c-4d)-(a+2b+3c+4d)} \\ &= \frac{2(a-3c)}{2(-2b+4d)} = \frac{2(a+3c)}{2(-2b-4d)} \end{aligned}$$

$$\frac{a-3c}{-2b+4d} = \frac{a+3c}{-2b-4d}$$

$$\frac{a - 2b - 3c + 4d}{a + 2b - 3c - 4d} = \frac{a - 2b + 3c - 4d}{a + 2b + 3c + 4d}$$

Applying componendo and dividendo,

$$\begin{aligned} \frac{(a - 2b - 3c + 4d) + (a + 2b - 3c - 4d)}{(a - 2b - 3c + 4d) - (a + 2b - 3c - 4d)} \\ = \frac{(a - 2b + 3c - 4d) + (a + 2b + 3c + 4d)}{(a - 2b + 3c - 4d) - (a + 2b + 3c + 4d)} \end{aligned}$$

$$\frac{2(a - 3c)}{2(-2b + 4d)} = \frac{2(a + 3c)}{2(-2b - 4d)}$$

$$\frac{a - 3c}{a + 3c} = \frac{-2b + 4d}{-2b - 4d}$$

Applying componendo and dividendo,

$$\frac{a - 3c + a + 3c}{a - 3c - a - 3c} = \frac{-2b + 4d - 2b - 4d}{-2b + 4d + 2b + 4d}$$

$$\frac{2a}{-6c} = \frac{-4b}{8d}$$

$$\frac{a}{-3c} = \frac{-b}{2d}$$

$$2ad = 3bc$$

Question 9.

If $(a^2 + b^2)(x^2 + y^2) = (ax + by)^2$; prove that: $\frac{a}{x} = \frac{b}{y}$.

Solution:

$$\text{Given, } (a^2 + b^2)(x^2 + y^2) = (ax + by)^2$$

$$a^2x^2 + a^2y^2 + b^2x^2 + b^2y^2 = a^2x^2 + b^2y^2 + 2abxy$$

$$a^2y^2 + b^2x^2 - 2abxy = 0$$

$$(ay - bx)^2 = 0$$

$$ay - bx = 0$$

$$ay = bx$$

$$\frac{a}{x} = \frac{b}{y}$$

Question 10.

If a, b and c are in continued proportion, prove that:

$$(i) \frac{a^2 + ab + b^2}{b^2 + bc + c^2} = \frac{a}{c}$$

$$(ii) \frac{a^2 + b^2 + c^2}{(a+b+c)^2} = \frac{a-b+c}{a+b+c}$$

Solution:

Given, a, b and c are in continued proportion.

$$\Rightarrow \frac{a}{b} = \frac{b}{c} = k \text{ (say)}$$

$$\Rightarrow a = bk, b = ck$$

$$\Rightarrow a = (ck)k = dk^2, b = dk$$

$$\begin{aligned} (i) \text{ L.H.S.} &= \frac{a^2 + ab + b^2}{b^2 + bc + c^2} \\ &= \frac{(dk^2)^2 + (dk^2)(dk) + (dk)^2}{(dk)^2 + (dk)c + c^2} \\ &= \frac{c^2k^4 + c^2k^3 + c^2k^2}{c^2k^2 + c^2k + c^2} \\ &= \frac{c^2k^2(k^2 + k + 1)}{c^2(k^2 + k + 1)} \\ &= k^2 \end{aligned}$$

$$\text{R.H.S.} = \frac{a}{c} = \frac{dk^2}{c} = k^2$$

$$\therefore \text{L.H.S.} = \text{R.H.S.}$$

$$\begin{aligned}
 \text{(ii) L.H.S.} &= \frac{a^2 + b^2 + c^2}{(a + b + c)^2} \\
 &= \frac{(dk^2)^2 + (dk)^2 + c^2}{(dk^2 + dk + c)^2} \\
 &= \frac{c^2k^4 + c^2k^2 + c^2}{c^2(k^2 + k + 1)^2} \\
 &= \frac{c^2(k^4 + k^2 + 1)}{c^2(k^2 + k + 1)^2} \\
 &= \frac{k^4 + k^2 + 1}{(k^2 + k + 1)^2}
 \end{aligned}$$

$$\begin{aligned}
 \text{R.H.S.} &= \frac{a - b + c}{a + b + c} \\
 &= \frac{dk^2 - dk + c}{dk^2 + dk + c} \\
 &= \frac{k^2 - k + 1}{k^2 + k + 1} \\
 &= \frac{(k^2 - k + 1)(k^2 + k + 1)}{(k^2 + k + 1)^2} \\
 &= \frac{k^4 + k^3 + k^2 - k^3 - k^2 - k + k^2 + k + 1}{(k^2 + k + 1)^2} \\
 &= \frac{k^4 + k^2 + 1}{(k^2 + k + 1)^2}
 \end{aligned}$$

$$\therefore \text{L.H.S.} = \text{R.H.S.}$$

Question 11.

Using properties of proportion, solve for x:

$$\text{(i)} \frac{\sqrt{x+5} + \sqrt{x-16}}{\sqrt{x+5} - \sqrt{x-16}} = \frac{7}{3}$$

$$\text{(ii)} \frac{\sqrt{x+1} + \sqrt{x-1}}{\sqrt{x+1} - \sqrt{x-1}} = \frac{4x-1}{2}$$

$$\text{(iii)} \frac{3x + \sqrt{9x^2 - 5}}{3x - \sqrt{9x^2 - 5}} = 5$$

Solution:

$$(i) \frac{\sqrt{x+5} + \sqrt{x-16}}{\sqrt{x+5} - \sqrt{x-16}} = \frac{7}{3}$$

Applying componendo and dividendo,

$$\frac{\sqrt{x+5} + \sqrt{x-16} + \sqrt{x+5} - \sqrt{x-16}}{\sqrt{x+5} + \sqrt{x-16} - \sqrt{x+5} + \sqrt{x-16}} = \frac{7+3}{7-3}$$

$$\frac{2\sqrt{x+5}}{2\sqrt{x-16}} = \frac{10}{4}$$

$$\frac{\sqrt{x+5}}{\sqrt{x-16}} = \frac{5}{2}$$

Squaring both sides,

$$\frac{x+5}{x-16} = \frac{25}{4}$$

$$4x + 20 = 25x - 400$$

$$21x = 420$$

$$x = \frac{420}{21} = 20$$

$$(ii) \frac{\sqrt{x+1} + \sqrt{x-1}}{\sqrt{x+1} - \sqrt{x-1}} = \frac{4x-1}{2}$$

Applying componendo and dividendo,

$$\frac{\sqrt{x+1} + \sqrt{x-1} + \sqrt{x+1} - \sqrt{x-1}}{\sqrt{x+1} + \sqrt{x-1} - \sqrt{x+1} + \sqrt{x-1}} = \frac{4x-1+2}{4x-1-2}$$

$$\frac{2\sqrt{x+1}}{2\sqrt{x-1}} = \frac{4x+1}{4x-3}$$

Squaring both sides,

$$\frac{x+1}{x-1} = \frac{16x^2 + 1 + 8x}{16x^2 + 9 - 24x}$$

Applying componendo and dividendo,

$$\frac{x+1+x-1}{x+1-x+1} = \frac{16x^2+1+8x+16x^2+9-24x}{16x^2+1+8x-16x^2-9+24x}$$

$$\frac{2x}{2} = \frac{32x^2+10-16x}{32x-8}$$

$$x = \frac{16x^2+5-8x}{16x-4}$$

$$16x^2-4x = 16x^2+5-8x$$

$$4x = 5$$

$$x = \frac{5}{4}$$

$$(iii) \frac{3x + \sqrt{9x^2 - 5}}{3x - \sqrt{9x^2 - 5}} = 5$$

Applying componendo and dividendo,

$$\frac{3x + \sqrt{9x^2 - 5} + 3x - \sqrt{9x^2 - 5}}{3x + \sqrt{9x^2 - 5} - 3x + \sqrt{9x^2 - 5}} = \frac{5+1}{5-1}$$

$$\frac{6x}{2\sqrt{9x^2 - 5}} = \frac{6}{4}$$

$$\frac{x}{\sqrt{9x^2 - 5}} = \frac{1}{2}$$

Squaring both sides,

$$\frac{x^2}{9x^2 - 5} = \frac{1}{4}$$

$$4x^2 = 9x^2 - 5$$

$$5x^2 = 5$$

$$x^2 = 1$$

$$x = 1$$

Question 12.

If $x = \frac{\sqrt{a+3b} + \sqrt{a-3b}}{\sqrt{a+3b} - \sqrt{a-3b}}$, prove that: $3bx^2 - 2ax + 3b = 0$.

Solution:

$$\text{Since, } \frac{x}{1} = \frac{\sqrt{a+3b} + \sqrt{a-3b}}{\sqrt{a+3b} - \sqrt{a-3b}}$$

Applying componendo and dividendo, we get,

$$\frac{x+1}{x-1} = \frac{\sqrt{a+3b} + \sqrt{a-3b} + \sqrt{a+3b} - \sqrt{a-3b}}{\sqrt{a+3b} + \sqrt{a-3b} - \sqrt{a+3b} + \sqrt{a-3b}}$$

$$\frac{x+1}{x-1} = \frac{2\sqrt{a+3b}}{2\sqrt{a-3b}}$$

Squaring both sides,

$$\frac{x^2 + 2x + 1}{x^2 - 2x + 1} = \frac{a+3b}{a-3b}$$

Again applying componendo and dividendo,

$$\frac{x^2 + 2x + 1 + x^2 - 2x + 1}{x^2 + 2x + 1 - x^2 + 2x - 1} = \frac{a+3b + a-3b}{a+3b - a+3b}$$

$$\frac{2(x^2 + 1)}{2(2x)} = \frac{2(a)}{2(3b)}$$

$$3b(x^2 + 1) = 2ax$$

$$3bx^2 + 3b = 2ax$$

$$3bx^2 - 2ax + 3b = 0.$$

Question 13.

Using the properties of proportion, solve for x,

$$\text{given } \frac{x^4 + 1}{2x^2} = \frac{17}{8}$$

Solution:

$$\frac{x^4 + 1}{2x^2} = \frac{17}{8}$$

Applying componendo and dividendo, we get

$$\frac{x^4 + 1 + 2x^2}{x^4 + 1 - 2x^2} = \frac{17 + 8}{17 - 8}$$

$$\Rightarrow \frac{(x^2)^2 + (1)^2 + 2 \times x^2 \times 1}{(x^2)^2 + (1)^2 - 2 \times x^2 \times 1} = \frac{25}{9}$$

$$\Rightarrow \frac{(x^2 + 1)^2}{(x^2 - 1)^2} = \frac{5^2}{3^2}$$

$$\Rightarrow \left(\frac{x^2 + 1}{x^2 - 1} \right)^2 = \left(\frac{5}{3} \right)^2$$

$$\Rightarrow \frac{x^2 + 1}{x^2 - 1} = \frac{5}{3}$$

Applying componendo and dividendo, we get

$$\frac{x^2 + 1 + x^2 - 1}{x^2 + 1 - x^2 + 1} = \frac{5 + 3}{5 - 3}$$

$$\Rightarrow \frac{2x^2}{2} = \frac{8}{2}$$

$$\Rightarrow x^2 = 4$$

$$\Rightarrow x = \pm 2$$

Question 14.

If $x = \frac{\sqrt{m+n} + \sqrt{m-n}}{\sqrt{m+n} - \sqrt{m-n}}$, express n in terms of x and m .

Solution:

$$x = \frac{\sqrt{m+n} + \sqrt{m-n}}{\sqrt{m+n} - \sqrt{m-n}}$$

Applying componendo and dividendo,

$$\frac{x+1}{x-1} = \frac{\sqrt{m+n} + \sqrt{m-n} + \sqrt{m+n} - \sqrt{m-n}}{\sqrt{m+n} + \sqrt{m-n} - \sqrt{m+n} + \sqrt{m-n}}$$

$$\frac{x+1}{x-1} = \frac{2\sqrt{m+n}}{2\sqrt{m-n}}$$

$$\frac{x+1}{x-1} = \frac{\sqrt{m+n}}{\sqrt{m-n}}$$

Squaring both sides,

$$\frac{x^2 + 2x + 1}{x^2 - 2x + 1} = \frac{m+n}{m-n}$$

Applying componendo and dividendo,

$$\frac{x^2 + 2x + 1 + x^2 - 2x + 1}{x^2 + 2x + 1 - x^2 + 2x - 1} = \frac{m+n+m-n}{m+n-m+n}$$

$$\frac{2x^2 + 2}{4x} = \frac{2m}{2n}$$

$$\frac{x^2 + 1}{2x} = \frac{m}{n}$$

$$\frac{x^2 + 1}{2mx} = \frac{1}{n}$$

$$n = \frac{2mx}{x^2 + 1}$$

Question 15.

If $\frac{x^3 + 3xy^2}{3x^2y + y^3} = \frac{m^3 + 3mn^2}{3m^2n + n^3}$, show that:

$$nx = my.$$

Solution:

$$\frac{x^3 + 3xy^2}{3x^2y + y^3} = \frac{m^3 + 3mn^2}{3m^2n + n^3}$$

Applying componendo and dividendo,

$$\frac{x^3 + 3xy^2 + 3x^2y + y^3}{x^3 + 3xy^2 - 3x^2y - y^3} = \frac{m^3 + 3mn^2 + 3m^2n + n^3}{m^3 + 3mn^2 - 3m^2n - n^3}$$

$$\frac{(x+y)^3}{(x-y)^3} = \frac{(m+n)^3}{(m-n)^3}$$

$$\frac{x+y}{x-y} = \frac{m+n}{m-n}$$

Applying componendo and dividendo,

$$\frac{x+y+x-y}{x+y-x+y} = \frac{m+n+m-n}{m+n-m+n}$$

$$\frac{2x}{2y} = \frac{2m}{2n}$$

$$\frac{x}{y} = \frac{m}{n}$$

$$nx = my$$

Exercise 7D

Question 1.

If $a : b = 3 : 5$, find:

$(10a + 3b) : (5a + 2b)$

Solution:

$$\text{Given, } \frac{a}{b} = \frac{3}{5}$$

$$\begin{aligned} & \frac{10a + 3b}{5a + 2b} \\ &= \frac{10\left(\frac{a}{b}\right) + 3}{5\left(\frac{a}{b}\right) + 2} \end{aligned}$$

$$\begin{aligned}
 &= \frac{10\left(\frac{3}{5}\right) + 3}{5\left(\frac{3}{5}\right) + 2} \\
 &= \frac{6 + 3}{3 + 2} \\
 &= \frac{9}{5}
 \end{aligned}$$

Question 2.

If $5x + 6y : 8x + 5y = 8 : 9$, find $x : y$.

Solution:

$$\begin{aligned}
 \frac{5x + 6y}{8x + 5y} &= \frac{8}{9} \\
 45x + 54y &= 64x + 40y \\
 64x - 45x &= 54y - 40y \\
 19x &= 14y \\
 \frac{x}{y} &= \frac{14}{19}
 \end{aligned}$$

Question 3.

If $(3x - 4y) : (2x - 3y) = (5x - 6y) : (4x - 5y)$, find $x : y$.

Solution:

$$\begin{aligned}
 (3x - 4y) : (2x - 3y) &= (5x - 6y) : (4x - 5y) \\
 \frac{3x - 4y}{2x - 3y} &= \frac{5x - 6y}{4x - 5y} \\
 \text{Applying componendo and dividendo,} \\
 \frac{3x - 4y + 2x - 3y}{3x - 4y - 2x + 3y} &= \frac{5x - 6y + 4x - 5y}{5x - 6y - 4x + 5y} \\
 \frac{5x - 7y}{x - y} &= \frac{9x - 11y}{x - y}
 \end{aligned}$$

$$5x - 7y = 9x - 11y$$

$$11y - 7y = 9x - 5x$$

$$4y = 4x$$

$$\frac{x}{y} = \frac{1}{1}$$

$$x : y = 1 : 1$$

Question 4.

Find the:

(i) duplicate ratio of $2\sqrt{2} : 3\sqrt{5}$

(ii) triplicate ratio of $2a : 3b$

(iii) sub-duplicate ratio of $9x^2a^4 : 25y^6b^2$

(iv) sub-triplicate ratio of $216 : 343$

(v) reciprocal ratio of $3 : 5$

(vi) ratio compounded of the duplicate ratio of $5 : 6$, the reciprocal ratio of $25 : 42$ and the sub-duplicate ratio of $36 : 49$.

Solution:

$$(i) \text{ Duplicate ratio of } 2\sqrt{2} : 3\sqrt{5} = (2\sqrt{2})^2 : (3\sqrt{5})^2 = 8 : 45$$

$$(ii) \text{ Triplicate ratio of } 2a : 3b = (2a)^3 : (3b)^3 = 8a^3 : 27b^3$$

$$(iii) \text{ Sub-duplicate ratio of } 9x^2a^4 : 25y^6b^2 = \sqrt{9x^2a^4} : \sqrt{25y^6b^2} = 3xa^2 : 5y^3b$$

$$(iv) \text{ Sub-triplicate ratio of } 216 : 343 = \sqrt[3]{216} : \sqrt[3]{343} = 6 : 7$$

$$(v) \text{ Reciprocal ratio of } 3 : 5 = 5 : 3$$

$$(vi) \text{ Duplicate ratio of } 5 : 6 = 25 : 36$$

$$\text{Reciprocal ratio of } 25 : 42 = 42 : 25$$

$$\text{Sub-duplicate ratio of } 36 : 49 = 6 : 7$$

$$\text{Required compound ratio} = \frac{25 \times 42 \times 6}{36 \times 25 \times 7} = 1 : 1$$

Question 5.

Find the value of x , if:

(i) $(2x + 3) : (5x - 38)$ is the duplicate ratio of $\sqrt{5} : \sqrt{6}$

(ii) $(2x + 1) : (3x + 13)$ is the sub-duplicate ratio of $9 : 25$.

(iii) $(3x - 7) : (4x + 3)$ is the sub-triplicate ratio of $8 : 27$.

Solution:

(i) $(2x + 3) : (5x - 38)$ is the duplicate ratio of $\sqrt{5} : \sqrt{6}$

Duplicate ratio of $\sqrt{5} : \sqrt{6} = 5 : 6$

$$\frac{2x + 3}{5x - 38} = \frac{5}{6}$$

$$12x + 18 = 25x - 190$$

$$25x - 12x = 190 + 18$$

$$13x = 208$$

$$x = \frac{208}{13} = 16$$

(ii) $(2x + 1) : (3x + 13)$ is the sub-duplicate ratio of 9: 25

Sub-duplicate ratio of 9: 25 = 3: 5

$$\frac{2x + 1}{3x + 13} = \frac{3}{5}$$

$$10x + 5 = 9x + 39$$

$$10x - 9x = 39 - 5$$

$$x = 34$$

(iii) $(3x - 7) : (4x + 3)$ is the sub-triplicate ratio of 8: 27

Sub-triplicate ratio of 8: 27 = 2: 3

$$\frac{3x - 7}{4x + 3} = \frac{2}{3}$$

$$9x - 21 = 8x + 6$$

$$9x - 8x = 6 + 21$$

$$x = 27$$

Question 6.

What quantity must be added to each term of the ratio $x : y$ so that it may become equal to $c : d$?

Solution:

Let the required quantity which is to be added be p .

Then, we have:

$$\frac{x+p}{y+p} = \frac{c}{d}$$

$$dx + pd = cy + cp$$

$$pd - cp = cy - dx$$

$$p(d - c) = cy - dx$$

$$p = \frac{cy - dx}{d - c}$$

Question 7.

A woman reduces her weight in the ratio 7 : 5. What does her weight become if originally it was 84 kg?

Solution:

Let the reduced weight be x.

Original weight = 84 kg

Thus, we have

$$84 : x = 7 : 5$$

$$\Rightarrow \frac{84}{x} = \frac{7}{5}$$

$$\Rightarrow 84 \times 5 = 7 \times x$$

$$\Rightarrow x = \frac{84 \times 5}{7}$$

$$\Rightarrow x = 60$$

Thus, her reduced weight is 60 kg.

Question 8.

If $15(2x^2 - y^2) = 7xy$, find x : y; if x and y both are positive.

Solution:

$$15(2x^2 - y^2) = 7xy$$

$$\frac{2x^2 - y^2}{xy} = \frac{7}{15}$$

$$\frac{2x}{y} - \frac{y}{x} = \frac{7}{15}$$

$$\text{Let } \frac{x}{y} = a$$

$$\therefore 2a - \frac{1}{a} = \frac{7}{15}$$

$$\frac{2a^2 - 1}{a} = \frac{7}{15}$$

$$30a^2 - 15 = 7a$$

$$30a^2 - 7a - 15 = 0$$

$$30a^2 - 25a + 18a - 15 = 0$$

$$5a(6a - 5) + 3(6a - 5) = 0$$

$$(6a - 5)(5a + 3) = 0$$

$$a = \frac{5}{6}, -\frac{3}{5}$$

But, a cannot be negative.

$$\therefore a = \frac{5}{6}$$

$$\Rightarrow \frac{x}{y} = \frac{5}{6}$$

$$\Rightarrow x : y = 5 : 6$$

Question 9.

Find the:

- (i) fourth proportional to $2xy$, x^2 and y^2 .
- (ii) third proportional to $a^2 - b^2$ and $a + b$.
- (iii) mean proportional to $(x - y)$ and $(x^3 - x^2y)$.

Solution:

(i) Let the fourth proportional to $2xy$, x^2 and y^2 be n .

$$\Rightarrow 2xy : x^2 = y^2 : n$$

$$\Rightarrow 2xy \times n = x^2 \times y^2$$

$$\Rightarrow n = \frac{x^2 y^2}{2xy} = \frac{xy}{2}$$

(ii) Let the third proportional to $a^2 - b^2$ and $a + b$ be n .

$$\Rightarrow a^2 - b^2, a + b \text{ and } n \text{ are in continued proportion.}$$

$$\Rightarrow a^2 - b^2 : a + b = a + b : n$$

$$\Rightarrow n = \frac{(a+b)^2}{a^2 - b^2} = \frac{(a+b)^2}{(a+b)(a-b)} = \frac{a+b}{a-b}$$

(iii) Let the mean proportional to $(x-y)$ and $(x^3 - x^2y)$ be n .

$\Rightarrow (x-y), n, (x^3 - x^2y)$ are in continued proportion

$\Rightarrow (x-y) : n = n : (x^3 - x^2y)$

$$\Rightarrow n^2 = (x-y)(x^3 - x^2y)$$

$$\Rightarrow n^2 = x^2(x-y)(x-y)$$

$$\Rightarrow n^2 = x^2(x-y)^2$$

$$\Rightarrow n = x(x-y)$$

Question 10.

Find two numbers such that the mean proportional between them is 14 and third proportional to them is 112.

Solution:

Let the required numbers be a and b .

Given, 14 is the mean proportional between a and b .

$$\Rightarrow a : 14 = 14 : b$$

$$\Rightarrow ab = 196$$

$$\Rightarrow a = \frac{196}{b} \dots (1)$$

Also, given, third proportional to a and b is 112.

$$\Rightarrow a : b = b : 112$$

$$\Rightarrow b^2 = 112a \dots (2)$$

Using (1), we have:

$$b^2 = 112 \times \frac{196}{b}$$

$$b^3 = (14)^3(2)^3$$

$$b = 28$$

From (1),

$$a = \frac{196}{28} = 7$$

Thus, the two numbers are 7 and 28.

Question 11.

If x and y be unequal and $x : y$ is the duplicate ratio of $x + z$ and $y + z$, prove that z is mean proportional between x and y .

Solution:

$$\text{Given, } \frac{x}{y} = \frac{(x+z)^2}{(y+z)^2}$$

$$x(y^2 + z^2 + 2yz) = y(x^2 + z^2 + 2xz)$$

$$xy^2 + xz^2 + 2xyz = x^2y + yz^2 + 2xyz$$

$$xy^2 + xz^2 = x^2y + yz^2$$

$$xy^2 - x^2y = yz^2 - xz^2$$

$$xy(y - x) = z^2(y - x)$$

$$xy = z^2$$

Hence, z is mean proportional between x and y .

Question 12.

If $x = \frac{2ab}{a+b}$, find the value of $\frac{x+a}{x-a} + \frac{x+b}{x-b}$.

Solution:

$$x = \frac{2ab}{a+b}$$

$$\frac{x}{a} = \frac{2b}{a+b}$$

Applying componendo and dividendo,

$$\frac{x+a}{x-a} = \frac{2b+a+b}{2b-a-b}$$

$$\frac{x+a}{x-a} = \frac{3b+a}{b-a} \quad \dots (1)$$

$$\text{Also, } x = \frac{2ab}{a+b}$$

$$\frac{x}{b} = \frac{2a}{a+b}$$

Applying componendo and dividendo,

$$\frac{x+b}{x-b} = \frac{2a+a+b}{2a-a-b}$$

$$\frac{x+b}{x-b} = \frac{3a+b}{a-b} \quad \dots (2)$$

From (1) and (2),

$$\frac{x+a}{x-a} + \frac{x+b}{x-b} = \frac{3b+a}{b-a} + \frac{3a+b}{a-b}$$

$$\frac{x+a}{x-a} + \frac{x+b}{x-b} = \frac{-3b-a+3a+b}{a-b}$$

$$\frac{x+a}{x-a} + \frac{x+b}{x-b} = \frac{2a-2b}{a-b} = 2$$

Question 13.

If $(4a + 9b)(4c - 9d) = (4a - 9b)(4c + 9d)$, prove that:
 $a : b = c : d$.

Solution:

$$\text{Given, } \frac{4a+9b}{4a-9b} = \frac{4c+9d}{4c-9d}$$

Applying componendo and dividendo,

$$\frac{4a+9b+4a-9b}{4a+9b-4a+9b} = \frac{4c+9d+4c-9d}{4c+9d-4c+9d}$$

$$\frac{8a}{18b} = \frac{8c}{18d}$$

$$\frac{a}{b} = \frac{c}{d}$$

Question 14.

If $\frac{a}{b} = \frac{c}{d}$, show that:

$$(a+b) : (c+d) = \sqrt{a^2+b^2} : \sqrt{c^2+d^2}$$

Solution:

$$\text{Let } \frac{a}{b} = \frac{c}{d} = k(\text{say})$$

$$\Rightarrow a = bk, c = dk$$

$$\begin{aligned}\text{L.H.S.} &= \frac{a+b}{c+d} \\ &= \frac{bk+b}{dk+d} \\ &= \frac{b(k+1)}{d(k+1)} \\ &= \frac{b}{d}\end{aligned}$$

$$\begin{aligned}\text{R.H.S.} &= \frac{\sqrt{a^2+b^2}}{\sqrt{c^2+d^2}} \\ &= \frac{\sqrt{(bk)^2+b^2}}{\sqrt{(dk)^2+d^2}} \\ &= \frac{\sqrt{b^2(k^2+1)}}{\sqrt{d^2(k^2+1)}} \\ &= \frac{\sqrt{b^2}}{\sqrt{d^2}} \\ &= \frac{b}{d}\end{aligned}$$

$$\therefore \text{L.H.S.} = \text{R.H.S.}$$

Question 15.

There are 36 members in a student council in a school and the ratio of the number of boys to the number of girls is 3: 1. How many more girls should be added to the council so that the ratio of the number of boys to the number of girls may be 9: 5?

Solution:

Ratio of number of boys to the number of girls = 3: 1

Let the number of boys be $3x$ and number of girls be x .

$$3x + x = 36$$

$$4x = 36$$

$$x = 9$$

∴ Number of boys = 27

Number of girls = 9

Let n number of girls be added to the council.

From given information, we have:

$$\begin{aligned}\frac{27}{9+n} &= \frac{9}{5} \\ 135 &= 81 + 9n \\ 9n &= 54 \\ n &= 6\end{aligned}$$

Thus, 6 girls are added to the council.

Question 16.

If $7x - 15y = 4x + y$, find the value of $x : y$. Hence, use componendo and dividendo to find the values of:

$$\begin{aligned}\text{(i)} & \frac{9x + 5y}{9x - 5y} \\ \text{(ii)} & \frac{3x^2 + 2y^2}{3x^2 - 2y^2}\end{aligned}$$

Solution:

$$\begin{aligned}7x - 15y &= 4x + y \\ 7x - 4x &= y + 15y \\ 3x &= 16y \\ \frac{x}{y} &= \frac{16}{3}\end{aligned}$$

$$\text{(i)} \frac{x}{y} = \frac{16}{3}$$

$$\Rightarrow \frac{9x}{5y} = \frac{144}{15}$$

$$\Rightarrow \frac{9x + 5y}{9x - 5y} = \frac{144 + 15}{144 - 15}$$

$$\Rightarrow \frac{9x + 5y}{9x - 5y} = \frac{159}{129} = \frac{53}{43}$$

(Multiplying both sides by $\frac{9}{5}$)

(Applying componendo and dividendo)

$$(ii) \frac{x}{y} = \frac{16}{3}$$

$$\Rightarrow \frac{x^2}{y^2} = \frac{256}{9}$$

$$\Rightarrow \frac{3x^2}{2y^2} = \frac{768}{18} = \frac{128}{3} \quad \left(\text{Multiplying both sides by } \frac{3}{2} \right)$$

$$\Rightarrow \frac{3x^2 + 2y^2}{3x^2 - 2y^2} = \frac{128 + 3}{128 - 3} \quad (\text{Applying componendo and dividendo})$$

$$\Rightarrow \frac{3x^2 + 2y^2}{3x^2 - 2y^2} = \frac{131}{125}$$

Question 17.

If $\frac{4m + 3n}{4m - 3n} = \frac{7}{4}$, use properties of proportion to find:

(i) $m : n$

$$(ii) \frac{2m^2 - 11n^2}{2m^2 + 11n^2}$$

Solution:

$$(i) \text{ Given, } \frac{4m + 3n}{4m - 3n} = \frac{7}{4}$$

Applying componendo and dividendo,

$$\frac{4m + 3n + 4m - 3n}{4m + 3n - 4m + 3n} = \frac{7 + 4}{7 - 4}$$

$$\frac{8m}{6n} = \frac{11}{3}$$

$$\frac{m}{n} = \frac{11}{4}$$

$$(ii) \frac{m}{n} = \frac{11}{4}$$

$$\frac{m^2}{n^2} = \frac{121}{16}$$

$$\frac{2m^2}{11n^2} = \frac{2 \times 121}{11 \times 16} \quad \left(\text{Multiplying both sides by } \frac{2}{11} \right)$$

$$\frac{2m^2}{11n^2} = \frac{11}{8}$$

$$\frac{2m^2 + 11n^2}{2m^2 - 11n^2} = \frac{11 + 8}{11 - 8} \quad (\text{Applying componendo and dividendo})$$

$$\frac{2m^2 + 11n^2}{2m^2 - 11n^2} = \frac{19}{3}$$

$$\frac{2m^2 - 11n^2}{2m^2 + 11n^2} = \frac{3}{19} \quad (\text{Applying invertendo})$$

Question 18.

If x, y, z are in continued proportion, prove that $\frac{(x+y)^2}{(y+z)^2} = \frac{x}{z}$.

Solution:

$\because x, y, z$ are in continued proportion,

$$\therefore \frac{x}{y} = \frac{y}{z} \Rightarrow y^2 = zx \dots (1)$$

Therefore,

$$\frac{x+y}{y} = \frac{y+z}{z} \quad (\text{By componendo})$$

$$\Rightarrow \frac{x+y}{y+z} = \frac{y}{z} \quad (\text{By alternendo})$$

$$\Rightarrow \frac{(x+y)^2}{(y+z)^2} = \frac{y^2}{z^2} \quad (\text{squaring both sides})$$

$$\Rightarrow \frac{(x+y)^2}{(y+z)^2} = \frac{zx}{z^2} \quad [\text{from (1)}]$$

$$\Rightarrow \frac{(x+y)^2}{(y+z)^2} = \frac{x}{z}$$

Hence Proved.

Question 19.

$$\text{Given } x = \frac{\sqrt{a^2 + b^2} + \sqrt{a^2 - b^2}}{\sqrt{a^2 + b^2} - \sqrt{a^2 - b^2}}.$$

Use componendo and dividendo to prove that $b^2 = \frac{2a^2x}{x^2 + 1}$.

Solution:

$$x = \frac{\sqrt{a^2 + b^2} + \sqrt{a^2 - b^2}}{\sqrt{a^2 + b^2} - \sqrt{a^2 - b^2}}$$

By componendo and dividendo,

$$\frac{x+1}{x-1} = \frac{\sqrt{a^2 + b^2} + \sqrt{a^2 - b^2} + \sqrt{a^2 + b^2} - \sqrt{a^2 - b^2}}{\sqrt{a^2 + b^2} + \sqrt{a^2 - b^2} - \sqrt{a^2 + b^2} + \sqrt{a^2 - b^2}}$$

$$\frac{x+1}{x-1} = \frac{2\sqrt{a^2 + b^2}}{2\sqrt{a^2 - b^2}}$$

Squaring both sides,

$$\frac{x^2 + 2x + 1}{x^2 - 2x + 1} = \frac{a^2 + b^2}{a^2 - b^2}$$

By componendo and dividendo,

$$\frac{(x^2 + 2x + 1) + (x^2 - 2x + 1)}{(x^2 + 2x + 1) - (x^2 - 2x + 1)} = \frac{(a^2 + b^2) + (a^2 - b^2)}{(a^2 + b^2) - (a^2 - b^2)}$$

$$\Rightarrow \frac{2(x^2 + 1)}{4x} = \frac{2a^2}{2b^2}$$

$$\Rightarrow \frac{x^2 + 1}{2x} = \frac{a^2}{b^2}$$

$$\Rightarrow b^2 = \frac{2a^2x}{x^2 + 1}$$

Hence Proved.

Question 20.

If $\frac{x^2 + y^2}{x^2 - y^2} = 2\frac{1}{8}$, find:

(i) $\frac{x}{y}$

(ii) $\frac{x^3 + y^3}{x^3 - y^3}$

Solution:

(i) Given, $\frac{x^2 + y^2}{x^2 - y^2} = 2\frac{1}{8}$

$$\frac{x^2 + y^2}{x^2 - y^2} = \frac{17}{8}$$

Applying componendo and dividendo,

$$\frac{x^2 + y^2 + x^2 - y^2}{x^2 + y^2 - x^2 + y^2} = \frac{17 + 8}{17 - 8}$$

$$\frac{2x^2}{2y^2} = \frac{25}{9}$$

$$\frac{x^2}{y^2} = \frac{25}{9}$$

$$\frac{x}{y} = \frac{5}{3} = 1\frac{2}{3}$$

(ii) $\frac{x^3 + y^3}{x^3 - y^3}$

$$= \frac{\left(\frac{x}{y}\right)^3 + 1}{\left(\frac{x}{y}\right)^3 - 1}$$

$$= \frac{\left(\frac{5}{3}\right)^3 + 1}{\left(\frac{5}{3}\right)^3 - 1}$$

$$= \frac{\left(\frac{5}{3}\right)^3 + 1}{\left(\frac{5}{3}\right)^3 - 1}$$

$$= \frac{\frac{125}{27} + 1}{\frac{125}{27} - 1}$$

$$= \frac{\frac{125}{27} + 1}{\frac{125}{27} - 1}$$

$$\begin{aligned}
 & \frac{125+27}{27} \\
 &= \frac{125-27}{27} \\
 &= \frac{125+27}{125-27} \\
 &= \frac{76}{49} = 1\frac{27}{49}
 \end{aligned}$$

Question 21.

Using componendo and dividendo find the value of x:

$$\frac{\sqrt{3x+4} + \sqrt{3x-5}}{\sqrt{3x+4} - \sqrt{3x-5}} = 9$$

Solution:

$$\frac{\sqrt{3x+4} + \sqrt{3x-5}}{\sqrt{3x+4} - \sqrt{3x-5}} = \frac{9}{1}$$

Applying componendo and dividendo, we have

$$\frac{\sqrt{3x+4} + \sqrt{3x-5} + \sqrt{3x+4} - \sqrt{3x-5}}{\sqrt{3x+4} + \sqrt{3x-5} - \sqrt{3x+4} + \sqrt{3x-5}} = \frac{9+1}{9-1}$$

$$\Rightarrow \frac{2\sqrt{3x+4}}{2\sqrt{3x-5}} = \frac{10}{8}$$

$$\Rightarrow \frac{\sqrt{3x+4}}{\sqrt{3x-5}} = \frac{5}{4}$$

Squaring both sides, we have

$$\frac{3x+4}{3x-5} = \frac{25}{16}$$

$$\Rightarrow 16(3x+4) = 25(3x-5)$$

$$\Rightarrow 48x + 64 = 75x - 125$$

$$\Rightarrow 75x - 48x = 64 + 125$$

$$\Rightarrow 27x = 189$$

$$\Rightarrow x = \frac{189}{27}$$

$$\Rightarrow x = 7$$

Question 22.

If $x = \frac{\sqrt{a+1} + \sqrt{a-1}}{\sqrt{a+1} - \sqrt{a-1}}$, using properties of proportion,

show that:

$$x^2 - 2ax + 1 = 0$$

Solution:

$$\text{Given that, } x = \frac{\sqrt{a+1} + \sqrt{a-1}}{\sqrt{a+1} - \sqrt{a-1}}$$

By applying Componendo-Dividendo,

$$\frac{x+1}{x-1} = \frac{(\sqrt{a+1} + \sqrt{a-1}) + (\sqrt{a+1} - \sqrt{a-1})}{(\sqrt{a+1} + \sqrt{a-1}) - (\sqrt{a+1} - \sqrt{a-1})}$$

$$\Rightarrow \frac{x+1}{x-1} = \frac{2\sqrt{a+1}}{2\sqrt{a-1}}$$

$$\Rightarrow \frac{x+1}{x-1} = \frac{\sqrt{a+1}}{\sqrt{a-1}}$$

Squaring both the sides of the equation, we have,

$$\Rightarrow \left(\frac{x+1}{x-1}\right)^2 = \frac{a+1}{a-1}$$

$$\Rightarrow (x+1)^2 (a-1) = (x-1)^2 (a+1)$$

$$\Rightarrow (x^2 + 2x + 1)(a-1) = (x^2 - 2x + 1)(a+1)$$

$$\Rightarrow a(x^2 + 2x + 1) - (x^2 + 2x + 1) = a(x^2 - 2x + 1) + (x^2 - 2x + 1)$$

$$\Rightarrow 4ax = 2x^2 + 2$$

$$\Rightarrow 2ax = x^2 + 1$$

$$\Rightarrow x^2 - 2ax + 1 = 0$$

Question 23.

$$\text{Given } \frac{x^3 + 12x}{6x^2 + 8} = \frac{y^3 + 27y}{9y^2 + 27}.$$

Using componendo and dividendo, find $x : y$.

Solution:

$$\frac{x^3 + 12x}{6x^2 + 8} = \frac{y^3 + 27y}{9y^2 + 27}$$

Applying componendo and dividendo, we get

$$\frac{x^3 + 12x + 6x^2 + 8}{x^3 + 12x - 6x^2 - 8} = \frac{y^3 + 27y + 9y^2 + 27}{y^3 + 27y - 9y^2 - 27}$$

$$\Rightarrow \frac{x^3 + 3(1)(4)x + 3(1)(2)x^2 + 2^3}{x^3 + 3(1)(4)x - 3(1)(2)x^2 - 2^3} = \frac{y^3 + 3(1)(9)y + 3(1)(3)y^2 + 3^3}{y^3 + 3(1)(9)y - 3(1)(3)y^2 - 3^3}$$

$$\Rightarrow \frac{x^3 + 3(1)(4)x + 3(1)(2)x^2 + 2^3}{x^3 - 3(1)(2)x^2 + 3(1)(4)x - 2^3} = \frac{y^3 + 3(1)(9)y + 3(1)(3)y^2 + 3^3}{y^3 - 3(1)(3)y^2 + 3(1)(9)y - 3^3}$$

$$\Rightarrow \frac{(x+2)^3}{(x-2)^3} = \frac{(y+3)^3}{(y-3)^3}$$

$$\Rightarrow \frac{x+2}{x-2} = \frac{y+3}{y-3}$$

Again applying componendo and dividendo, we get

$$\frac{x+2+x-2}{x+2-x+2} = \frac{y+3+y-3}{y+3-y+3}$$

$$\Rightarrow \frac{2x}{4} = \frac{2y}{6}$$

$$\Rightarrow \frac{x}{2} = \frac{y}{3}$$

Applying alternendo, we get

$$\frac{x}{y} = \frac{2}{3}$$

$$\Rightarrow x : y = 2 : 3$$

Question 24.

$$\text{Let } \frac{x}{a} = \frac{y}{b} = \frac{z}{c} = k$$

$$\Rightarrow x = ak, y = bk, z = ck$$

$$\text{L.H.S.} = \frac{x^3}{a^3} + \frac{y^3}{b^3} + \frac{z^3}{c^3}$$

$$\begin{aligned}
&= \frac{(ak)^3}{a^3} + \frac{(bk)^3}{b^3} + \frac{(ck)^3}{c^3} \\
&= \frac{a^3 k^3}{a^3} + \frac{b^3 k^3}{b^3} + \frac{c^3 k^3}{c^3} \\
&= k^3 + k^3 + k^3 \\
&= 3k^3
\end{aligned}$$

$$\begin{aligned}
\text{R.H.S.} &= \frac{3xyz}{abc} \\
&= \frac{3(ak)(bk)(ck)}{abc} \\
&= 3k^3
\end{aligned}$$

$$\Rightarrow \text{L.H.S.} = \text{R.H.S.}$$

$$\text{i.e. } \frac{x^3}{a^3} + \frac{y^3}{b^3} + \frac{z^3}{c^3} = \frac{3xyz}{abc}$$

Question 25.

Given that b is the mean proportion between a and c .

$$\therefore \frac{a}{b} = \frac{b}{c} = k$$

$$\Rightarrow b = ck ; a = bk = (ck)k = ck^2$$

$$\begin{aligned}
\text{L.H.S.} &= \frac{a^4 + a^2 b^2 + b^4}{b^4 + b^2 c^2 + c^4} \\
&= \frac{(ck^2)^4 + (ck^2)^2 (ck)^2 + (ck)^4}{(ck)^4 + (ck)^2 c^2 + c^4} \\
&= \frac{c^4 k^8 + c^4 k^6 + c^4 k^4}{c^4 k^4 + c^4 k^2 + c^4} \\
&= \frac{c^4 k^4 (k^4 + k^2 + 1)}{c^4 (k^4 + k^2 + 1)} \\
&= k^4 \quad \dots (i)
\end{aligned}$$

$$\text{R. H. S.} = \frac{a^2}{c^2} = \frac{(ck^2)^2}{c^2} = \frac{c^2 k^4}{c^2} = k^4 \quad \dots (ii)$$

From (i) and (ii), we get

$$\text{L.H.S} = \text{R.H.S}$$

$$\Rightarrow \frac{a^4 + a^2b^2 + b^4}{b^4 + b^2c^2 + c^4} = \frac{a^2}{c^2}$$

Hence proved.

Question 26.

$$\frac{7m + 2n}{7m - 2n} = \frac{5}{3}$$

Applying Componendo and Dividendo, we get

$$\frac{7m + 2n + 7m - 2n}{7m + 2n - 7m + 2n} = \frac{5 + 3}{5 - 3}$$

$$\Rightarrow \frac{14m}{4n} = \frac{8}{2}$$

$$\Rightarrow \frac{m}{n} = \frac{8 \times 4}{2 \times 14}$$

$$\Rightarrow \frac{m}{n} = \frac{8}{7}$$

$$\Rightarrow m : n = 8 : 7$$

ii.

From (i),

$$\frac{m}{n} = \frac{8}{7}$$

$$\Rightarrow \frac{m^2}{n^2} = \frac{64}{49}$$

Applying Componendo and Dividendo, we get

$$\frac{m^2 + n^2}{m^2 - n^2} = \frac{64 + 49}{64 - 49}$$

$$\Rightarrow \frac{m^2 + n^2}{m^2 - n^2} = \frac{64 + 49}{64 - 49}$$

$$\Rightarrow \frac{m^2 + n^2}{m^2 - n^2} = \frac{113}{15} = 7 \frac{8}{15}$$

Question 27.

i. $(2x^2 - 5y^2): xy = 1:3$

$$\Rightarrow \frac{2x^2 - 5y^2}{xy} = \frac{1}{3}$$

$$\Rightarrow \frac{2x}{y} - \frac{5y}{x} = \frac{1}{3}$$

Put $\frac{x}{y} = a$, we get

$$\Rightarrow 2a - 5\frac{1}{a} = \frac{1}{3}$$

$$\Rightarrow 3(2a^2 - 5) = a$$

$$\Rightarrow 6a^2 - a - 15 = 0$$

$$\Rightarrow 6a^2 + 9a - 10a - 15 = 0$$

$$\Rightarrow 3a(2a + 3) - 5(2a + 3) = 0$$

$$\Rightarrow (2a + 3)(3a - 5) = 0$$

$$\Rightarrow (2a + 3) = 0 \text{ or } (3a - 5) = 0$$

$$\Rightarrow a = -\frac{3}{2} \text{ or } a = \frac{5}{3}$$

$a = -\frac{3}{2}$ is not acceptable, as x and y both are positive.

$$\therefore a = \frac{5}{3} \Rightarrow \frac{x}{y} = \frac{5}{3}$$

$$\Rightarrow x : y = 5 : 3$$

ii.

$$16\left(\frac{a - x}{a + x}\right)^3 = \frac{a + x}{a - x}$$

$$\Rightarrow 16 = \left(\frac{a + x}{a - x}\right)^4$$

$$\Rightarrow (2)^4 = \left(\frac{a + x}{a - x}\right)^4$$

$$\Rightarrow \frac{a + x}{a - x} = \pm 2$$

$$\Rightarrow \frac{a+x}{a-x} = \frac{2}{1} \quad \text{or} \quad \frac{a+x}{a-x} = \frac{-2}{1}$$

Applying Componendo and Dividendo, we get

$$\Rightarrow \frac{a+x+a-x}{a+x-a+x} = \frac{3}{1} \quad \text{or} \quad \frac{a+x+a-x}{a+x-a+x} = \frac{-1}{-3}$$

$$\Rightarrow \frac{2a}{2x} = 3 \quad \text{or} \quad \frac{2a}{2x} = \frac{1}{3}$$

$$\Rightarrow x = \frac{a}{3} \quad \text{or} \quad x = 3a$$

Question 28.

If $\frac{x}{a} = \frac{y}{b} = \frac{z}{c}$, prove that:

$$\frac{ax-by}{(a+b)(x-y)} + \frac{by-cz}{(b+c)(y-z)} + \frac{cz-ax}{(c+a)(z-x)} = 3$$

Solution:

$$\text{Let } \frac{x}{a} = \frac{y}{b} = \frac{z}{c} = k(\text{say})$$

$$\Rightarrow x = ak, y = bk, z = ck$$

L.H.S.

$$\begin{aligned} &= \frac{ax-by}{(a+b)(x-y)} + \frac{by-cz}{(b+c)(y-z)} + \frac{cz-ax}{(c+a)(z-x)} \\ &= \frac{a(ak)-b(bk)}{(a+b)(ak-bk)} + \frac{b(bk)-c(ck)}{(b+c)(bk-ck)} + \frac{c(ck)-a(ak)}{(c+a)(ck-ak)} \\ &= \frac{k(a^2-b^2)}{k(a+b)(a-b)} + \frac{k(b^2-c^2)}{k(b+c)(b-c)} + \frac{k(c^2-a^2)}{k(c+a)(c-a)} \\ &= \frac{k(a^2-b^2)}{k(a^2-b^2)} + \frac{k(b^2-c^2)}{k(b^2-c^2)} + \frac{k(c^2-a^2)}{k(c^2-a^2)} \\ &= 1+1+1=3=R.H.S. \end{aligned}$$

Question 29.

If q is the mean proportional between p and r , prove that:

$$\frac{p^3 + q^3 + r^3}{p^2 q^2 r^2} = \frac{1}{p^3} + \frac{1}{q^3} + \frac{1}{r^3}.$$

Solution:

Since, q is the mean proportional between p and r ,

$$q^2 = pr$$

$$\begin{aligned} \text{L.H.S.} &= \frac{p^3 + q^3 + r^3}{p^2 q^2 r^2} \\ &= \frac{p^3 + (pr)q + r^3}{p^2 (pr)^2} \\ &= \frac{p^3 + pqr + r^3}{p^3 r^3} \\ &= \frac{1}{r^3} + \frac{q}{p^2 r^2} + \frac{1}{p^3} \\ &= \frac{1}{r^3} + \frac{q}{(q^2)^2} + \frac{1}{p^3} \\ &= \frac{1}{r^3} + \frac{1}{q^3} + \frac{1}{p^3} \\ &= \text{R.H.S.} \end{aligned}$$

Question 30.

If a , b and c are in continued proportion, prove that:

$$a : c = (a^2 + b^2) : (b^2 + c^2)$$

Solution:

Given, a , b and c are in continued proportion.

$$\Rightarrow a : b = b : c$$

$$\text{Let } \frac{a}{b} = \frac{b}{c} = k \text{ (say)}$$

$$\Rightarrow a = bk, b = ck$$

$$\Rightarrow a = ck^2, b = ck$$

$$\text{Now, L.H.S.} = \frac{a}{c} = \frac{ck^2}{c} = k^2$$

$$\begin{aligned}
 \text{R.H.S.} &= \frac{a^2 + b^2}{b^2 + c^2} \\
 &= \frac{(ck^2)^2 + (ck)^2}{(ck)^2 + c^2} \\
 &= \frac{c^2k^4 + c^2k^2}{c^2k^2 + c^2} \\
 &= \frac{c^2k^2(k^2 + 1)}{c^2(k^2 + 1)} \\
 &= k^2
 \end{aligned}$$

$$\therefore \text{L.H.S.} = \text{R.H.S.}$$