ICSE Paper 2013

CHEMISTRY

SECTION-I (40 Marks)

Attempt all questions from this Section

Question 1.

を の としてのは関係といいを変し、例如

| (a) | From the list given below, select the word(s) required to correctly complete |
|-----|--|
| | blanks (i) to (v) in the following passage. The words from the list are to be used |
| | only once. Write the answers as (a) (i), (ii), (iii) and so on. Do not copy the |
| | passage. |

[ammonia, ammonium, carbonate, carbon dioxide, hydrogen, hydronium, hydroxide, precipitate, salt water]:

- (ii) When solutions **M** and **O** are mixed together, the products will be (iii) and (iv)
- (iii) If a piece of magnesium was put into a solution M, (v) gas would be evolved. [5]
- **(b)** Identify the gas evolved in the following reactions when:
 - (i) Sodium propionate is heated with soda lime.
 - (ii) Potassium suphite is treated with dilute hydrochloric acid.
 - (iii) Sulphur is treated with concentrated nitric acid.
 - (iv) A few crystals of KNO3 are heated in a hard glass test tube.
 - (v) Concentrated hydrochloric acid is made to react with manganese dioxide. [5]
- (c) State one appropriate observation for each of the following:
 - (i) Concentrated sulphuric acid is added drop wise to a crystal of hydrated copper sulphate.
 - (ii) Copper sulphide is treated with dilute hydrochloric acid.
 - (iii) Excess of chlorine gas is reacted with ammonia gas.
 - (iv) A few drops of dilute hydrochloric acid are added to silver nitrate solution, followed by addition of ammonium hydroxide solution.
 - (v) Electricity is passed through molten lead bromide. [5]
- (d) Give suitable chemical terms for the following: [5]
 - (i) A bond formed by a shared pair of electrons with both electrons coming from the same atom.
 - (ii) A salt formed by incomplete neutralisation of an acid by a base.
 - (iii) A reaction in which hydrogen of an alkane is replaced by a halogen.
 - (iv) A definite number of water molecules bound to some salts.

| | | (v) | The process in which a substan | ce absorbs moisture from the atmospher | ic | | | |
|------|--|---------|---|--|-------|--|--|--|
| 20 | (e) | Gin | an to become moist, and uttime | itely dissolves in the absorbed water. | | | | |
| | (0) | au | e a chemicai test to distinguish be | tween the following pairs of compounds | : | | | |
| | , | (i) | Sodium Alasi I a la como | [5 | 6] | | | |
| | | 67 1570 | Sodium chloride solution and s | | | | | |
| | | (ii) | Hydrogen chloride gas and hyd | rogen sulphide gas. | | | | |
| | | | ii) Ethene gas and ethane gas. | | | | | |
| | | (iv) | the second die control and the | nc nitrate solution. | | | | |
| | . | | (v) Carbon dioxide gas and sulphur dioxide gas. | | | | | |
| | (f) | Cho | oose the most appropriate answer p | from the following options: [10 |)] | | | |
| | | (i) | Among the period 2 elements affinity is: | , the element which has high electro | n | | | |
| | | 2 | (A) Lithium | (B) Carbon | | | | |
| ¥ | | | (C) Chlorine | (D) Fluorine | | | | |
| | | (ii) | Among the following compour | ds identify the compound that has al | 77 | | | |
| | | | three bonds (ionic, covalent and | coordinate hond) | i | | | |
| | | | (A) Ammonia | (B) Ammonium chloride | | | | |
| | | | (C) Sodium hydroxide | (D) Calcium chloride | | | | |
| | | (iii) | Identify the statement that is in | Correct about albanes | | | | |
| 1= | | | (A) They are hydrocarbons. | correct about atkanes. | | | | |
| | | | | ond between carbon and hydrogen. | | | | |
| | | 4. | (C) They can undergo both sub- | stitution as well as addition reactions. | | | | |
| | | | (D) On complete combustion the | ey produce carbon dioxide and water. | 120 | | | |
| | | (iv) | Which of these will act as a non- | electrolyte? | | | | |
| | | | (A) Liquid carbon tetrachloride | oreon object. | ٠ | | | |
| | | | (B) Acetic acid | * | | | | |
| | , | | (C) Sodium hydroxide aqueous | solution soid | | | | |
| . 67 | | | (D) Potassium chloride aqueous | solution acta | | | | |
| | ş | (v) | Which one of the following will | not made and the same of the s | | | | |
| | | (.,) | with water ? | not produce an acid when made to react | react | | | |
| | | | (A) Carbon monoxide | (B) Carbon dioxide | | | | |
| | | 100 | (C) Nitrogen dioxide | (D) Sulphur trioxide | • | | | |
| | | (vi) | Identify the metallic oxide which | is amphoteric in nature: | | | | |
| | A | | (A) Calcium oxide | (B) Barium oxide | | | | |
| | | | (C) Zinc oxide | (D) Copper(II) oxide | 4 | | | |
| | | (vii) | In the given equation identify th | e role played by concentrated sulphuric | | | | |
| * | $acid\ S + 2H_2SO_4 \longrightarrow 3SO_2 + 2H_2O$: | | | | | | | |
| | 100 | U U | (A) Non-volatile acid | (B) Oxidising agent | | | | |
| | | | (C) Dehydrating agent | (D) None of the above | | | | |
| | | (viii) | Nitrogen gas can be obtained by | | | | | |
| | | | (A) Ammonium nitrate | (B) Ammonium nitrite | | | | |
| | | 8 3 C | (C) Magnesium nitride | (D) Ammonium chloride | | | | |
| | | (ix) | | pical property of an ionic compound? | | | | |
| | | | (A) High melting point. | erous property of an ionic compound? | | | | |

| | | (B) Conducts electricity in the molten and in the aqueous solution state. |
|------------|---------|---|
| | | (C) They are insoluble in water. |
| | | (D) They exist as oppositely charged ions even in the solid state. |
| | (x) | The metals zinc and tin are present in the alloy: |
| | Č | (A) Solder (B) Brass |
| | | (C) Bronze (D) Duralumin |
| (g) | Solve | e the following: |
| | (i) | What volume of oxygen is required to burn completely 90 dm ³ of butane under similar conditions of temperature and pressure? [2] |
| | | under stitutal conditions of temperature F. |
| | ··· | $2C_4H_{10} + 13O_2 \longrightarrow 8CO_2 + 10H_2O$ The vapour density of a gas is 8. What would be the volume occupied by |
| | (ii) | 24.0g of the gas at STP? [2] |
| | (iii) | A vessel contains X number of molecules of hydrogen gas at a certain |
| | (111) | temperature and pressure. How many molecules of nitrogen gas would be |
| R | | present in the same vessel under the same conditions of temperature and |
| | | pressure? [1] |
| Ans | wer. | |
| (a) | (i) | Hydronium (ii) Hydroxide (iii) Salt |
| | (iv) | Water (v) Hydrogen |
| (b) | (i) | Ethane (C ₂ H ₆) (ii) Sulphur dioxide (SO ₂) (iii) Nitrogen dioxide (NO ₂) |
| | (iv) | Oxygen (O_2) (v) Chlorine (Cl_2) |
| (c) | (i) | The blue colour of the solution changes to white. |
| | (ii) | A gas evolved which has the smell of rotten eggs. |
| | (iii) | A yellow coloured explosive is formed. |
| | (iv) | Curdy white ppt. observed which is soluble in NH ₄ OH. |
| | (v) | Reddish brown vapours of bromine are evolved. |
| (d) | (i) | Co-ordinate bond (ii) Acidic salt (iii) Halogenation Water of crystallisation (v) Deliquescence |
| | (iv) | Water of Crystallication |
| (e) | (i). | Sodium chloride gives white ppt. with silver nitrate solution while |
| | | sodium nitrate does not. Dense white fumes observed when a rod dipped in NH ₃ is brought to the |
| | (ii) | mouth of t.t containing HCl gas. Whereas no such fumes observed in |
| | | case of H_2S gas. |
| | (iii) | - Land Land Land Land Land Land Land Land |
| | (111) | observed with ethane. |
| | (iv) | the second with |
| | | Zn(NO ₃) ₂ solution which is soluble in excess. Whereas white ppt. |
| | | observed with $Ca(NO_3)_2$ which is sparingly soluble in water. |
| | (v) | SO_2 turns acidified potassium dichromate solution green whereas CO_2 |
| | | does not change the colour. |
| (f) | (i) | $D \qquad \text{(ii)} B \qquad \text{(iii)} C \qquad \text{(iv)} A \qquad \text{(v)} A$ |
| | (vi) | |
| (g) | (i) | $2C_4H_{10} + 13O_2 \longrightarrow 8CO_2 + 10H_2O$ |
| | | 2 Volume of butane = 90 dm^3 |
| | | |

1 Volume of butane = 45 dm^3 Volume of O_2 required = 13×45 $= 585 \, dm^3$ Ans. (ii) Molecular Weight = $2 \times VD$ = 16gVolume of 16g gas = 22.4LVolume of 1g gas = Volume of 24g gas = $\frac{22.4}{16} \times 24$ = 33.6 litres. Ans. It will contain X no. of molecules. SECTION-II (40 Marks) (Answer any four questions from this section) Question 2. Group IA IIA IIIA IVA VA VIA VIIA 0 number 1 2 13 14 *15* 16 17 18 2^{nd} period Li \boldsymbol{D} \boldsymbol{J} 0 Ne \boldsymbol{A} Mg \boldsymbol{E} SiHM \boldsymbol{R} TQ \boldsymbol{u} y In this table H does not represent hydrogen. Some elements are given in their own symbol and position in the periodic table. While others are shown with a letter. With reference to the table answer the following questions: (i) Identify the most electronegative element. [1] (ii) Identify the most reactive element of group 1. [1] Identify the element from perod 3 with least atomic size. (iii) [1] How many valence electrons are present in Q? (iv) [1] (v) Which element from group 2 would have the least ionization energy? [1] Identify the noble gas of the fourth period. [1] (vii) In the compound between A and B what type of bond would be formed and give the molecular formula for the same. [2] Compare the compounds carbon tetrachloride and sodium chloride with regard to solubility in water and electrical conductivity. [2]Answer: (i) J (ii) R (iii) M (iv) 5 electron (v) T (vi) y (vii) Ionic Bond. Α

Thus, Molecular formula is A_2H .

+1

 A_2H

(a)

(a)

Carbon tetrachloride is insoluble in water and behave as a bad conductor of **(b)** electricity while sodium chloride is soluble in water and behave as a good conductor of electricity in their aqueous state.

[4]

Question 3.

Choosing the substances from the list given below, write balanced chemical equations for the reactions which would be used in the laboratory to obtain the following salts:

Copper(II) carbonate Dilute Sulphuric acid Copper Sodium carbonate Iron Sodium chloride Sodium Zinc nitrate

(i) Sodium sulphate

Zinc carbonate (ii)

Copper(II) sulphate (iii)

Iron(II) sulphate. (iv)

State two relevant observations for each of the following:

Ammonium hydroxide solution is added to copper(II) nitrate solution in small quantities and then in excess.

Ammonium hydroxide solution is added to zinc nitrate solution in (ii) minimum quantities and then in excess.

Lead nitrate crystals are heated in a hard glass test tube. [6] (iii)

Answer.

(b)

 $2NaCl + H_2SO_4(dil.) \longrightarrow Na_2SO_4 +$ (i) (a) Sod. Sulphate Hydrogen chloride Sodium chloride Sulphuric acid

+ 2NaNO₃ $Na_2CO_3 \longrightarrow ZnCO_3$ $Zn(NO_3)_2$ + (ii) Sodium Carbonate Zinc Carbonate **Sodium Nitrate** Zinc Nitrate

 $CuCO_3 + H_2SO_4 \, (dil.) \,\, \longrightarrow \,\, CuSO_4 + H_2O + CO_2$ (iii) Copper Carbonate Sulphuric Acid Copper Sulphate

 $Fe + H_2SO_4 (dil.) \longrightarrow FeSO_4$ (iv) Iron (II) Sulphate Sulphuric Acid

(a) When NH₄OH is added to copper (II) nitrate solution in small quan-**(b)** (i) tities then a pale blue ppt. is observed.

(b) When added in excess it disolves to give an inky blue solution forming a complex salt.

(ii) (a) When added in small quantities is forms a gelatinous white ppt.


(b) When added in excess it dissolves to form a complex salt.

(iii) (a) Reddish brown gas is evolved.

(b) Colourless, odourless gas evolved which rekindles a glowing splinter.

Question 4.

Copper sulphate solution is electrolysed using copper electrodes. Study the diagram given below and answer the question that follows:



Copper (II) Sulphate Solution

- Which electrode to your left or right is knwon as the oxidising electrode (i) and why? [2]
- Write the equation representing the reaction that occurs. (ii)

[1]

State two appropriate observations for the above electrolysis reaction. [2] (iii)

| * | X | Y |
|---|---|--|
| Normal Electronic Configuration | 2, 8, 7 | 2, 8, 2 |
| Nature of oxide | Dissolves in water and turns blue litmus red | Very low solubility water. Dissolves hydrochloric acid |
| Tendency for oxidising and reducing reactions Electrical and Thermal conductivity | Tends to oxidise ele- ments and compounds Very poor electrical conductor | Tends to act as reducing agent |
| Tendency to form | Poor thermal conductivity No tendency to form alloys | Good Thermal conductor Forms alloys |

Using the information above, complete the following:

- (i) is the metallic element.
- (ii) Metal atoms tend to have a maximum of electrons in the outermost energy level.
- Non-metallic elements tend to form oxides while metals tend to (iii) form oxides.
- Non-metallic elements tend to be conductors of heat and (iv) electricity.
- Metals tend to electrons and act as agents in their reactions with elements and compounds. [5]

Answer:

- (a) Right electrode is known as oxidising electrode because copper ion gain (i) electron to form copper metal i.e., reduction takes place.
 - (ii) At Catode:

 $Cu^{2+} + 2e^{-} \longrightarrow Cu$

At Anode:

 $Cu - 2e^- \longrightarrow Cu^{2+}$

- (a) Reddish brown metal copper is deposited at cathode. (iii)
 - (b) Blue colour of electrolytic solution i.e., CuSO₄ does not fades during the process.
- **(b)** (i) Y
- (ii) 2
- (iii) Acidic, Basic
- (iv) Bad

(v) lose, reducing

Question 5.

- Give balanced equations for each of the following:
 - Reduction of hot Copper(II) oxide to copper using ammonia gas. (i)
 - Oxidation of carbon with concentrated nitric acid. (ii)
 - Dehydration of concentrated sulphuric acid with sugar crystals.

Copy and complete the following table relating to important industrial **(b)** [3] process: Equation for the Catalyst Name of the **Temperature** catalyzed reaction process Haber's process The following questions relate to the extraction of aluminium by electrolysis: (c) Name the other aluminium containing compound added to alumina and state its significance. Give the equation for the reaction that takes place at the cathode. Explain why is it necessary to renew the anode periodically. **[4]** (iii) Answer: $3\text{CuO} + 2\text{NH}_3 \longrightarrow 3\text{Cu} + \text{N}_2 \uparrow + 3\text{H}_2\text{O}$ (a) (i) (ii) $C + 4HNO_3 \longrightarrow CO_2 + 2H_2O + 4NO_2$ $\xrightarrow{\text{Conc. H}_2\text{SO}_4} 12\text{C} + 11\text{H}_2\text{O}$ Sugar Temperature: 450-500 °C **(b)** Catalyst: Finely divided iron. **Equation:** $N_2(g) + 3H_2(g)$ $2NH_3(l)$ (i) Cryolite (Na_3AlF_6) (c) It reduces the temperature and enhance conductivity. At Cathode: $Al^{3+} + 3e^{-} \longrightarrow Al$ (iii) The anode has got to be replaced periodically, as it gets oxidised by the oxygen evolved at the anode. Question 6. Give balanced equations for the laboratory preparations of the following organic compounds: A saturated hydrocarbon from iodomethane. (i) An unsaturated hydrocarbon from an alcohol. (ii) (iii) An unsaturated hydrocarbon from calcium carbide. [4] (iv) An alcohol from ethyl bromide. Give the structural formulae for the following: **(b)** An isomer of n-butane. (i) (ii) 2-propanol. [3] (iii) Diethyl ether. Give reasons for the following: **(c)** Methane does not undergo addition reactions, but ethene does. (i) (ii) Ethyne is more reactive than ethane. [3] Hydrocarbons are excellent fuels. (iii) Answer: $CH_3I + 2[H] \longrightarrow CH_4 + HI$ (a) (i)


Iodomethane

C₂H₅OH Alcohol

(ii)

 $\xrightarrow{\text{Conc. } H_2SO_4} C_2H_4 + H_2O$

- $\begin{array}{ccc} \text{(iii)} & CaC_2 + 2H_2O & \longrightarrow & C_2H_2 + Ca(OH)_2 \\ & & \text{Calcium carbide} \end{array}$
- (iv) $C_2H_5Br + KOH(aq) \longrightarrow C_2H_5OH + KBr$ Ethyl bromide
- **(b)** (i) Isomer of n-butane:



(ii) 2-Propanol:

(iii) Diethyl ether:

- (c) (i) Methane does not undergo addition reaction, because it is bonded with four H-atom while in ethene double bond breaks and provide site for addition.
 - (ii) Due to the presence of triple bond it provide site for addition, hence ethyne is more reactive than ethane.
 - (iii) Hydrocarbon are excellent fuels as they produces lot of heat during combustion.

Question 7.

(a) O_2 is evolved by heating $KClO_3$ using MnO_2 as a catalyst

$$2KClO_3 \xrightarrow{MnO_2} 2KCl + 3O_2$$

- (i) Calculate the mass of $KClO_3$ required to produce 6.72 litre of O_2 at STP. [atomic masses of K = 39, Cl = 35.5, O = 16]
- (ii) Calculate the number of moles of oxygen present in the above volume and also the number of molecules. [2]
- (iii) Calculate the volume occupied by 0.01 mole of CO₂ at STP. [1]
- (b) Identify the following substances which are underlined:
 - (i) An alkaline gas which produces dense white fumes when reacted with hydrogen chloride gas.
 - (ii) An acid which is present in vinegar.
 - (iii) <u>A gas</u> which does not conduct electricity in the liquid state but conducts electricity when dissolved in water.

barium chloride solution. The element which has the highest ionization potential. [5] (v) Answer. $\xrightarrow{\rm MnO_2} ~ 2 \rm KCl$ 30_{2} 2KClO₃ -(i) (a) $2 \times (39 + 35.5 + 3 \times 16) \ 2 (39 + 35.5)$ $= 2 \times 74.5 = 149g$ $3 \times 22.4 = 67.2l$ $=2\times122\cdot5=245g$ 67.2l of O_2 produced by $KClO_3 = 245 g$ $1l \text{ of } O_2 \text{ produced by } KClO_3 = \frac{245}{67 \cdot 2}$ 6.72 of O_2 produced by $KClO_3$ = Ans. = 24.5 g245g of KClO₃ contains = 3 moles of oxygen (ii) 24.5g of KClO₃ contains = Ans. = 0.3 mole. 1 mole of oxygen contain = 6.022×10^{23} molecules Since, 0.3 mole of oxygen contain = $6.022 \times 10^{23} \times 0.3$ Therefore, = 1.8066×10^{23} molecules 1 mole of CO_2 occupied = 22.4 l(iii) $0.01 \text{ mole of CO}_2 \text{ occupied} = 22.4 \times 0.01$ = 0.224 l(ii) Acetic Acid (CH₃COOH) Ammonia gas (NH₃) **(b)** (i) (iv) Sulphuric Acid (H₂SO₄) Hydrogen chloride (HCl)

A dilute mineral acid which forms a white precipitate when treated with

(iv)

(iii)

(v)

Helium (He)