

# ICSE Paper 2008

## CHEMISTRY

### SECTION—I (40 Marks)

(Compulsory : Attempt all questions.)


#### Question 1.

- (a) For part (a) (i) – (a) (x), select the correct answer from the choices A, B, C, D which are given. Write down only the letter corresponding to the correct answer.
- (i) With reference to the variation of properties in the Periodic Table, which of the following is generally true ?
- (A) Atomic size increases from left to right across a period.  
(B) Ionization potential increases from left to right across a period.  
(C) Electron affinity increases going down a group.  
(D) Electro-negativity increases going down a group.
- (ii) Which of the following is **not** a common characteristic of an electrovalent compound ?
- (A) High melting point.  
(B) Conducts electricity when molten.  
(C) Consists of oppositely charged ions.  
(D) Ionizes when dissolved in water.
- (iii) Dilute sulphuric acid will produce a white precipitate when added to a solution of :
- (A) Copper nitrate (B) Zinc nitrate  
(C) Lead nitrate (D) Sodium nitrate
- (iv) The salt which in solution gives a pale green precipitate with sodium hydroxide solution and a white precipitate with barium chloride solution is :
- (A) Iron (III) sulphate (B) Iron (II) sulphate  
(C) Iron (II) chloride (D) Iron (III) chloride
- (v) The gas law which relates the volume of a gas to the number of molecules of the gas is :
- (A) Avogadro's Law (B) Gay-Lussac's Law  
(C) Boyle's Law (D) Charles' Law
- (vi) During the electrolysis of molten lead bromide, which of the following takes place ?
- (A) Bromine is released at the cathode  
(B) Lead is deposited at the anode  
(C) Bromine ions gain electrons  
(D) Lead is deposited at the cathode

- (vii) Brass is an alloy of :
- (A) Copper and tin (B) Copper and zinc  
(C) Zinc and lead (D) Lead and tin
- (viii) Ammonia can be obtained by adding water to :
- (A) Ammonium chloride (B) Ammonium nitrate  
(C) Magnesium nitrate (D) Magnesium nitrate
- (ix) Which of the following reactions is used to prepare sulphuryl chloride ?\*\*
- (A) Adding concentrated sulphuric acid to a chloride.  
(B) Passing sulphur dioxide through a solution of chlorine.  
(C) Reacting dry sulphur dioxide and dry chlorine.  
(D) Reacting dilute sulphuric acid with a solution of chlorine.
- (x) The formation of 1, 2-dibromethane from ethene and bromine is an example of :
- (A) Substitution (B) Dehydration  
(C) Dehydrohalogenation (D) Addition

[10]

(b) The equation for the burning of octane is :



- (i) How many moles of carbon dioxide are produced when **one** mole of octane burns ?
- (ii) What volume, at stp, is occupied by the number of moles determined in (b) (i) ?
- (iii) If the relative molecular mass of carbon dioxide is 44, what is the mass of carbon dioxide produced by burning **two** moles of octane ?
- (iv) What is the empirical formula of octane ?

[5]

(c) **Name** the organic compound prepared by each of the following reactions :

- (i)  $C_2H_5COONa + NaOH \longrightarrow$
- (ii)  $CH_3I + 2\bar{H} \longrightarrow$
- (iii)  $C_2H_5Br + KOH$  (alcoholic solution)  $\longrightarrow$
- (iv)  $CO + 2H_2$  (Zinc oxide catalyst)  $\longrightarrow$
- (v)  $CaC_2 + 2H_2O \longrightarrow$

[5]




(d) Identify the following substances :

- (i) An alkaline gas A which gives dense white fumes with hydrogen chloride.
- (ii) A dilute acid B which does not normally give hydrogen when reacted with metals but does give a gas when it reacts with copper.
- (iii) Gas C has an offensive smell like rotten eggs.
- (iv) Gas D is a colourless gas which can be used as a bleaching agent. \*\*
- (v) Liquid E can be dehydrated to produce ethane.

[5]

\*\* Answer has not given due to out of present syllabus.



(f) (i) This reaction will take place at anode.  
It is an example of oxidation.(ii)  $Cu^{++}$  ions.

(iii) Carbon tetrachloride is a covalent compound.

(g) (i) Covalent bond (ii) Co-ordinate bond.

(iv) Alkali

(v) Substitution.

(iii) Complex salt.

**Section—II (40 marks)**

(Answer any four questions from this section)

**Question 2.**Copy and complete the following table relating to important industrial processes. Output refers to the product of the process **not** the intermediate steps.

Name of process	Inputs	Catalyst	Equation for catalysed reaction	Output
Haber Process	Hydrogen + Ammonia + air			
Contact Process	Sulphur dioxide + oxygen			Nitric acid

**Answer.**

Name of process	Inputs	Catalyst	Equation for catalysed reaction	Output
Haber Process	Hydrogen + Nitrogen	Finely dioxided iron	$N_2 + 3H_2 \longrightarrow 2NH_3$	Ammonia
Ostwald's Process	Ammonia + air	Platinum	$4NH_3 + 5O_2 \longrightarrow 4NO + 6H_2O$	Nitric acid
Contact Process	Sulphur dioxide + oxygen	Vanadium pentoxide	$2SO_2 + O_2 \longrightarrow 2SO_3$	Sulphuric acid

**Question 3.**

The following questions refer to the Periodic Table :

- (a) (i) Name the first and last element in period 2.  
(ii) What happens to the atomic size of elements moving from top to bottom of a group?



- (iii) Which of the elements has the greatest electron affinity among the halogens ?
- (iv) What is the common feature of the electronic configurations of the elements in group 7 ? [5]
- (b) Supply the missing word from those in the brackets (Do not write out the sentence).
- (i) If an element has a low ionization energy then it is likely to be ..... (metallic/non-metallic).
- (ii) If an element has seven electrons in its outermost shell then it is likely to have the ..... (largest/smallest) atomic size among all the elements in the same period. [2]
- (c) (i) The metals of Group 2 from top to bottom are : Be, Mg, Ca, Sr, Ba. Which of these metals will form ions most readily and why ?
- (ii) What property of an element is measured by electronegativity ? [3]

**Answer.**

- (a) (i) First element → Lithium.  
Last element → Neon
- (ii) It increases.
- (iii) Chlorine.
- (iv) Elements of seventh group have seven electrons in their valence shell.
- (b) (i) metallic (ii) smallest.
- (c) (i) Ba will form ions most readily because it has lowest ionisation potential in group 2.
- (ii) Electronegativity of an element is its relative tendency to attract the shared pair of electrons towards itself in a covalent bond.

**Question 4.**

- (a) Distinguish between the saturated hydrocarbon ethane and the unsaturated hydrocarbon ethene by drawing their structural formulae. [2]
- (b) Addition reactions and substitution reactions are types of organic reactions. Which type of reaction is shown by :
- (i) ethane (ii) ethene. [2]
- (c) (i) Write the equation for the complete combustion of ethane.
- (ii) Using appropriate catalysts, ethane can be oxidized to an alcohol, an aldehyde and an acid. Name the alcohol, aldehyde and acid formed when ethane is oxidized. [4]
- (d) (i) Why is pure acetic acid known as glacial acetic acid ?
- (ii) What type of compound is formed by the reaction between acetic acid and an alcohol ? [2]

**Answer.**

**Saturated hydrocarbon :** Ethane has single covalent bond between carbon atoms.


**Unsaturated hydrocarbon :** Ethene has a double covalent bond between carbon atoms.

- (b) (i) ethane  $\longrightarrow$  substitution reactions.  
 (ii) ethene  $\longrightarrow$  addition reactions.
- (c) (i)  $2C_2H_6 + 7O_2 \longrightarrow 4CO_2 + 6H_2O$   
 (ii) When ethene is oxidised, ethanol ( $C_2H_5OH$ ), ethanal ( $CH_3CHO$ ) and ethanoic acid ( $CH_3COOH$ ) are formed.
- (d) (i) Because on cooling pure acetic acid forms ice like crystals.  
 (ii) An ester is formed.

**Question 5.**

- (a) (i) A compound has the following percentage composition by mass : carbon 14.4%, hydrogen 1.2% and chlorine 84.5%. Determine the empirical formula of this compound. Work correct to 1 decimal place. ( $H = 1$ ;  $C = 12$ ;  $Cl = 35.5$ ).  
 (ii) The relative molecular mass of this compound is 168, so what is its molecular formula ?  
 (iii) By what type of reaction could this compound be obtained from ethyne ?

(b) From the equation



Calculate :

- (i) The mass of carbon oxidized by 49 g of sulphuric acid ( $C = 12$ ; relative molecular mass of sulphuric acid = 98).  
 (ii) The volume of sulphur dioxide measured at stp. liberated at the same time.

Volume occupied by 1 mole of a gas at stp is  $22.4 \text{ dm}^3$ .

**Answer.**

(a) (i)

Element	Percentage	At. Mass	Relative No. of atoms	Simplest ratio
C	14.4	12	$14.4/12 = 1.2$	$1.2/1.2 = 1$
H	1.2	1	$1.2/1 = 1.2$	$1.2/1.2 = 1$
Cl	84.5	35.5	$84.5/35.5 = 2.4$	$2.4/1.2 = 2$

Empirical formula =  $CHCl_2$

(ii)

Relative molecular mass = 168

Empirical formula mass =  $12 + 1 + 71$

= 84

$n = \frac{\text{Relative molecular mass}}{\text{empirical formula mass}}$

=  $\frac{168}{84} = 2$

[4]

[6]

$$\begin{aligned} \text{Molecular formula} &= (\text{Empirical formula})_n \\ &= (\text{CHCl}_2)_2 \\ &= \text{C}_2\text{H}_2\text{Cl}_4. \end{aligned}$$

(iii) By addition reaction.

(b) (i) From equation :

$2 \times 98$  g of sulphuric acid oxidises 12g of carbon

$$\begin{aligned} \therefore 49\text{g of sulphuric acid will oxidise} &= \frac{12 \times 49}{2 \times 98} \\ &= 3\text{g of C.} \end{aligned}$$

**Ans.**

(ii) From equation :


$2 \times 98$  of sulphuric acid liberates  $2 \times 22.4$  l of  $\text{SO}_2$ .

$$\begin{aligned} \therefore 49\text{ g of sulphuric acid will liberate} &= \frac{2 \times 22.4 \times 49}{2 \times 98} \\ &= 11.2\text{ l of SO}_2. \end{aligned}$$

**Ans.**

### Question 6.

(a) The following is a sketch of an electrolytic cell used in the extraction of aluminium :



(i) What is the substance of which the electrodes A and B are made ?

(ii) At which electrode (A or B) is the aluminium formed ?

(iii) What are the two aluminium compounds in the electrolyte C ?

(iv) Why is it necessary for electrode B to be continuously replaced ? [5]

(b) Making use only of substances chosen from those given below :

Dilute sulphuric acid

sodium carbonate

Zinc

sodium sulphite

Lead

calcium carbonate

give the equations for the reactions by which you could obtain :

(i) hydrogen

(ii) sulphur dioxide\*\*

(iii) carbon dioxide

(iv) zinc carbonate (two steps required) [5]

### Answer.

(a) (i) Graphite or carbon.

(ii) At electrode A.



(iii) Alumina (Aluminium oxide) and Cryolite (sodium aluminium fluoride).

(iv) Because electrode B gets burnt away with oxygen produced.

(b) (i)  $\text{Zn} + \text{H}_2\text{SO}_4 \longrightarrow \text{ZnSO}_4 + \text{H}_2$

(iii)  $\text{Na}_2\text{CO}_3 + \text{H}_2\text{SO}_4 \longrightarrow \text{Na}_2\text{SO}_4 + \text{H}_2\text{O} + \text{CO}_2$ .

\*\* Answer has not given due to out of present syllabus.

**Question 7.**

- (a) (i) *What is the property of concentrated sulphuric acid which allows it to be used in the preparation of hydrogen chloride and nitric acid ?*
- (ii) *What property of hydrogen chloride is demonstrated when it is collected by downward delivery (upward displacement) ?*
- (iii) *Why is hydrogen chloride not collected over water ?*
- (iv) *What is the property of nitric acid which allows it to react with copper ?*
- (v) *What property of concentrated sulphuric acid is in action when sugar turns black in its presence ?* [5]

(b) *Write the equations for the following reactions :*

- (i) *Dilute nitric acid and copper.*
- (ii) *Dilute sulphuric acid and barium chloride.*
- (iii) *Dilute hydrochloric acid and sodium thiosulphate.*
- (iv) *Dilute hydrochloric acid and lead nitrate solution.*
- (v) *Dilute sulphuric acid and sodium sulphide.* [5]

**Answer.**

- (a) (i) It is non-volatile.
- (ii) Hydrogen chloride is heavier than air.
- (iii) Because Hydrogen chloride is highly soluble in water.
- (iv) It is a strong oxidising agent.
- (v) Conc. sulphuric acid is a dehydrating agent.
- (b) (i)  $3\text{Cu} + 8\text{HNO}_3 \longrightarrow 3\text{Cu}(\text{NO}_3)_2 + 2\text{NO} + 4\text{H}_2\text{O}$
- (ii)  $\text{H}_2\text{SO}_4 + \text{BaCl}_2 \longrightarrow \text{BaSO}_4 + 2\text{HCl}$
- (iii)  $\text{Na}_2\text{S}_2\text{O}_3 + 2\text{HCl} \longrightarrow 2\text{NaCl} + \text{H}_2\text{O} + \text{SO}_2 + \text{S}$
- (iv)  $\text{Pb}(\text{NO}_3)_2 + 2\text{HCl} \longrightarrow \text{PbCl}_2 + 2\text{HNO}_3$
- (v)  $\text{Na}_2\text{S} + \text{H}_2\text{SO}_4 \longrightarrow \text{Na}_2\text{SO}_4 + \text{H}_2\text{S}$ .