

# Binomial Theorem

## Short Answer Type Questions

**Q. 1** Find the term independent of  $x$ , where  $x \neq 0$ ,

in the expansion of  $\left(\frac{3x^2}{2} - \frac{1}{3x}\right)^{15}$ .

### Thinking Process

The general term in the expansion of  $(x-a)^n$  i.e.,  $T_{r+1} = {}^nC_r(x)^{n-r}(-a)^r$ . For the term independent of  $x$ , put  $n-r=0$ , then we get the value of  $r$ .

**Sol.** Given expansion is  $\left(\frac{3x^2}{2} - \frac{1}{3x}\right)^{15}$ .

Let  $T_{r+1}$  term is the general term.

$$\begin{aligned} \text{Then, } T_{r+1} &= {}^{15}C_r \left(\frac{3x^2}{2}\right)^{15-r} \left(-\frac{1}{3x}\right)^r \\ &= {}^{15}C_r 3^{15-r} x^{30-2r} 2^{r-15} (-1)^r \cdot 3^{-r} \cdot x^{-r} \\ &= {}^{15}C_r (-1)^r 3^{15-2r} 2^{r-15} x^{30-3r} \end{aligned}$$

For independent of  $x$ ,

$$\begin{aligned} 30 - 3r &= 0 \\ 3r &= 30 \Rightarrow r = 10 \end{aligned}$$

$\therefore T_{r+1} = T_{10+1} = 11\text{th term is independent of } x.$

$$\begin{aligned} \therefore T_{10+1} &= {}^{15}C_{10} (-1)^{10} 3^{15-20} 2^{10-15} \\ &= {}^{15}C_{10} 3^{-5} 2^{-5} \\ &= {}^{15}C_{10} (6)^{-5} \\ &= {}^{15}C_{10} \left(\frac{1}{6}\right)^5 \end{aligned}$$

**Q. 2** If the term free from  $x$  in the expansion of  $\left(\sqrt{x} - \frac{k}{x^2}\right)^{10}$  is 405, then find the value of  $k$ .

**Sol.** Given expansion is  $\left(\sqrt{x} - \frac{k}{x^2}\right)^{10}$ .

Let  $T_{r+1}$  is the general term.

$$\begin{aligned} \text{Then, } T_{r+1} &= {}^{10}C_r (\sqrt{x})^{10-r} \left(\frac{-k}{x^2}\right)^r \\ &= {}^{10}C_r (x)^{\frac{1}{2}(10-r)} (-k)^r \cdot x^{-2r} \\ &= {}^{10}C_r x^{5-\frac{r}{2}} (-k)^r \cdot x^{-2r} \\ &= {}^{10}C_r x^{5-\frac{r}{2}-2r} (-k)^r \\ &= {}^{10}C_r x^{\frac{10-5r}{2}} (-k)^r \end{aligned}$$

For free from  $x$ ,  $\frac{10-5r}{2} = 0$

$\Rightarrow 10-5r = 0 \Rightarrow r = 2$

Since,  $T_{2+1} = T_3$  is free from  $x$ .

$\therefore T_{2+1} = {}^{10}C_2 (-k)^2 = 405$

$\Rightarrow \frac{10 \times 9 \times 8!}{2! \times 8!} (-k)^2 = 405$

$\Rightarrow 45k^2 = 405 \Rightarrow k^2 = \frac{405}{45} = 9$

$\therefore k = \pm 3$

**Q. 3** Find the coefficient of  $x$  in the expansion of  $(1 - 3x + 7x^2)(1 - x)^{16}$ .

**Sol.** Given, expansion =  $(1 - 3x + 7x^2)(1 - x)^{16}$ .

$$\begin{aligned} &= (1 - 3x + 7x^2)({}^{16}C_0 1^{16} - {}^{16}C_1 1^{15} x^1 + {}^{16}C_2 1^{14} x^2 + \dots + {}^{16}C_{16} x^{16}) \\ &= (1 - 3x + 7x^2)(1 - 16x + 120x^2 + \dots) \end{aligned}$$

$\therefore$  Coefficient of  $x = -3 - 16 = -19$

**Q. 4** Find the term independent of  $x$  in the expansion of  $\left(3x - \frac{2}{x^2}\right)^{15}$ .

💡 **Thinking Process**

The general term in the expansion of  $(x-a)^n$  i.e.,  $T_{r+1} = {}^nC_r (x)^{n-r} (-a)^r$ .

**Sol.** Given expansion is  $\left(3x - \frac{2}{x^2}\right)^{15}$ .

Let  $T_{r+1}$  is the general term.

$$\begin{aligned} \therefore T_{r+1} &= {}^{15}C_r (3x)^{15-r} \left(\frac{-2}{x^2}\right)^r = {}^{15}C_r (3x)^{15-r} (-2)^r x^{-2r} \\ &= {}^{15}C_r 3^{15-r} x^{15-3r} (-2)^r \end{aligned}$$

For independent of  $x$ ,  $15 - 3r = 0 \Rightarrow r = 5$

Since,  $T_{5+1} = T_6$  is independent of  $x$ .

$$\begin{aligned} \therefore T_{5+1} &= {}^{15}C_5 \cdot 3^{15-5} \cdot (-2)^5 \\ &= -\frac{15 \times 14 \times 13 \times 12 \times 11 \times 10!}{5 \times 4 \times 3 \times 2 \times 1 \times 10!} \cdot 3^{10} \cdot 2^5 \\ &= -3003 \cdot 3^{10} \cdot 2^5 \end{aligned}$$

**Q. 5** Find the middle term (terms) in the expansion of

(i)  $\left(\frac{x}{a} - \frac{a}{x}\right)^{10}$

(ii)  $\left(3x - \frac{x^3}{6}\right)^9$

🔑 **Thinking Process**

In the expansion of  $(a + b)^n$ , if  $n$  is even, then this expansion has only one middle term i.e.,  $\left(\frac{n}{2} + 1\right)$ th term is the middle term and if  $n$  is odd, then this expansion has two middle terms i.e.,  $\left(\frac{n+1}{2}\right)$ th and  $\left(\frac{n+1}{2} + 1\right)$ th are two middle terms.

**Sol.** (i) Given expansion is  $\left(\frac{x}{a} - \frac{a}{x}\right)^{10}$ .

Here, the power of Binomial i.e.,  $n = 10$  is even.

Since, it has one middle term  $\left(\frac{10}{2} + 1\right)$ th term i.e., 6th term.

$$\begin{aligned} \therefore T_6 = T_{5+1} &= {}^{10}C_5 \left(\frac{x}{a}\right)^{10-5} \left(\frac{-a}{x}\right)^5 \\ &= -{}^{10}C_5 \left(\frac{x}{a}\right)^5 \left(\frac{a}{x}\right)^5 \\ &= -\frac{10 \times 9 \times 8 \times 7 \times 6 \times 5!}{5! \times 5 \times 4 \times 3 \times 2 \times 1} \left(\frac{x}{a}\right)^5 \left(\frac{x}{a}\right)^{-5} \\ &= -9 \times 4 \times 7 = -252 \end{aligned}$$

(ii) Given expansion is  $\left(3x - \frac{x^3}{6}\right)^9$ .

Here,  $n = 9$

Since, the Binomial expansion has two middle terms i.e.,  $\left(\frac{9+1}{2}\right)$ th and  $\left(\frac{9+1}{2} + 1\right)$ th [odd]

i.e., 5th term and 6th term.

$$\begin{aligned} \therefore T_5 = T_{(4+1)} &= {}^9C_4 (3x)^{9-4} \left(-\frac{x^3}{6}\right)^4 \\ &= \frac{9 \times 8 \times 7 \times 6 \times 5!}{4 \times 3 \times 2 \times 1 \times 5!} 3^5 x^5 x^{12} 6^{-4} \\ &= \frac{7 \times 6 \times 3 \times 3^1}{2^4} x^{17} = \frac{189}{8} x^{17} \end{aligned}$$

$$\begin{aligned}
 \therefore T_6 = T_{5+1} &= {}^9C_5(3x)^{9-5} \left(-\frac{x^3}{6}\right)^5 \\
 &= -\frac{9 \times 8 \times 7 \times 6 \times 5!}{5! \times 4 \times 3 \times 2 \times 1} \cdot 3^4 \cdot x^4 \cdot x^{15} \cdot 6^{-5} \\
 &= \frac{-21 \times 6}{3 \times 2^5} x^{19} = \frac{-21}{16} x^{19}
 \end{aligned}$$

**Q. 6** Find the coefficient of  $x^{15}$  in the expansion of  $(x - x^2)^{10}$ .

**Sol.** Given expansion is  $(x - x^2)^{10}$ .

Let the term  $T_{r+1}$  is the general term.

$$\begin{aligned}
 \therefore T_{r+1} &= {}^{10}C_r x^{10-r} (-x^2)^r \\
 &= (-1)^r \cdot {}^{10}C_r \cdot x^{10-r} \cdot x^{2r} \\
 &= (-1)^r {}^{10}C_r x^{10+r}
 \end{aligned}$$

For the coefficient of  $x^{15}$ ,

$$10 + r = 15 \Rightarrow r = 5$$

$$T_{5+1} = (-1)^5 {}^{10}C_5 x^{15}$$

$$\begin{aligned}
 \therefore \text{Coefficient of } x^{15} &= -1 \frac{10 \times 9 \times 8 \times 7 \times 6 \times 5!}{5 \times 4 \times 3 \times 2 \times 1 \times 5!} \\
 &= -3 \times 2 \times 7 \times 6 = -252
 \end{aligned}$$

**Q. 7** Find the coefficient of  $\frac{1}{x^{17}}$  in the expansion of  $\left(x^4 - \frac{1}{x^3}\right)^{15}$ .

**Thinking Process**

In this type of questions, first of all find the general terms, in the expansion  $(x-y)^n$  using the formula  $T_{r+1} = {}^nC_r x^{n-r} (-y)^r$  and then put  $n-r$  equal to the required power of  $x$  of which coefficient is to be find out.

**Sol.** Given expansion is  $\left(x^4 - \frac{1}{x^3}\right)^{15}$ .

Let the term  $T_{r+1}$  contains the coefficient of  $\frac{1}{x^{17}}$  i.e.,  $x^{-17}$ .

$$\begin{aligned}
 \therefore T_{r+1} &= {}^{15}C_r (x^4)^{15-r} \left(-\frac{1}{x^3}\right)^r \\
 &= {}^{15}C_r x^{60-4r} (-1)^r x^{-3r} \\
 &= {}^{15}C_r x^{60-7r} (-1)^r
 \end{aligned}$$

For the coefficient  $x^{-17}$ ,

$$60 - 7r = -17$$

$$\Rightarrow 7r = 77 \Rightarrow r = 11$$

$$\Rightarrow T_{11+1} = {}^{15}C_{11} x^{60-77} (-1)^{11}$$

$$\begin{aligned}
 \therefore \text{Coefficient of } x^{-17} &= \frac{-15 \times 14 \times 13 \times 12 \times 11!}{11! \times 4 \times 3 \times 2 \times 1} \\
 &= -15 \times 7 \times 13 = -1365
 \end{aligned}$$

**Q. 8** Find the sixth term of the expansion  $(y^{1/2} + x^{1/3})^n$ , if the Binomial coefficient of the third term from the end is 45.

**Sol.** Given expansion is  $(y^{1/2} + x^{1/3})^n$ .

The sixth term of this expansion is

$$T_6 = T_{5+1} = {}^n C_5 (y^{1/2})^{n-5} (x^{1/3})^5 \quad \dots(i)$$

Now, given that the Binomial coefficient of the third term from the end is 45.

We know that, Binomial coefficient of third term from the end = Binomial coefficient of third term from the beginning =  ${}^n C_2$

$$\therefore {}^n C_2 = 45$$

$$\Rightarrow \frac{n(n-1)(n-2)!}{2!(n-2)!} = 45$$

$$\Rightarrow n(n-1) = 90$$

$$\Rightarrow n^2 - n - 90 = 0$$

$$\Rightarrow n^2 - 10n + 9n - 90 = 0$$

$$\Rightarrow n(n-10) + 9(n-10) = 0$$

$$\Rightarrow (n-10)(n+9) = 0$$

$$\Rightarrow (n+9) = 0 \text{ or } (n-10) = 0$$

$$\therefore n = 10 \quad [\because n \neq -9]$$

From Eq. (i),

$$T_6 = {}^{10} C_5 y^{5/2} x^{5/3} = 252 y^{5/2} \cdot x^{5/3}$$

**Q. 9** Find the value of  $r$ , if the coefficients of  $(2r+4)$ th and  $(r-2)$ th terms in the expansion of  $(1+x)^{18}$  are equal.

**Thinking Process**

*Coefficient of  $(r+1)$ th term in the expansion of  $(1+x)^n$  is  ${}^n C_r$ . Use this formula to solve the above problem.*

**Sol.** Given expansion is  $(1+x)^{18}$ .

Now,  $(2r+4)$ th term i.e.,  $T_{2r+3+1}$ .

$$\begin{aligned} \therefore T_{2r+3+1} &= {}^{18} C_{2r+3} (1)^{18-2r-3} (x)^{2r+3} \\ &= {}^{18} C_{2r+3} x^{2r+3} \end{aligned}$$

Now,  $(r-2)$ th term i.e.,  $T_{r-3+1}$ .

$$\therefore T_{r-3+1} = {}^{18} C_{r-3} x^{r-3}$$

$$\text{As, } {}^{18} C_{2r+3} = {}^{18} C_{r-3} \quad [\because {}^n C_x = {}^n C_y \Rightarrow x + y = n]$$

$$\Rightarrow 2r+3 + r-3 = 18$$

$$\Rightarrow 3r = 18$$

$$\therefore r = 6$$

**Q. 10** If the coefficient of second, third and fourth terms in the expansion of  $(1+x)^{2n}$  are in AP, then show that  $2n^2 - 9n + 7 = 0$ .

**Thinking Process**

*In the expansion of  $(x+y)^n$ , the coefficient of  $(r+1)$ th term is  ${}^n C_r$ . Use this formula to get the required coefficient. If  $a, b$  and  $c$  are in AP, then  $2b = a + c$ .*

**Sol.** Given expansion is  $(1 + x)^{2n}$ .

Now, coefficient of 2nd term =  ${}^{2n}C_1$

Coefficient of 3rd term =  ${}^{2n}C_2$

Coefficient of 4th term =  ${}^{2n}C_3$

Given that,  ${}^{2n}C_1$ ,  ${}^{2n}C_2$  and  ${}^{2n}C_3$  are in AP.

Then,  $2 \cdot {}^{2n}C_2 = {}^{2n}C_1 + {}^{2n}C_3$

$$\Rightarrow 2 \left[ \frac{2n(2n-1)(2n-2)!}{2 \times 1 \times (2n-2)!} \right] = \frac{2n(2n-1)!}{(2n-1)!} + \frac{2n(2n-1)(2n-2)(2n-3)!}{3!(2n-3)!}$$

$$\Rightarrow n(2n-1) = n + \frac{n(2n-1)(2n-2)}{6}$$

$$\Rightarrow n(12n-6) = n(6+4n^2-4n-2n+2)$$

$$\Rightarrow 12n-6 = (4n^2-6n+8)$$

$$\Rightarrow 6(2n-1) = 2(2n^2-3n+4)$$

$$\Rightarrow 3(2n-1) = 2n^2-3n+4$$

$$\Rightarrow 2n^2-3n+4-6n+3=0$$

$$\Rightarrow 2n^2-9n+7=0$$

**Q. 11** Find the coefficient of  $x^4$  in the expansion of  $(1 + x + x^2 + x^3)^{11}$ .

**Sol.** Given, expansion =  $(1 + x + x^2 + x^3)^{11} = [(1 + x) + x^2(1 + x)]^{11}$   
 $= [(1 + x)(1 + x^2)]^{11} = (1 + x)^{11} \cdot (1 + x^2)^{11}$

Now, above expansion becomes

$$= ({}^{11}C_0 + {}^{11}C_1x + {}^{11}C_2x^2 + {}^{11}C_3x^3 + {}^{11}C_4x^4 + \dots)({}^{11}C_0 + {}^{11}C_1x^2 + {}^{11}C_2x^4 + \dots)$$

$$= (1 + 11x + 55x^2 + 165x^3 + 330x^4 + \dots)(1 + 11x^2 + 55x^4 + \dots)$$

$\therefore$  Coefficient of  $x^4 = 55 + 605 + 330 = 990$

## Long Answer Type Questions

**Q. 12** If  $p$  is a real number and the middle term in the expansion of  $\left(\frac{p}{2} + 2\right)^8$  is 1120, then find the value of  $p$ .

**Sol.** Given expansion is  $\left(\frac{p}{2} + 2\right)^8$ .

Here,  $n = 8$

[even]

Since, this Binomial expansion has only one middle term i.e.,  $\left(\frac{p}{2} + 2\right)^8$  th = 5th term

$$T_5 = T_{4+1} = {}^8C_4 \left(\frac{p}{2}\right)^{8-4} \cdot 2^4$$

$$\Rightarrow 1120 = {}^8C_4 p^4 \cdot 2^{-4} \cdot 2^4$$

$$\Rightarrow 1120 = \frac{8 \times 7 \times 6 \times 5 \times 4!}{4! \times 4 \times 3 \times 2 \times 1} p^4$$

$$\begin{aligned} \Rightarrow & 1120 = 7 \times 2 \times 5 \times p^4 \\ \Rightarrow & p^4 = \frac{1120}{70} = 16 \Rightarrow p^4 = 2^4 \\ \Rightarrow & p^2 = 4 \Rightarrow p = \pm 2 \end{aligned}$$

**Q. 13** Show that the middle term in the expansion of  $\left(x - \frac{1}{x}\right)^{2n}$  is

$$\frac{1 \times 3 \times 5 \times \dots \times (2n-1)}{n!} \times (-2)^n.$$

**Sol.** Given, expansion is  $\left(x - \frac{1}{x}\right)^{2n}$ . This Binomial expansion has even power. So, this has one middle term.

*i.e.*,  $\left(\frac{2n}{2} + 1\right)$ th term =  $(n + 1)$ th term

$$\begin{aligned} T_{n+1} &= {}^{2n}C_n (x)^{2n-n} \left(-\frac{1}{x}\right)^n = {}^{2n}C_n x^n (-1)^n x^{-n} \\ &= {}^{2n}C_n (-1)^n = (-1)^n \frac{(2n)!}{n!n!} = \frac{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \dots (2n-1)(2n)}{n!n!} (-1)^n \\ &= \frac{1 \cdot 3 \cdot 5 \dots (2n-1) \cdot 2 \cdot 4 \cdot 6 \dots (2n)}{12 \cdot 3 \dots n(n!)} (-1)^n \\ &= \frac{1 \cdot 3 \cdot 5 \dots (2n-1) \cdot 2^n (1 \cdot 2 \cdot 3 \dots n)}{(1 \cdot 2 \cdot 3 \dots n)(n!)} (-1)^n \\ &= \frac{[1 \cdot 3 \cdot 5 \dots (2n-1)]}{n!} (-2)^n \end{aligned}$$

Hence proved.

**Q. 14** Find  $n$  in the Binomial  $\left(\sqrt[3]{2} + \frac{1}{\sqrt[3]{3}}\right)^n$ , if the ratio of 7th term from the beginning to the 7th term from the end is  $\frac{1}{6}$ .

**Sol.** Here, the Binomial expansion is  $\left(\sqrt[3]{2} + \frac{1}{\sqrt[3]{3}}\right)^n$ .

$$\text{Now, 7th term from beginning } T_7 = T_{6+1} = {}^nC_6 (\sqrt[3]{2})^{n-6} \left(\frac{1}{\sqrt[3]{3}}\right)^6 \quad \dots(i)$$

and 7th term from end *i.e.*,  $T_7$  from the beginning of  $\left(\frac{1}{\sqrt[3]{3}} + \sqrt[3]{2}\right)^n$

$$\text{i.e., } T_7 = {}^nC_6 \left(\frac{1}{\sqrt[3]{3}}\right)^{n-6} (\sqrt[3]{2})^6 \quad \dots(ii)$$

$$\text{Given that, } \frac{{}^nC_6 (\sqrt[3]{2})^{n-6} \left(\frac{1}{\sqrt[3]{3}}\right)^6}{{}^nC_6 \left(\frac{1}{\sqrt[3]{3}}\right)^{n-6} (\sqrt[3]{2})^6} = \frac{1}{6} \Rightarrow \frac{2^{\frac{n-6}{3}} \cdot 3^{-6/3}}{3^{-\left(\frac{n-6}{3}\right)} \cdot 2^{6/3}} = \frac{1}{6}$$

$$\Rightarrow \left(2^{\frac{n-6}{3}} \cdot 2^{-\frac{6}{3}}\right) \left(\frac{-6}{3} \cdot 3^{\frac{(n-6)}{3}}\right) = 6^{-1}$$

$$\Rightarrow \left(2^{\frac{n-6}{3}-\frac{6}{3}}\right) \cdot \left(3^{\frac{n-6}{3}-\frac{6}{3}}\right) = 6^{-1} \Rightarrow (2 \cdot 3)^{\frac{n}{3}-4} = 6^{-1}$$

$$\Rightarrow \frac{n}{3} - 4 = -1 \Rightarrow \frac{n}{3} = 3$$

$$\therefore n = 9$$

**Q. 15** In the expansion of  $(x+a)^n$ , if the sum of odd terms is denoted by  $O$  and the sum of even term by  $E$ . Then, prove that

$$(i) O^2 - E^2 = (x^2 - a^2)^n.$$

$$(ii) 4OE = (x+a)^{2n} - (x-a)^{2n}.$$

**Sol.** (i) Given expansion is  $(x+a)^n$ .

$$\therefore (x+a)^n = {}^nC_0 x^n a^0 + {}^nC_1 x^{n-1} a^1 + {}^nC_2 x^{n-2} a^2 + {}^nC_3 x^{n-3} a^3 + \dots + {}^nC_n a^n$$

Now, sum of odd terms

$$i.e., O = {}^nC_0 x^n + {}^nC_2 x^{n-2} a^2 + \dots$$

and sum of even terms

$$i.e., E = {}^nC_1 x^{n-1} a + {}^nC_3 x^{n-3} a^3 + \dots$$

$$\therefore (x+a)^n = O + E \quad \dots(i)$$

$$\text{Similarly, } (x-a)^n = O - E \quad \dots(ii)$$

$$\therefore (O+E)(O-E) = (x+a)^n (x-a)^n \quad [\text{on multiplying Eqs. (i) and (ii)}]$$

$$\Rightarrow O^2 - E^2 = (x^2 - a^2)^n$$

$$(ii) 4OE = (O+E)^2 - (O-E)^2 = [(x+a)^n]^2 - [(x-a)^n]^2 \quad [\text{from Eqs. (i) and (ii)}]$$

$$= (x+a)^{2n} - (x-a)^{2n} \quad \text{Hence proved.}$$

**Q. 16** If  $x^p$  occurs in the expansion of  $\left(x^2 + \frac{1}{x}\right)^{2n}$ , then prove that its

$$\text{coefficient is } \frac{2n!}{3! (4n-p)! (2n+p)!}.$$

**Sol.** Given expansion is  $\left(x^2 + \frac{1}{x}\right)^{2n}$ .

Let  $x^p$  occur in the expansion of  $\left(x^2 + \frac{1}{x}\right)^{2n}$ .

$$T_{r+1} = {}^{2n}C_r (x^2)^{2n-r} \left(\frac{1}{x}\right)^r$$

$$= {}^{2n}C_r x^{4n-2r} x^{-r} = {}^{2n}C_r x^{4n-3r}$$

$$\text{Let } 4n - 3r = p$$

$$\Rightarrow 3r = 4n - p \Rightarrow r = \frac{4n-p}{3}$$

$$\therefore \text{Coefficient of } x^p = {}^{2n}C_r = \frac{(2n)!}{r!(2n-r)!} = \frac{(2n)!}{\left(\frac{4n-p}{3}\right)! \left(2n - \frac{4n-p}{3}\right)!}$$

$$= \frac{(2n)!}{\left(\frac{4n-p}{3}\right)! \left(\frac{6n-4n+p}{3}\right)!} = \frac{(2n)!}{\left(\frac{4n-p}{3}\right)! \left(\frac{2n+p}{3}\right)!}$$

**Q. 17** Find the term independent of  $x$  in the expansion of

$$(1 + x + 2x^3) \left( \frac{3}{2}x^2 - \frac{1}{3x} \right)^9.$$

**Sol.** Given expansion is  $(1 + x + 2x^3) \left( \frac{3}{2}x^2 - \frac{1}{3x} \right)^9$ .

Now, consider  $\left( \frac{3}{2}x^2 - \frac{1}{3x} \right)^9$

$$\begin{aligned} T_{r+1} &= {}^9C_r \left( \frac{3}{2}x^2 \right)^{9-r} \left( -\frac{1}{3x} \right)^r \\ &= {}^9C_r \left( \frac{3}{2} \right)^{9-r} x^{18-2r} \left( -\frac{1}{3} \right)^r x^{-r} = {}^9C_r \left( \frac{3}{2} \right)^{9-r} \left( -\frac{1}{3} \right)^r x^{18-3r} \end{aligned}$$

Hence, the general term in the expansion of  $(1 + x + 2x^3) \left( \frac{3}{2}x^2 - \frac{1}{3x} \right)^9$

$$= {}^9C_r \left( \frac{3}{2} \right)^{9-r} \left( -\frac{1}{3} \right)^r x^{18-3r} + {}^9C_r \left( \frac{3}{2} \right)^{9-r} \left( -\frac{1}{3} \right)^r x^{19-3r} + 2 \cdot {}^9C_r \left( \frac{3}{2} \right)^{9-r} \left( -\frac{1}{3} \right)^r x^{21-3r}$$

For term independent of  $x$ , putting  $18 - 3r = 0$ ,  $19 - 3r = 0$  and  $21 - 3r = 0$ , we get

$$r = 6, r = 19/3, r = 7$$

Since, the possible value of  $r$  are 6 and 7.

Hence, second term is not independent of  $x$ .

$\therefore$  The term independent of  $x$  is  ${}^9C_6 \left( \frac{3}{2} \right)^{9-6} \left( -\frac{1}{3} \right)^6 + 2 \cdot {}^9C_7 \left( \frac{3}{2} \right)^{9-7} \left( -\frac{1}{3} \right)^7$

$$\begin{aligned} &= \frac{9 \times 8 \times 7 \times 6!}{6! \times 3 \times 2} \cdot \frac{3^3}{2^3} \cdot \frac{1}{3^6} - 2 \cdot \frac{9 \times 8 \times 7!}{7! \times 2 \times 1} \cdot \frac{3^2}{2^2} \cdot \frac{1}{3^7} \\ &= \frac{84}{8} \cdot \frac{1}{3^3} - \frac{36}{4} \cdot \frac{2}{3^5} = \frac{7}{18} - \frac{2}{27} = \frac{21-4}{54} = \frac{17}{54} \end{aligned}$$

## Objective Type Questions

**Q. 18** The total number of terms in the expansion of  $(x + a)^{100} + (x - a)^{100}$  after simplification is

- (a) 50                      (b) 202                      (c) 51                      (d) None of these

**Sol. (c)** Here,  $(x + a)^{100} + (x - a)^{100}$

Total number of terms is 102 in the expansion of  $(x + a)^{100} + (x - a)^{100}$

50 terms of  $(x + a)^{100}$  cancel out 50 terms of  $(x - a)^{100}$ . 51 terms of  $(x + a)^{100}$  get added to the 51 terms of  $(x - a)^{100}$ .

**Alternate Method**

$$\begin{aligned} (x + a)^{100} + (x - a)^{100} &= {}^{100}C_0 x^{100} + {}^{100}C_1 x^{99}a + \dots + {}^{100}C_{100} a^{100} \\ &\quad + {}^{100}C_0 x^{100} - {}^{100}C_1 x^{99}a + \dots + {}^{100}C_{100} a^{100} \\ &= 2 \left[ \underbrace{{}^{100}C_0 x^{100} + {}^{100}C_2 x^{98} a^2 + \dots + {}^{100}C_{100} a^{100}}_{51 \text{ terms}} \right] \end{aligned}$$

**Q. 19** If the integers  $r > 1$ ,  $n > 2$  and coefficients of  $(3r)$ th and  $(r + 2)$ nd terms in the Binomial expansion of  $(1 + x)^{2n}$  are equal, then

- (a)  $n = 2r$  (b)  $n = 3r$   
(c)  $n = 2r + 1$  (d) None of these

**💡 Thinking Process**

In the expansion of  $(x + y)^n$ , the coefficient of  $(r + 1)$ th term is  ${}^n C_r$ .

**Sol. (a)** Given that,  $r > 1$ ,  $n > 2$  and the coefficients of  $(3r)$ th and  $(r + 2)$ th term are equal in the expansion of  $(1 + x)^{2n}$ .

Then,  $T_{3r} = T_{3r-1+1} = {}^{2n} C_{3r-1} x^{3r-1}$

and  $T_{r+2} = T_{r+1+1} = {}^{2n} C_{r+1} x^{r+1}$

Given,  ${}^{2n} C_{3r-1} = {}^{2n} C_{r+1}$  [ $\because {}^n C_x = {}^n C_y \Rightarrow x + y = n$ ]

$\Rightarrow 3r - 1 + r + 1 = 2n$

$\Rightarrow 4r = 2n \Rightarrow n = \frac{4r}{2}$

$\therefore n = 2r$

**Q. 20** The two successive terms in the expansion of  $(1 + x)^{24}$  whose coefficients are in the ratio 1 : 4 are

- (a) 3rd and 4th (b) 4th and 5th  
(c) 5th and 6th (d) 6th and 7th

**Sol. (c)** Let two successive terms in the expansion of  $(1 + x)^{24}$  are  $(r + 1)$ th and  $(r + 2)$ th terms.

$\therefore T_{r+1} = {}^{24} C_r x^r$

and  $T_{r+2} = {}^{24} C_{r+1} x^{r+1}$

Given that,  $\frac{{}^{24} C_r}{{}^{24} C_{r+1}} = \frac{1}{4}$

$\Rightarrow \frac{\frac{(24)!}{r!(24-r)!}}{(24)!} = \frac{1}{4}$

$\Rightarrow \frac{(r+1)!(24-r-1)!}{r!(24-r)(23-r)!} = \frac{1}{4}$

$\Rightarrow \frac{r+1}{24-r} = \frac{1}{4} \Rightarrow 4r + 4 = 24 - r$

$\Rightarrow 5r = 20 \Rightarrow r = 4$

$\therefore T_{4+1} = T_5$  and  $T_{4+2} = T_6$

Hence, 5th and 6th terms.

**Q. 21** The coefficient of  $x^n$  in the expansion of  $(1+x)^{2n}$  and  $(1+x)^{2n-1}$  are in the ratio

(a) 1 : 2

(b) 1 : 3

(c) 3 : 1

(d) 2 : 1

**Sol. (d)**  $\therefore$  Coefficient of  $x^n$  in the expansion of  $(1+x)^{2n} = {}^{2n}C_n$   
and coefficient of  $x^n$  in the expansion of  $(1+x)^{2n-1} = {}^{2n-1}C_n$

$$\begin{aligned} \therefore \frac{{}^{2n}C_n}{{}^{2n-1}C_n} &= \frac{\frac{(2n)!}{n!n!}}{\frac{(2n-1)!}{n!(n-1)!}} \\ &= \frac{(2n)!n!(n-1)!}{n!n!(2n-1)!} \\ &= \frac{2n(2n-1)!n!(n-1)!}{n!n(n-1)!(2n-1)!} \\ &= \frac{2n}{n} = \frac{2}{1} = 2 : 1 \end{aligned}$$

**Q. 22** If the coefficients of 2nd, 3rd and the 4th terms in the expansion of  $(1+x)^n$  are in AP, then the value of  $n$  is

(a) 2

(b) 7

(c) 11

(d) 14

**Sol. (b)** The expansion of  $(1+x)^n$  is  ${}^nC_0 + {}^nC_1x + {}^nC_2x^2 + {}^nC_3x^3 + \dots + {}^nC_nx^n$

$\therefore$  Coefficient of 2nd term =  ${}^nC_1$ ,

Coefficient of 3rd term =  ${}^nC_2$ ,

and coefficient of 4th term =  ${}^nC_3$ .

Given that,  ${}^nC_1$ ,  ${}^nC_2$  and  ${}^nC_3$  are in AP.

$$\therefore 2 {}^nC_2 = {}^nC_1 + {}^nC_3$$

$$\Rightarrow 2 \left[ \frac{(n)!}{(n-2)!2!} \right] = \frac{(n)!}{(n-1)!} + \frac{(n)!}{3!(n-3)!}$$

$$\Rightarrow \frac{2 \cdot n(n-1)(n-2)!}{(n-2)!2!} = \frac{n(n-1)!}{(n-1)!} + \frac{n(n-1)(n-2)(n-3)!}{3 \cdot 2 \cdot 1(n-3)!}$$

$$\Rightarrow n(n-1) = n + \frac{n(n-1)(n-2)}{6}$$

$$\Rightarrow 6n - 6 = 6 + n^2 - 3n + 2$$

$$\Rightarrow n^2 - 9n + 14 = 0$$

$$\Rightarrow n^2 - 7n - 2n + 14 = 0$$

$$\Rightarrow n(n-7) - 2(n-7) = 0$$

$$\Rightarrow (n-7)(n-2) = 0$$

$$\therefore n = 2 \text{ or } n = 7$$

Since,  $n = 2$  is not possible.

$$\therefore n = 7$$

**Q. 23** If  $A$  and  $B$  are coefficient of  $x^n$  in the expansions of  $(1+x)^{2n}$  and  $(1+x)^{2n-1}$  respectively, then  $\frac{A}{B}$  equals to

- (a) 1 (b) 2  
(c)  $\frac{1}{2}$  (d)  $\frac{1}{n}$

**Sol. (b)** Since, the coefficient of  $x^n$  in the expansion of  $(1+x)^{2n}$  is  ${}^{2n}C_n$ .

$$\therefore A = {}^{2n}C_n$$

Now, the coefficient of  $x^n$  in the expansion of  $(1+x)^{2n-1}$  is  ${}^{2n-1}C_n$ .

$$\therefore B = {}^{2n-1}C_n$$

$$\text{Now, } \frac{A}{B} = \frac{{}^{2n}C_n}{{}^{2n-1}C_n} = \frac{2}{1} = 2$$

Same as solution No. 21.

**Q. 24** If the middle term of  $\left(\frac{1}{x} + x \sin x\right)^{10}$  is equal to  $7\frac{7}{8}$ , then the value of  $x$  is

- (a)  $2n\pi + \frac{\pi}{6}$  (b)  $n\pi + \frac{\pi}{6}$   
(c)  $n\pi + (-1)^n \frac{\pi}{6}$  (d)  $n\pi + (-1)^n \frac{\pi}{3}$

**Sol. (c)** Given expansion is  $\left(\frac{1}{x} + x \sin x\right)^{10}$ .

Since,  $n = 10$  is even, so this expansion has only one middle term i.e., 6th term.

$$\therefore T_6 = T_{5+1} = {}^{10}C_5 \left(\frac{1}{x}\right)^{10-5} (x \sin x)^5$$

$$\Rightarrow \frac{63}{8} = {}^{10}C_5 x^{-5} x^5 \sin^5 x$$

$$\Rightarrow \frac{63}{8} = \frac{10 \cdot 9 \cdot 8 \cdot 7 \cdot 6 \cdot 5!}{5 \cdot 4 \cdot 3 \cdot 2 \cdot 1 \cdot 5!} \sin^5 x$$

$$\Rightarrow \frac{63}{8} = 2 \cdot 9 \cdot 2 \cdot 7 \cdot \sin^5 x$$

$$\Rightarrow \sin^5 x = \frac{1}{32}$$

$$\Rightarrow \sin^5 x = \left(\frac{1}{2}\right)^5$$

$$\Rightarrow \sin x = \frac{1}{2}$$

$$\therefore x = n\pi + (-1)^n \pi / 6$$

## Fillers

**Q. 25** The largest coefficient in the expansion of  $(1+x)^{30}$  is .....

**Thinking Process**

In the expansion of  $(1+x)^n$ , the largest coefficient is  ${}^nC_{n/2}$  (when  $n$  is even).

**Sol.** Largest coefficient in the expansion of  $(1+x)^{30} = {}^{30}C_{30/2} = {}^{30}C_{15}$

**Q. 26** The number of terms in the expansion of  $(x+y+z)^n$  .....

**Sol.** Given expansion is  $(x+y+z)^n = [x+(y+z)]^n$ .

$$[x+(y+z)]^n = {}^nC_0x^n + {}^nC_1x^{n-1}(y+z) + {}^nC_2x^{n-2}(y+z)^2 + \dots + {}^nC_n(y+z)^n$$

$\therefore$  Number of terms =  $1+2+3+\dots+n+(n+1)$

$$= \frac{(n+1)(n+2)}{2}$$

**Q. 27** In the expansion of  $\left(x^2 - \frac{1}{x^2}\right)^{16}$ , the value of constant term is .....

**Sol.** Let constant be  $T_{r+1}$ .

$$\begin{aligned} \therefore T_{r+1} &= {}^{16}C_r(x^2)^{16-r}\left(-\frac{1}{x^2}\right)^r \\ &= {}^{16}C_r x^{32-2r}(-1)^r x^{-2r} \\ &= {}^{16}C_r x^{32-4r}(-1)^r \end{aligned}$$

For constant term,  $32-4r=0 \Rightarrow r=8$

$$\therefore T_{8+1} = {}^{16}C_8$$

**Q. 28** If the seventh term from the beginning and the end in the expansion of  $\left(\sqrt[3]{2} + \frac{1}{\sqrt[3]{3}}\right)^n$  are equal, then  $n$  equals to .....

**Sol.** Given expansions is  $\left(\sqrt[3]{2} + \frac{1}{\sqrt[3]{3}}\right)^n$ .

$$\therefore T_7 = T_{6+1} = {}^nC_6(\sqrt[3]{2})^{n-6}\left(\frac{1}{\sqrt[3]{3}}\right)^6 \quad \dots(i)$$

Since,  $T_7$  from end is same as the  $T_7$  from beginning of  $\left(\frac{1}{\sqrt[3]{3}} + \sqrt[3]{2}\right)^n$ .

$$\text{Then, } T_7 = {}^nC_6\left(\frac{1}{\sqrt[3]{3}}\right)^{n-6}(\sqrt[3]{2})^6 \quad \dots(ii)$$

$$\text{Given that, } {}^nC_6(2)^{\frac{n-6}{3}}(3)^{-6/3} = {}^nC_6(3)^{-\frac{(n-6)}{3}}2^{6/3}$$

$$\Rightarrow (2)^{\frac{n-12}{3}} = \left(\frac{1}{3^{1/3}}\right)^{n-12}$$

which is true, when  $\frac{n-12}{3} = 0$ .

$$\Rightarrow n-12=0 \Rightarrow n=12$$

**Q. 29** The coefficient of  $a^{-6}b^4$  in the expansion of  $\left(\frac{1}{a} - \frac{2b}{3}\right)^{10}$  is .....

**Thinking Process**

In the expansion of  $(x-a)^n$ ,  $T_{r+1} = {}^nC_r x^{n-r}(-a)^r$

**Sol.** Given expansion is  $\left(\frac{1}{a} - \frac{2b}{3}\right)^{10}$ .

Let  $T_{r+1}$  has the coefficient of  $a^{-6}b^4$ .

$$\therefore T_{r+1} = {}^{10}C_r \left(\frac{1}{a}\right)^{10-r} \left(-\frac{2b}{3}\right)^r$$

For coefficient of  $a^{-6}b^4$ ,  $10 - r = 6 \Rightarrow r = 4$

$$\text{Coefficient of } a^{-6}b^4 = {}^{10}C_4 (-2/3)^4$$

$$\therefore = \frac{10 \cdot 9 \cdot 8 \cdot 7 \cdot 6!}{6! \cdot 4 \cdot 3 \cdot 2 \cdot 1} \cdot \frac{2^4}{3^4} = \frac{1120}{27}$$

**Q. 30** Middle term in the expansion of  $(a^3 + ba)^{28}$  is .....

**Sol.** Given expansion is  $(a^3 + ba)^{28}$ .

$$\therefore n = 28 \quad \text{[even]}$$

$$\therefore \text{Middle term} = \left(\frac{28}{2} + 1\right)\text{th term} = 15\text{th term}$$

$$\begin{aligned} \therefore T_{15} &= T_{14+1} \\ &= {}^{28}C_{14} (a^3)^{28-14} (ba)^{14} \\ &= {}^{28}C_{14} a^{42} b^{14} a^{14} \\ &= {}^{28}C_{14} a^{56} b^{14} \end{aligned}$$

**Q. 31** The ratio of the coefficients of  $x^p$  and  $x^q$  in the expansion of  $(1+x)^{p+q}$  is .....

**Sol.** Given expansion is  $(1+x)^{p+q}$ .

$$\therefore \text{Coefficient of } x^p = {}^{p+q}C_p$$

$$\text{and coefficient of } x^q = {}^{p+q}C_q$$

$$\therefore \frac{{}^{p+q}C_p}{{}^{p+q}C_q} = \frac{{}^{p+q}C_p}{{}^{p+q}C_p} = 1:1$$

**Q. 32** The position of the term independent of  $x$  in the expansion of

$$\left(\sqrt{\frac{x}{3}} + \frac{3}{2x^2}\right)^{10} \text{ is .....$$

**Sol.** Given expansion is  $\left(\sqrt{\frac{x}{3}} + \frac{3}{2x^2}\right)^{10}$ .

Let the constant term be  $T_{r+1}$ .

Then,

$$\begin{aligned}
 T_{r+1} &= {}^{10}C_r \left( \sqrt{\frac{x}{3}} \right)^{10-r} \left( \frac{3}{2x^2} \right)^r \\
 &= {}^{10}C_r \cdot x^{\frac{10-r}{2}} \cdot 3^{\frac{-10+r}{2}} \cdot 3^r \cdot 2^{-r} \cdot x^{-2r} \\
 &= {}^{10}C_r \cdot x^{\frac{10-5r}{2}} \cdot 3^{\frac{-10+3r}{2}} \cdot 2^{-r}
 \end{aligned}$$

For constant term,  $10 - 5r = 0 \Rightarrow r = 2$   
Hence, third term is independent of  $x$ .

**Q. 33** If  $25^{15}$  is divided by 13, then the remainder is .....

**Sol.** Let

$$\begin{aligned}
 25^{15} &= (26 - 1)^{15} \\
 &= {}^{15}C_0 26^{15} - {}^{15}C_1 26^{14} + \dots - {}^{15}C_{15} \\
 &= {}^{15}C_0 26^{15} - {}^{15}C_1 26^{14} + \dots - 1 - 13 + 13 \\
 &= {}^{15}C_0 26^{15} - {}^{15}C_1 26^{14} + \dots - 13 + 12
 \end{aligned}$$

It is clear that, when  $25^{15}$  is divided by 13, then remainder will be 12.

## True/False

**Q. 34** The sum of the series  $\sum_{r=0}^{10} {}^{20}C_r$  is  $2^{19} + \frac{{}^{20}C_{10}}{2}$ .

**Sol. False**

$$\begin{aligned}
 \text{Given series} &= \sum_{r=0}^{10} {}^{20}C_r = {}^{20}C_0 + {}^{20}C_1 + {}^{20}C_2 + \dots + {}^{20}C_{10} \\
 &= {}^{20}C_0 + {}^{20}C_1 + \dots + {}^{20}C_{10} + {}^{20}C_{11} + \dots + {}^{20}C_{20} - ({}^{20}C_{11} + \dots + {}^{20}C_{20}) \\
 &= 2^{20} - ({}^{20}C_{11} + \dots + {}^{20}C_{20})
 \end{aligned}$$

Hence, the given statement is false.

**Q. 35** The expression  $7^9 + 9^7$  is divisible by 64.

**Sol. True**

$$\begin{aligned}
 \text{Given expression} &= 7^9 + 9^7 = (1 + 8)^7 - (1 - 8)^9 \\
 &= ({}^7C_0 + {}^7C_1 8 + {}^7C_2 8^2 + \dots + {}^7C_7 8^7) - ({}^9C_0 - {}^9C_1 8 + {}^9C_2 8^2 \dots - {}^9C_9 8^9) \\
 &= (1 + 7 \times 8 + 21 \times 8^2 + \dots) - (1 - 9 \times 8 + 36 \times 8^2 + \dots - 8^9) \\
 &= (7 \times 8 + 9 \times 8) + (21 \times 8^2 - 36 \times 8^2) + \dots \\
 &= 2 \times 64 + (21 - 36)64 + \dots
 \end{aligned}$$

which is divisible by 64.

Hence, the statement is true.

**Q. 36** The number of terms in the expansion of  $[(2x + y^3)^4]^7$  is 8.

**Sol. False**

$$\text{Given expansion is } [(2x + y^3)^4]^7 = (2x + y^3)^{28}.$$

Since, this expansion has 29 terms.

So, the given statement is false.

**Q. 37** The sum of coefficients of the two middle terms in the expansion of  $(1+x)^{2n-1}$  is equal to  ${}^{2n-1}C_n$ .

**Sol. False**

Here, the Binomial expansion is  $(1+x)^{2n-1}$ .

Since, this expansion has two middle term i.e.,  $\left(\frac{2n-1+1}{2}\right)$ th term and  $\left(\frac{2n-1+1}{2}+1\right)$ th

term i.e.,  $n$ th term and  $(n+1)$ th term.

$$\therefore \text{Coefficient of } n\text{th term} = {}^{2n-1}C_{n-1}$$

$$\text{Coefficient of } (n+1)\text{th term} = {}^{2n-1}C_n$$

$$\text{Sum of coefficients} = {}^{2n-1}C_{n-1} + {}^{2n-1}C_n$$

$$= {}^{2n-1+1}C_n = {}^{2n}C_n \quad [{}^nC_r + {}^nC_{r-1} = {}^{n+1}C_r]$$

**Q. 38** The last two digits of the numbers  $3^{400}$  are 01.

**Sol. True**

$$\text{Given that, } 3^{400} = 9^{200} = (10-1)^{200}$$

$$\Rightarrow (10-1)^{200} = {}^{200}C_0 10^{200} - {}^{200}C_1 10^{199} + \dots - {}^{200}C_{199} 10^1 + {}^{200}C_{200} 1^{200}$$

$$\Rightarrow (10-1)^{200} = 10^{200} - 200 \times 10^{199} + \dots - 10 \times 200 + 1$$

So, it is clear that the last two digits are 01.

**Q. 39** If the expansion of  $\left(x - \frac{1}{x^2}\right)^{2n}$  contains a term independent of  $x$ , then  $n$  is a multiple of 2.

**Sol. False**

Given Binomial expansion is  $\left(x - \frac{1}{x^2}\right)^{2n}$ .

Let  $T_{r+1}$  term is independent of  $x$ .

$$\begin{aligned} \text{Then, } T_{r+1} &= {}^{2n}C_r (x)^{2n-r} \left(-\frac{1}{x^2}\right)^r \\ &= {}^{2n}C_r x^{2n-r} (-1)^r x^{-2r} = {}^{2n}C_r x^{2n-3r} (-1)^r \end{aligned}$$

For independent of  $x$ ,

$$2n - 3r = 0$$

$$\therefore r = \frac{2n}{3},$$

which is not a integer.

So, the given expansion is not possible.

**Q. 40** The number of terms in the expansion of  $(a+b)^n$ , where  $n \in N$ , is one less than the power  $n$ .

**Sol. False**

We know that, the number of terms in the expansion of  $(a+b)^n$ , where  $n \in N$ , is one more than the power  $n$ .