

## Rolle's Theorem

---

**Q.1.** Verify Rolle's theorem for the function :  $f(x) = e^{2x} (\sin 2x - \cos 2x)$  , defined in the interval  $[\pi/8, 5\pi/8]$  .

### Solution : 1

We have ,  $f(x) = e^{2x}(\sin^{2x} - \cos^{2x}) \times \epsilon [\pi/8, 5\pi/8]$

(1) As sine, cosine and exponential function are always continuous, hence given function  $f(x)$  is continuous in  $[\pi/8, 5\pi/8]$  .

$$\begin{aligned}(2) f'(x) &= e^{2x} \times 2 (\sin^{2x} - \cos^{2x}) + e^{2x} (2 \cos^{2x} + 2 \sin^{2x}) \\&= 2 e^{2x} (\sin^{2x} - \cos^{2x} + \cos^{2x} + \sin^{2x}) \\&= 2 e^{2x} (2 \sin^{2x}) = 4 e^{2x} \sin^{2x}.\end{aligned}$$

Thus derivatives exists in the given interval and function is differentiable.

$$(3) f(\pi/8) = e^{\pi/4} (\sin \pi/4 - \cos \pi/4) = e^{\pi/4} \times 0 = 0 .$$

$$f(5\pi/8) = e^{5\pi/4} (\sin 5\pi/4 - \cos 5\pi/4) = e^{5\pi/4} \times 0 = 0 .$$

$$\text{Therefore , } f(\pi/8) = f(5\pi/8)$$

$$\text{Now } f'(c) = 0$$

$$\text{Or, } 4 e^{2c} \sin 2c = 0$$

$$\text{Or, } \sin 2c = 0 \text{ [As } e^{2c} \neq 0] \text{ Hence, } 2c = 0, \pi, 2\pi, 3\pi \dots\dots\dots .$$

$$\text{Or, } c = 0, \pi/2, \pi, 3\pi/2 \dots\dots\dots .$$

$$\text{Therefore , } \pi/2 \epsilon (\pi/8, 5\pi/8) .$$

Hence Rolle's theorem is verified.

**Q.2.** Examine the validity and conclusion of Rolle's theorem for the function :

$$f(x) = e^x \cdot \sin^x , \text{ for all } x \epsilon [0, \pi] .$$

**Solution : 2**

We have ,  $f(x) = e^x \sin x$  , for all  $x \in [0, \pi]$

(1) As exponential function and trigonometric function are continuous , hence their product is also continuous in  $[0, \pi]$  i.e.  $f(x)$  is continuous in the given interval.

$$(2) f(x) = e^x \sin x$$

$$\text{Hence , } f'(x) = e^x \sin x + e^x \cos x .$$

Clearly  $f(x)$  exists in the open interval  $(0, \pi)$  .

$$(3) f(0) = e^0 \cdot \sin 0 = 0 ; f(\pi) = e^\pi \cdot \sin \pi = 0 .$$

$$\text{Since } f(0) = f(\pi) = 0 .$$

Hence all condition of Rolle's theorem is satisfied. Hence there exist 'c' , in  $0 < c < \pi$  such that  $f'(c) = 0$

$$\text{Or, } e^c (\sin c + \cos c) = 0$$

$$\text{Or, } \sin c + \cos c = 0 \text{ [As, } e^c \neq 0]$$

$$\text{Or, } \tan c = -1 = \tan 3\pi/4 .$$

As ,  $3\pi/4 \in [0, \pi]$  , **Rolle's theorem is verified.**

**Q.3.** Verify Rolle's theorem for the function  $f(x) = \log [(x^2 + ab)/\{x(a + b)\}]$  ,  $x \in [a, b]$  and  $x \neq 0$  .

**Solution : 3**

$$(1) f(x) = \log [(x^2 + ab)/\{x(a + b)\}] = \log (x^2 + ab) - \log(a + b) - \log x$$

Therefore ,  $f(x)$  is continuous in  $a \leq x \leq b$  .

$$(2) f'(x) = 2x/(x^2 + ab) - 1/x$$

$$= (2x^2 - x^2 - ab)/\{x(x^2 + ab)\}$$

$$= (x^2 - ab)/\{x(x^2 + ab)\} , \text{ which exist in } a < x < b .$$

$$(3) f(a) = \log (a^2 + ab) - \log (a + b) - \log a$$

$$= \log a + \log (a + b) - \log (a + b) - \log a = 0$$

$$f(b) = \log (b^2 + ab) - \log (a + b) - \log b$$

$$= \log b + \log (a + b) - \log (a + b) - \log b = 0$$

Hence ,  $f(a) = f(b)$  .

Hence , there exist  $c \in [a, b]$  , such that  $f'(c) = 0$  ,

$$\text{Or, } f'(c) = (c^2 - ab) / \{c(c^2 + ab)\} = 0$$

$$\text{Or, } c^2 - ab = 0 \Rightarrow c^2 = ab \Rightarrow c = \sqrt{ab} .$$

As,  $c = \sqrt{ab} \in [a, b]$  , **Rolle's theorem is verified.**

**Q.4.** Taking the function  $f(x) = (x - 3) \log x$ , prove that there is at least one value of  $x$  in  $(1, 3)$  which satisfies  $x \log x = 3 - x$ .

#### **Solution : 4**

(1) As both  $(x - 3)$  and  $\log x$  are continuous functions , hence  $f(x)$  is also a continuous in  $[1, 3]$  .

(2)  $f'(x) = (x - 3)/x + \log x$  which exist in  $(1, 3)$  .

(3)  $f(1) = 0 = f(3)$ . Therefore ,  $f(x)$  satisfies all three conditions. Rolle's theorem is applicable. As such there is at least one value of  $x$  in  $(1, 3)$  for which  $f'(x) = 0$ .

$$\text{Or, } (x - 3)/x + \log x = 0 \Rightarrow x \log x = 3 - x .$$

Or, one of the roots of  $x \log x = 3 - x$  will be in the interval  $(1, 3)$  . **[Proved.]**

**Q.5.** Apply Rolle's theorem to find point (or points) on the curve  $y = -1 + \cos x$  where the tangent is parallel to the  $x$ -axis in  $[0, 2\pi]$  .

#### **Solution : 5**

$$y = f(x) = -1 + \cos x \text{ ----- (1)}$$

As , cosine function is continuous function for all values of ,  $f(x)$  is continuous in  $[0, 2\pi]$  and differentiable in  $(0, 2\pi)$  .

Also  $f(0) = 0 - 1 + \cos 0 = -1 + 1 = 0$  ;  $f(2\pi) = -1 + 1 = 0$  .

Hence ,  $f(0) = f(2\pi)$  .

Thus all the three conditions of Rolle's theorem are satisfied by  $f(x)$  in  $[0, 2\pi]$ .

Hence , by Rolle's theorem there exist at least one real number  $x$  in  $(0, 2\pi)$  such that  $f'(x) = 0$ .

Now ,  $f'(x) = -\sin x$  , and  $f'(x) = 0 \Rightarrow -\sin x = 0 \Rightarrow \sin x = 0 \Rightarrow x = \pi \in [0, 2\pi]$  .

When  $x = \pi$ , (i)  $y = -1 + \cos \pi = -1 - 1 = -2$  . So, there exist a point  $(\pi, -2)$  on the given curve  $y = -1 + \cos x$  , where the tangent is parallel to the  $x$ -axis.

**Q.6.** Verify Rolle's theorem for the function  $f(x) = x^3 - 7x^2 + 16x - 12$  in the interval  $[2, 3]$ .

### **Solution : 6**

We have  $f(x) = x^3 - 7x^2 + 16x - 12$  in  $[2, 3]$  .

The function being a polynomial is continuous in  $[2, 3]$  and differentiable in  $(2, 3)$  . Also  $f(2) = 2^3 - 7 \times 2^2 + 16 \times 2 - 12 = 0 = f(3)$  .

Therefore , all the conditions of Rolle's theorem are satisfied , hence there exist at least one value  $c \in [2, 3]$  such that  $f'(c) = 0$  .

Now  $f'(x) = 3x^2 - 14x + 16$ .

And  $f'(x) = 0 \Rightarrow 3x^2 - 14x + 16 = 0$  .

Or,  $(x - 2)(3x - 8) = 0$

Or,  $x = 2, 8/3$ .

Clearly  $c = 8/3 \in (2, 3)$  and  $f'(c) = 0$  .

Hence , **Rolle's theorem is verified.**

**Q.7.** It is given that Rolle's theorem holds good for the function :  $f(x) = x^3 + ax^2 + bx$ ,  $x \in [1, 2]$  at the point  $x = 4/3$ . Find the values of  $a$  and  $b$ .

### **Solution : 7**

We have,  $f(x) = x^3 + ax^2 + bx$ ,  $x \in [1, 2]$

Then  $f(1) = 1 + a + b = 0$  ;  $f(2) = 8 + 4a + 2b = 0$

Adding the two we get,  $3a + b + 7 = 0$  ----- (i)

Differentiating we get,  $f'(x) = 3x^2 + 2ax + b = 0$

And  $f'(4/3) = 3(4/3)^2 + 2a(4/3) + b = 0$

Or,  $16/3 + 8a/3 + b = 0$

Or,  $8a + 3b + 16 = 0$  ----- (ii)

Solving (i) and (ii) we get,  $a = -5$ ,  $b = 8$ .