# Electrochemistry

### **Electrochemical cells:**

• Two types – Galvanic cell or voltaic cell, and Electrolytic cell

### Galvanic cell:

Converts the chemical energy of a spontaneous redox reaction into electrical energy

Daniell cell -

$$\Delta_r G^{\Theta} = -RT \ln K$$

The standard potential of a cell is given by

$$\Delta_r G^{\Theta} = -nFE_{cell}^{\odot}$$

The potential of an individual half cell cannot be measured.

Relation between the standard potential of a cell and standard Gibbs energy:

$$\Delta_r G^{\Theta} = -nFE_{cell}^{\odot}$$

Relation between standard Gibbs energy and the equilibrium constant:

$$\Delta_r G^{\Theta} = -RT \ln K$$

- Nernst equation:
  - Gives the concentration dependence of the potentials of the electrodes and the cells
  - For the electrode reaction

$$aA + bB \xrightarrow{ne^-} cC + dD$$

Nernst equation is given by

$$\mathbf{E}_{\text{cell}} = \mathbf{E}_{\text{cell}}^{\odot} - \frac{\mathbf{R}T}{nF} \ln \frac{\left[\mathbf{C}\right]^{\text{c}} \left[\mathbf{D}\right]^{\text{d}}}{\left[\mathbf{A}\right]^{\text{a}} \left[\mathbf{B}\right]^{\text{b}}}$$

Nernst equation is given by

$$aA + bB \xrightarrow{ne} cC + dD$$

Nernst equation is given by

$$\mathbf{E}_{\text{cell}} = \mathbf{E}_{\text{cell}}^{\odot} - \frac{\mathbf{R}T}{nF} \ln \frac{\left[\mathbf{C}\right]^{c} \left[\mathbf{D}\right]^{d}}{\left[\mathbf{A}\right]^{a} \left[\mathbf{B}\right]^{b}}$$

• For a general electrochemical reaction of the type

$$aA + bB \xrightarrow{ne} cC + dD$$

Nernst equation is given by

$$\mathbf{E}_{\text{cell}} = \mathbf{E}_{\text{cell}}^{\odot} - \frac{\mathbf{R}T}{nF} \ln \frac{\left[\mathbf{C}\right]^{c} \left[\mathbf{D}\right]^{a}}{\left[\mathbf{A}\right]^{a} \left[\mathbf{B}\right]^{b}}$$

# **Conductance of electrolytic solutions:**

Resistance,  $R = \rho \frac{l}{A}$ 

Where,

 $l \rightarrow \text{Length}$ 

 $A \rightarrow$  Area of cross-section

 $P \rightarrow Resistivity$  or specific resistance

Conductance, 
$$G = \frac{1}{R} = \frac{A}{\rho l} = \kappa \frac{A}{l}$$

Where, & Conductivity or specific conductance

The SI unit of conductance is  $\Omega^{-1}$  (siemens or mho).

The conductivity of an electrolyte depends upon

- nature of the solvent
- nature of the electrolyte added
- concentration of the electrolyte
- temperature

Molar conductivity,  $\Lambda_m = \frac{\kappa}{C}$ 

# Variation of conductivity

• For both strong and weak electrolytes, conductivity decreases with decrease in concentration.

# Variation of molar conductivity

• For both strong and weak electrolytes, molar conductivity increases with decrease in concentration.

**Limiting molar conductivity** – molar conductivity when concentration approaches zero

• Degree of dissociation, 
$$\alpha = \frac{\Lambda_m}{\Lambda_m^0}$$

• Kohlrausch law of independent migration of ions:

According to this law, for an electrolyte, the molar conductivity at infinite dilution is the sum of the contribution of the molar conductivity of the ions in which it dissociates.

# Electrolytic cells and electrolysis

### $1F = 96487 \text{ C mol}^{-1}$

- Faraday's first law of electrolysis: The amount of chemical reaction occurring at any electrode during the process of electrolysis by a current is proportional to the quantity of electricity passed through the electrolyte.
- Second law of electrolysis: The amounts of different substances liberated when same quantity of electricity is passed through the electrolytic solution are proportional to their chemical equivalent weights.

Battery is a galvanic cell in which chemical energy of the redox reaction is converted into electrical energy.

Mainly two types:

Primary batteries Secondary batteries

# • Primary Batteries

In primary batteries, reaction occurs only once.

After use over a period of time, these become dead and cannot be reused.

Examples:Dry cell (or Leclanche cell), Mercury cell

### Secondary Batteries

Secondary batteries can be recharged again by passing current through them in the opposite direction.


Examples:Lead storage battery, Nickel-cadmium cell

**Corrosion:** Oxidation of a metal by loss of electrons to oxygen and formation of oxides

### **Corrosion of Iron**

Known as rusting

The spot where oxidation takes place behaves as anode.



# **Prevention of Corrosion**

Preventing the surface of the metal from coming in contact with atmosphere By covering the surface with paint or chemicals such as bis-phenol By covering the surface with other metals such as Sn, Zn, Mg.

The Hydrogen Economy: Based on electrochemical principles