


Chemical Kinetics

For a reaction $R \rightarrow P$

$$r_{av} = \frac{-\Delta[R]}{\Delta t} = \frac{\Delta[P]}{\Delta t}$$




$$\text{Rate of reaction} = -\frac{1}{2} \frac{\Delta[\text{HI}]}{\Delta t} = \frac{\Delta[\text{H}_2]}{\Delta t} = \frac{\Delta[\text{I}_2]}{\Delta t}$$

Factors influencing the rate of a reaction:

Rate of a reaction depends upon the concentration of reactants (pressure in the case of gas), temperature and catalyst.

- Rate expression and rate constant



Rate expression

$$\therefore \text{Rate} \propto [A]^x [B]^y$$

Differential rate equation ®

$$\Rightarrow -\frac{d[R]}{dt} = k[A]^x [B]^y \quad (1)$$

Where, k is called rate constant

- Order of a reaction:


$$\text{Rate} = k[A]^x [B]^y$$

x is Order of the reaction with respect to A

y is Order of the reaction with respect to B

$x + y$ is Overall order of the reaction

1. Order of a reaction can be 0, 1, 2, 3 and even a fraction
2. Units of rate constant



$$\text{Rate} = k [A]^x [B]^y$$

$$x + y = n = \text{Order of the reaction}$$

$$k = \frac{[\text{Rate}]}{[A]^x [B]^y}$$

$$= \frac{\text{Concentration}}{\text{Time}} \times \frac{1}{(\text{Concentration})^n}$$

$$[[A] = [B] \text{ and } x + y = n = \text{Order of the reaction}]$$

1. For a zero-order reaction, $n = 0$

$$\text{Unit of } \frac{\text{mol L}^{-1}}{\text{s}} \times \frac{1}{(\text{mol L}^{-1})^0} = \text{mol L}^{-1} \text{s}^{-1}$$

1. For a first-order reaction, $n = 1$

$$\text{Unit of } \frac{\text{mol L}^{-1}}{\text{s}} \times \frac{1}{(\text{mol L}^{-1})^1} = \text{s}^{-1}$$

1. For a second-order reaction, $n = 2$

$$\text{Unit of } \frac{\text{mol L}^{-1}}{\text{s}} \times \frac{1}{(\text{mol L}^{-1})^2} = \text{mol}^{-1} \text{L s}^{-1}$$

- **Molecularity of a reaction:**


The number of reacting species (atoms, ions or molecules) taking part in an elementary reaction

- **Order versus molecularity**


1. Order can be zero and even a fraction. But molecularity cannot be zero or a non-integer.
2. Order is applicable to both elementary and complex reactions whereas molecularity is applicable to elementary reactions only.

Integrated rate equations:


- Zero-order reactions:



$$k = \frac{[R]_0 - [R]}{t}$$





- First-order reactions:




$$\text{Rate} = -\frac{d[R]}{dt} = k[R]$$

$$\therefore \frac{d[R]}{[R]} = -k dt$$



1. For a typical first-order gas phase reaction:



Here, p_i is initial pressure of A

$p_t = (p_A + p_B + p_C) = \text{Total pressure at time } t$

$$\therefore k = \frac{2.303}{t} \log \frac{p_i}{p_A}$$

$$\Rightarrow k = \frac{2.303}{t} \log \frac{p_i}{(2p_i - p_t)}$$

- **Half-life of a reaction:**

1. **For a zero-order reaction:**

$$t_{1/2} = \frac{[R]_0}{2k}$$


1. **For a first-order reaction:**

$$t_{1/2} = \frac{0.693}{k}$$

That is, $t_{1/2}$ is independent of $[R]_0$

- **Pseudo first-order reaction:**

Hydrolysis of ethyl acetate –



$$\text{Rate} = k' [\text{CH}_3\text{COOC}_2\text{H}_5] [\text{H}_2\text{O}]$$

Here, $[\text{H}_2\text{O}]$ is constant

So, rate = $k [\text{CH}_3\text{COOC}_2\text{H}_5]$

Where, $k = k' [\text{H}_2\text{O}]$

Temperature dependence of the rate of reaction:

- For a chemical reaction, with a rise in temperature by 10° , the rate constant is nearly doubled.
- **Arrhenius equation:**

$$k = A e^{-E_a/RT}$$


Where,

A is Arrhenius factor or frequency factor or pre-exponential factor

R is Gas constant

E_a is Activation energy

On taking natural logarithm on both sides, we have




If k_1 and k_2 are the values of rate constants at temperatures T_1 and T_2 respectively, then

$$\ln \frac{k_2}{k_1} = \frac{E_a}{R} \left[\frac{T_2 - T_1}{T_1 T_2} \right]$$

$$\text{or, } \log \frac{k_2}{k_1} = \frac{E_a}{2.303R} \left[\frac{T_2 - T_1}{T_1 T_2} \right]$$

- Effect of a catalyst:**

A catalyst provides an alternate pathway or reaction mechanism by reducing the activation energy.



- Collision frequency (Z) → The number of collisions per second per unit volume of the reaction mixture

For the reaction $A + B \rightarrow \text{Products}$,

Rate =

$$Z_{AB} e^{-E_a/RT}$$

Where,

Z_{AB} → Collision frequency of reactants A and B

$e^{-E_a/RT}$ → The fraction of molecules with energies equal to or greater than E_a

To account for effective collisions, another factor P (called the probability or steric factor) is introduced.

Then, rate =

$$PZ_{AB}e^{-E_a/RT}$$