


Haloalkanes and Haloarenes

Classification: Mono, di and polyhalogen (tri, tetra, etc.)








(i) Alkyl halides or haloalkanes ($R-X$) $\rightarrow$ They form homologous series of general formula $C_nH_{2n+1}X$. They are further classified into primary, secondary, and tertiary.

(ii) Allylic halides $\rightarrow$ Compounds containing halogen atom bonded to an allylic carbon

(iii) Benzylic halides $\rightarrow$ Compounds containing halogen atom bonded to an sp^3 hybridised carbon atom next to an aromatic ring


Methods of preparation:

- From alcohols –




- **From hydrocarbons –**


By free radical halogenations- Yields a complex mixture of isomeric mono- and polyhaloalkanes



- By electrophilic substitution




- **Sandmeyer's reaction –**



- **From alkenes –**


Addition of hydrogen halides to unsaturated hydrocarbons (Markovnikov's rule)



- Addition of halogens


(Method used for detecting double bond in a molecule)

In this method, a reddish-brown colour is discharged.




- **Halogen exchange –**

Finkelstein reaction




- **Swarts reaction (synthesis of alkyl fluoride) –**




Physical properties:

- Melting and boiling points –
- Halides have higher boiling points than hydrocarbons of comparable molecular mass because of having stronger dipole–dipole and van der Waals' forces of attraction.
- The order of increasing boiling points of the different haloalkanes is:



- The boiling points of isomeric haloalkanes decrease with increase in branching. The order of branching is



Hence, order of boiling points is opposite.

- Density –


The density of halides increases with increase in the number of carbon atoms, halogen atoms and atomic mass of the halogen atoms.

- Solubility –

Soluble in organic solvents, but only slightly soluble in water


Reactions of haloalkanes –

- Nucleophilic substitution reaction:**




Mechanism

- Substitution nucleophilic bimolecular ($\text{S}_{\text{N}}2$) (Inversion of configuration)




The increasing order of reactivity is




(Due to increase in hindrance by bulky substituent in the case of 2° and 3° halides)

- Substitution nucleophilic unimolecular (S_N1) (Two-step mechanism)


Step I is the slowest; it is reversible and rate-determining step




The increasing order of reactivity




- In both the mechanisms, for a given alkyl group, the increasing order of reactivity is



- Elimination reactions (β – elimination):**




- Reaction with metals:**
- Grignard reagent –




- Wurtz reaction –

(Preparation of hydrocarbons containing double the number of carbon atoms)




Nucleophilic substitution reaction:

- Aryl halides are less reactive towards nucleophilic substitution due to
 - i. Resonance effect
 - ii. sp^2 hybridisation
 - iii. Instability of phenyl cation
 - iv. Repulsion
- Replacement by alkyl group




Presence of electron-withdrawing group at *o*- and *m*-positions increases the reactivity of haloarenes.




- Electrophilic substitution reaction:**

The electron density at ortho-and para-position is more than at meta-position


Halogenation





Nitration



Sulphonation




Friedel-Crafts reaction




- Reaction with metals:**

Wurtz-Fitting reaction




Fitting reaction




Polyhalogen compounds:

- **Dichloromethane (CH₂Cl₂)** – Industrially used as a solvent
- **Trichloromethane (Chloroform, CHCl₃)** –
 - Used as a solvent for fats, alkaloids, iodine
 - Causes damage to liver, kidneys and skin
 - Stored in closed, dark-coloured container as it is oxidised to an extremely poisonous gas, phosgene, when exposed to light.



- **Triiodomethane (iodoform, CHI₃)** –
 - Earlier, used as an antiseptic
 - Its antiseptic properties are due to the liberation of free iodine
- **Tetrachloromethane** –
 - Used in the manufacture of refrigerants, chlorofluorocarbons
 - Causes liver cancer in humans
 - Causes depletion of ozone layer
- **Freons** – Collective chlorofluorocarbon compounds of methane and ethane
 - Stable, unreactive, non-toxic, non-corrosive and easily liquefiable gas
 - Used as refrigerants
 - Upset the natural ozone balance
- ***p,p'*-Dichlorodiphenyltrichloroethane (DDT)** –



The first chlorinated organic insecticide