

Matrices

- A **matrix** is an ordered rectangular array of numbers or functions. The numbers or functions are called the **elements** or the **entries** of the matrix.

For example, $\begin{bmatrix} -10 & \sin x & \log x \\ e^x & 2 & -9 \end{bmatrix}$ is a matrix having 6 elements. In this matrix, number of rows = 2 and number of columns = 3

- A matrix having m rows and n columns is called a matrix of order $m \times n$. In such a matrix, there are mn numbers of elements.

For example, the order of the matrix $\begin{bmatrix} \sin x & \cos x \\ -1 & 1 + \sin x \\ 0 & \cos x \end{bmatrix}$ is 3×2 as the numbers of rows and columns of this matrix are 3 and 2 respectively.

- A matrix A is said to be a **row matrix**, if it has only one row. In general, $A = [a_{ij}]_{1 \times n}$ is a row matrix of order $1 \times n$.

For example, $[-9 \ 6 \ 5 \ e \ \sin x]$ is a row matrix of order 1×5 .

- A matrix B is said to be a **column matrix**, if it has only one column. In general, $B = [b_{ij}]_{m \times 1}$ is a column matrix of order $m \times 1$.

For example, $B = \begin{bmatrix} -6 \\ 19 \\ 13 \end{bmatrix}$ is a column matrix of order 3×1 .

- A matrix C is said to be a **square matrix**, if the number of rows and columns of the matrix are equal. In general, $C = [c_{ij}]_{m \times n}$ is a square matrix, if $m = n$

For example, $C = \begin{bmatrix} -1 & 9 \\ 5 & 1 \end{bmatrix}$ is a square matrix.

- A square matrix A is said to be a **diagonal matrix**, if all its non-diagonal elements are zero. In general, $A = [a_{ij}]_{m \times n}$ is a diagonal matrix, if $a_{ij} = 0$ for $i \neq j$
- A matrix is said to be a **rectangular matrix**, if the number of rows is not equal to the number of columns.

For example: $\begin{bmatrix} 8 & 3 & 9 \\ 1 & 6 & 7 \end{bmatrix}$ is a rectangular matrix.

- If A is a matrix of order $m \times n$, then the matrix obtained by interchanging the rows and columns of A is called the transpose of matrix A . The transpose of A is denoted by A' or A^T . In other words, if $A = [a_{ij}]_{m \times n}$, then $A' = [a_{ij}]_{n \times m}$

For example, the transpose of the matrix $\begin{bmatrix} 2 & 8 & -3 \\ 1 & 11 & 9 \end{bmatrix}$ is $\begin{bmatrix} 2 & 1 \\ 8 & 11 \\ -3 & 9 \end{bmatrix}$.

- For any matrices A and B of suitable orders, the properties of transpose of matrices are given as:
 - $(A')' = A$
 - $(kA)' = kA'$, where k is a constant
 - $(A + B)' = A' + B'$
 - $(AB)' = B' A'$
- Two matrices $A = [a_{ij}]$ and $B = [b_{ij}]$ are said to be equal (denoted as $A = B$) if they are of the same order and each element of A is equal to the corresponding element of B i.e., $a_{ij} = b_{ij}$ for all i and j .

For example: $\begin{bmatrix} 15 & 11 \\ 7 & 2 \end{bmatrix}$ and $\begin{bmatrix} 15 & 11 \\ 7 & 2 \end{bmatrix}$ are equal but $\begin{bmatrix} 15 & 11 \\ 7 & 2 \end{bmatrix}$ and $\begin{bmatrix} 7 & 2 \\ 15 & 11 \end{bmatrix}$ are not equal.

Example: If $\begin{bmatrix} 7 & x-y \\ 13 & 3y+z \end{bmatrix} = \begin{bmatrix} 2x+y & 5 \\ 2x+y+z & 3 \end{bmatrix}$, then find the values of x , y and z .

Solution:

Since the corresponding elements of equal matrices are equal,

$$2x + y = 7 \dots (1)$$

$$x - y = 5 \dots (2)$$

$$2x + y + z = 13 \dots (3)$$

$$3y + z = 3 \dots (4)$$

On solving equations (1) and (2), we obtain $x = 4$ and $y = -1$.

On substituting the value of y in equation (4), we obtain $z = 6$.

Thus, the values of x , y and z are 4, -1 and 6 respectively.

- Two matrices $A = [a_{ij}]$ and $B = [b_{ij}]$ can be added, if they are of the same order.

The sum of two matrices A and B of same order $m \times n$ is defined as matrix $C = [c_{ij}]_{m \times n}$, where $c_{ij} = a_{ij} + b_{ij}$ for all possible values of i and j .

- The difference of two matrices A and B is defined, if and only if they are of same order. The difference of the matrices A and B is defined as $A - B = A + (-1)B$
- If A , B , and C are three matrices of same order, then they follow the following properties related to addition:
 - Commutative law: $A + B = B + A$
 - Associative law: $A + (B + C) = (A + B) + C$
 - Existence of additive identity: For every matrix A , there exists a matrix O such that $A + O = O + A = A$. In this case, O is called the additive identity for matrix addition.
 - Existence of additive inverse: For every matrix A , there exists a matrix $(-A)$ such that $A + (-A) = (-A) + A = O$. In this case, $(-A)$ is called the additive inverse or the negative of A .

Example: Find the value of x and y , if:

$$\begin{bmatrix} 2x+3y & 9 \\ -2 & 4x-7y \end{bmatrix} + 2 \begin{bmatrix} 3x+\frac{5}{2}y & -11 \\ -13 & 3x-\frac{3}{2}y \end{bmatrix} = \begin{bmatrix} 56 & -13 \\ -28 & 30 \end{bmatrix}$$

Solution:

$$\begin{aligned} & \begin{bmatrix} 2x+3y & 9 \\ -2 & 4x-7y \end{bmatrix} + 2 \begin{bmatrix} 3x+\frac{5}{2}y & -11 \\ -13 & 3x-\frac{3}{2}y \end{bmatrix} = \begin{bmatrix} 56 & -13 \\ -28 & 30 \end{bmatrix} \\ \Rightarrow & \begin{bmatrix} 2x+3y & 9 \\ -2 & 4x-7y \end{bmatrix} + \begin{bmatrix} 6x+5y & -22 \\ -26 & 6x-3y \end{bmatrix} = \begin{bmatrix} 56 & -13 \\ -28 & 30 \end{bmatrix} \\ \Rightarrow & \begin{bmatrix} 8(x+y) & -13 \\ -28 & 10(x-y) \end{bmatrix} = \begin{bmatrix} 56 & -13 \\ -28 & 30 \end{bmatrix} \end{aligned}$$

Therefore, we have

$$8(x+y) = 56 \text{ and } 10(x-y) = 30$$

$$\Rightarrow x+y = 7 \quad \dots (1)$$

And

$$x-y = 3 \quad \dots (2)$$

Solving equation (1) and (2), we obtain $x = 5$ and $y = 2$

- The multiplication of a matrix A of order $m \times n$ by a scalar k is defined as

$$kA = kA = k[a_{ij}]_{m \times n} = [k(a_{ij})]_{m \times n}$$

- If A and B are matrices of same order and k and l are scalars, then
 - $k(A+B) = kA + kB$
 - $(k+l)A = kA + lA$
- The negative of a matrix B is denoted by $-B$ and is defined as $(-1)B$.

- The product of two matrices A and B is defined, if the number of columns of A is equal to the number of rows of B .
- If $A = [a_{ij}]_{m \times n}$ and $B = [b_{jk}]_{n \times p}$ are two matrices, then their product is defined as $AB = C = [c_{ik}]_{m \times p}$, where $c_{ik} = \sum_{j=1}^n a_{ij}b_{jk}$

For example, if $A = \begin{bmatrix} 2 & -3 & 7 \\ 0 & 1 & -9 \end{bmatrix}$ and $B = \begin{bmatrix} -5 & 9 \\ 7 & 2 \\ 0 & 1 \end{bmatrix}$, then

$$\begin{aligned} AB &= \begin{bmatrix} 2 & -3 & 7 \\ 0 & 1 & -9 \end{bmatrix} \times \begin{bmatrix} -5 & 9 \\ 7 & 2 \\ 0 & 1 \end{bmatrix} \\ &= \begin{bmatrix} 2 \times (-5) + (-3) \times 7 + 7 \times 0 & 2 \times 9 + (-3) \times 2 + 7 \times 1 \\ 0 \times (-5) + 1 \times 7 + (-9) \times 0 & 0 \times 9 + 1 \times 2 + (-9) \times 1 \end{bmatrix} \\ &= \begin{bmatrix} -31 & 19 \\ 7 & -7 \end{bmatrix} \end{aligned}$$

- If A , B , and C are any three matrices, then they follow the following properties related to multiplication:
 - Associative law: $(AB)C = A(BC)$
 - Distribution law: $A(B + C) = AB + AC$ and $(A + B)C = AC + BC$, if both sides of equality are defined.
 - Existence of multiplicative identity: For every square matrix A , there exists an identity matrix I of same order such that $IA = AI = A$. In this case, I is called the multiplicative identity.
- Multiplication of two matrices is not commutative. There are many cases where the product AB of two matrices A and B is defined, but the product BA need not be defined.

For example, if $A = \begin{bmatrix} -1 & 5 \end{bmatrix}_{1 \times 2}$ and $B = \begin{bmatrix} 0 & 1 & -4 \\ 3 & 2 & -1 \end{bmatrix}_{2 \times 3}$, then AB is defined where as BA is not defined.