

Chaper 5. Heat Change Of State

PAGE NO-255:

Solution 1

Specific heat is the amount of heat required to raise the temperature of 1 kg of the substance through 1°C . It is not same as heat capacity.

Solution 2

SI unit of specific heat is $\text{J kg}^{-1}\text{K}^{-1}$.

Solution 3

Specific Heat.

Solution 4

Principle of Calorimetry: When a hot body is mixed or kept in contact with a cold body, there is a transfer of heat from hot body to cold body such that Total heat gained by colder body = Total heat lost by the hot body, if there is no loss of heat to the surroundings.

Solution 5

Thermal capacity of a body is the quantity of heat required to raise its temperature by 1°C .

Solution 6

The product of mass and specific heat is known as heat capacity.

Solution 7

No, specific heat does not depend on temperature. It is constant for a substance.

Solution 8

Copper rod becomes warmer than an aluminium rod of the same mass because copper has lower heat capacity than aluminium.

Solution 9

The amount of heat required to raise the temperature of a body by 1°C is called Heat capacity.

Solution 10

Heat capacity has units $\text{J}/^{\circ}\text{C}$.

Solution 11

Specific heat of water is $4200 \text{ J kg}^{-1}\text{K}^{-1}$

Solution 12

The substances like water which have high heat capacity warm up more slowly than substances like iron which have low heat capacity.

Solution 13

Latent heat is the quantity of heat absorbed or released by a substance undergoing a change of state, such as ice changing to water or water to steam, at constant temperature.

Solution 14

SI unit of latent heat is J/kg .

Solution 15

It means that 1 g of ice at 0°C absorbs 336 J of heat energy to convert into water at 0°C .

Solution 16

When a liquid is solidified, it may either expand or contract. As water freezes to form ice it expands and increases in volume by 10 per cent.

Solution 17

The melting point of ice decreases on addition of impurities in it.

Solution 18

The melting point of substances which contract on melting (like ice) decreases by the increase in pressure.

Solution 19

Regelation is the phenomenon of melting under pressure and freezing again when the pressure is reduced.

Solution 20

The amount of heat required to change a liquid at its boiling point to vapour at the same temperature is called its latent heat of vaporization.

Solution 21

The boiling point of a liquid increases with the increase in pressure and decreases with the decrease in pressure.

Solution 22

The latent heat of fusion of ice is the amount of heat energy required to change ice at 0°C into water at the same temperature.

Solution 23

The latent heat of vaporization of steam is the amount of heat energy required to change water at 100°C to steam at the same temperature.

Solution 24

The physical quantity which does not change during change of state is temperature of the body.

Solution 25

1 kg of ice at 0°C absorbs 336000 J of heat energy to convert into water at 0°C .

Therefore, 1 kg of water at 0°C has 336000 J heat energy more than 1 kg ice at 0°C . So, ice appears colder than water.

Solution 26

Steam causes more severe burns than water at 100°C because every gram of steam gives out 2260 J of heat energy while condensing. This much quantity of heat is additional to the heat contained in 1 g of boiling water.

Solution 27

The unit of heat capacity in CGS system is calorie/°C.

Solution 28

$1 \text{ cal g}^{-1} \text{ }^{\circ}\text{C}^{-1} = 239 \text{ J kg}^{-1} \text{ }^{\circ}\text{C}^{-1}$

Solution 29

This means that 0.2 cal g⁻¹ °C⁻¹ of heat is required to raise the temperature of 1g of the body by 1°C.

Solution 30

$m = 100 \text{ g}$
 $C = 0.04 \text{ cal g}^{-1} \text{ }^{\circ}\text{C}^{-1}$
Heat capacity = $m \times C = 4 \text{ cal }^{\circ}\text{C}^{-1}$

Solution 31

Heat capacity = mass x specific heat capacity

Solution 32

No, specific heat does not depend on mass of a substance. It is constant for a substance.

Solution 33

Specific heat of water in SI units is 4200 J kg⁻¹ K⁻¹.

Solution 34
Ammonia has the maximum value of specific heat.

Solution 35

Heat gained or lost by a substance depends on the mass of the substance and the nature of the substance.

Solution 36

Oceans cover more than 70% of Earth's surface, making them the world's largest solar collectors. The sun's heat warms the surface water a lot more than the deep ocean water, and this temperature difference creates heat energy. Thus, oceans are known as storehouse of heat energy. Just a small portion of the heat trapped in the ocean could power the world.

Solution 37

Water has a high specific heat capacity. So, water extracts much heat without much rise in temperature. By allowing water to flow in radiator pipes of the vehicles, heat energy from such parts is removed. Hence, it is used as a cooling agent in the radiators of automobiles.

PAGE NO-256:**Solution 38**

A change of state is a change in the object from a solid to liquid or from a solid to gas or from liquid to solid. Ice changing into water is an example of a change in state.

Solution 39

Heat energy is absorbed during melting of ice.

Solution 40

Heat energy is released during freezing of ice.

Solution 41

Temperature remains constant when ice melts at 0°C.

Solution 42

The molecules in a solid are held by strong intermolecular bonds. For the solid to melt, these bonds have to be broken. Since energy is needed to break the intermolecular bonds, the thermal energy supplied at the melting point is used to do the work to break the intermolecular bonds between the molecules of the solid. Once the intermolecular bonds are broken, the molecules can then move out of their fixed positions. Hence it can then be said that the solid has melted, which is the change of state from solid to liquid. This explains why temperature remains constant during the melting phases.

Solution 43

Whenever a substance goes through a phase change (like boiling), the energy goes into breaking up the interactions between molecules, and so the temperature stays constant until all the interactions are broken. Once whole of the substance has boiled, then any added heat will act to raise their temperature again.

Solution 44

Atmosphere is usually warm during snowfall because each kilogram of ice on melting absorbs 336000 J of heat from atmosphere.

Solution 45

Latent heat of fusion of ice, $L=336 \text{ J/g}$ Mass of ice, $m=2\text{g}$ Amount of heat required convert 2g of ice at 0°C into water at $0^\circ\text{C} = m \times L = 2 \times 336 = 672 \text{ J}$

Solution 46

Latent heat of vapourisation of water, $L=540 \text{ cal/g}$ Mass of water, $m=100\text{g}$ Amount of heat required convert 100g of water at 100°C into steam at $100^\circ\text{C} = m \times L = 100 \times 540 = 54 \text{ kJ}$

Solution 47

Ice melts under pressure. So, when the steel blades of the skates pressed on the ice, the ice melts. The water formed makes the skates slide easily over the ice, reducing friction. So, when we are skating on ice, we are skating on a thin film of water, which acts like lubricating oil. Nothing such happens in case of glass.

Solution 48

Bottled drinks are cooled more effectively when surrounded by lumps of ice than by cold water at 0°C because ice appears colder than water at 0°C .

Solution 49

Parts AB and CD correspond to the substance existing in two states.

Solution 50

(e) The first time when temperature is constant represents change of state from solid to liquid and the second time temperature is constant represents change of state from liquid to vapour.


Solution 51

(a) Boiling point of substance is 150°C (because the part BC represents condensation where the vapour changes into liquid without the change in temperature).

(b) DE represents freezing of the substance where the liquid changes into solid at a constant temperature of 100°C .

(c) Melting point is the temperature of the region DE where liquid changes into solid i.e., 100°C .

Solution 52



Solution 53

i. AB represents the change of state from solid to liquid i.e., AB represents melting of ice at 0°C. ii. CD represents the change of state from liquid to vapour i.e., CD represents boiling of water at 100°C. iii. The ice initially is in solid state at -10°C. On heating, its temperature rises to 0°C. It then takes some heat at 0°C to melt in water at 0°C which is its latent heat.